

PROBING THE INNER KPC OF MASSIVE GALAXIES WITH STRONG GRAVITATIONAL LENSING

YASHAR D. HEZAVEH, PHILIP J. MARSHALL, ROGER D. BLANDFORD
 Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, Stanford, CA, USA
Draft version October 25, 2014

ABSTRACT

We examine the prospects of detecting demagnified images of gravitational lenses in observations of strongly lensed mm-wave molecular emission lines with ALMA. We model the lensing galaxies as a superposition of a dark matter component, a stellar population, and a central supermassive black hole and forecast the detection of the central images for a range of relevant parameters (e.g. stellar core and black hole mass). We find that over a large range of acceptable parameters, future deep observations of lensed molecular lines with ALMA will be able to detect the central images at $\gtrsim 3\sigma$ significance. We use Fisher analysis to examine the constraints that could be placed on these parameters in various scenarios.

Subject headings: black hole physics — gravitational lensing: strong — galaxies: formation — galaxies: high-redshift

1. INTRODUCTION

Probing the inner 0.5 kpc is interesting because:

- 1) SMBHs are there
- 2) Various DM models have different predictions
- 3) The stellar population profile contains information about past mergers and BH-stellar population interactions.

Mapping the matter density and decomposing these different components can thus shed light on various astrophysical phenomena.

Stellar light profiles:

- 1) Massive galaxies often exhibit cored surface brightness profiles.
- 2) The most massive ellipticals are thought to form through gas-poor mergers.
- 3) The central structure of a merger without gas is dominated by the more concentrated of the two progenitors: the steeper central density cusp survives.
- 4) Since high-mass ellipticals are thought to be built from mergers of lower-mass ellipticals, which have steep central-density cusps, the cores in more massive ellipticals have to be the result of another (not merger) physical mechanism.
- 5) The existence of cores thus, represents a challenge to our understanding of the merging process.
- 6) “Black hole scouring” has been suggested to produce a core.

Explain “Black hole scouring”.

...
 ...
 ...
 ... whatever the mechanisms are, clear measurements of the central densities can be valuable to solve this

puzzle and to shed light on the connection of SMBHs and galaxies.

Strong lensing:

- 1) lensing is a well-known tool for mapping out mass
- 2) strong lenses have been used to measure the einstein mass (e.g., X et al.), and the density skopes (Y et al.).
- 3) Strong lensing formalism, predicts an odd number of images, with a central image that is *demagnified*
- 4) it has been known that the flux (i.e. demagnification) of the central image is very sensitive to the very inner regions of the density profiles: very peaky, singular density profiles, demagnify the image by very large amounts, whereas cored or shallow profile will result in bright central images.
- 4) typically only an even number of images are observed, suggesting that the central images are demagnified below the sensitivity of instruments.
- 5) Additionally, another reason for lack of detection is that the central image is located at the center of the lensing galaxy, where i) the sensitivity is lower, due to photon noise of the lens ii) it is difficult to distinguish the flux of the central image from the emission from the lens. iii) if in optical, since they pass through the centers of galaxies there’s a high chance of large absorption (in the lens)

a review of central image observations, claims, and studies:

Winn et al 2004
 Rusin et al 2005
 Inada et al 2005, etc.

Describe the newly discovered population of mm lenses + ALMA's observations:

- 1) strong lenses in mm were discovered (SPT, Herschel, etc.)
- 2) ALMA observations confirmed that they're all lensed and that the sources are at high z
- 3) because they were selected by flux, + the high sensitivity of ALMA \rightarrow SNR is very high
- 4) the sources have many molecular lines. If a central image of a *molecular line* is observed it will be easily identifiable since it corresponds to the redshift of the source, and there will be no confusion that it may be associated with the lens.
- 5) since these lines are in mm, there's very little (if any) absorption in the lens, so the flux doesn't decrease due to absorption

motivation and description of the paper:

- 1) Long ALMA observations of these molecular lines are likely to be carried out for various reasons (e.g. power spectrum of dark matter, measuring the mass of background black holes).
- 2) In this paper we explore the possibility of detection of these central images in such deep observations, and investigate what we could learn about the innermost regions of galaxies from detection or non-detection of such central images.
- 3) the paper is organized as: section 2 describes the simulations, section 3 presents the results and discuss them and conclude in section 5
- 4) we use Λ CDM cosmology of XXX

2. SIMULATIONS

description We generate mock data cubes of gas emission in the source galaxy, lens the data cubes with a foreground halo, predict the ALMA visibilities, and use the visibilities to estimate the detection significance of the central image for various parameters. The gas in the source galaxy is modeled as a collection of 5 Gaussian clumps with FWHM of XX, distributed over XX kpc and each separated in a different channel. and an intrinsic velocity dispersion of 10 km/s (FWHM) similar to observations of ?. The choice of two discrete rings is made to enforce resolving the inner 50 pc region for a black hole mass measurement. We point out, however, that assuming a more general case (e.g. an exponential disk) does not change the results (see Figure 2 of ?). The velocity integrated flux of the circumnuclear ring is set to 4 mJy km/s . This value is calculated by scaling the flux of the circumnuclear CO gas in Arp 220 to to $z=2$ (?). The rings are concentric and in circular orbits. The rotational velocity of each point in the rings is calculated based on the enclosed mass contributed from the galaxy mass model

and the mass of the central black hole. The black hole is given a mass of $M_{\text{BH}} = 5 \times 10^8 M_{\odot}$. We model the galaxy density with an isothermal profile ($\rho \propto r^{-2}$) with a central velocity dispersion of 200 km/s. The line-of-sight velocities are calculated to generate a frequency-position data cube. Figure ?? shows a velocity-position diagram extracted from the data cube. The data comprises 100 frequency channels covering 800 km/s.

Each layer of the data cube is lensed with the foreground halo to generate a lensed cube. We use a Singular Isothermal Ellipsoid (SIE) mass model for the lens galaxy ($\rho \propto r^{-2}$, ?), placed at $z_l = 0.5$ and place the source at $z_s = 2.0$. The central velocity dispersion of the lens is set to 180 km/s , which is typical for galaxy-galaxy lensing systems (e.g. ?).

To predict the visibilities we calculate the ALMA *uv*-coverage for a 5-hr long observation with the most extended antenna configuration (full array), using the *simobserve* task of Common Astronomy Software Applications package, which results in an angular resolution of $\sim 20 \text{ milli-arcsec}$ at an observing frequency of 240 GHz. The visibilities for each channel are calculated by computing the 2D Fourier transform of the corresponding layer of the data cube and resampling the Fourier transform maps over the *uv*-coverage. The noise is estimated using ALMA sensitivity calculator for a channel width of 8 km/s at 240 GHz. We use finite differencing of visibilities to calculate the Fisher information and the parameter covariance matrix to calculate the significance of black hole mass measurement.

When the background source only consists of the two rings described above, the lens model parameters have large uncertainties and are mildly degenerate with the mass of the black hole. A realistic source galaxy, however, will consist of many other components, extended over kilo-parsec scales, which if included in lens models, can strongly constrain the lens parameters. We run five simulations in which the background source is embedded in an extended Gaussian clump with a radius of 1 kpc and a flux of 1 Jy km/s (typical for SMGs, see ?). We find that in these simulations, the lens parameters are extremely constrained and the effect of marginalization over them barely reduces the significance of black hole mass measurements ($\lesssim 0.05 \sigma$). This suggests that when the full structure of the source galaxies are included in models, the lens parameters do not significantly contribute to the black hole mass uncertainties. Therefore, to reduce the computational cost of simulations, we do not marginalize over the lens parameters.

3. RESULTS AND DISCUSSIONS

3.1. Detection of central images

- 1) We evaluate the detection significance of central images over a range of relevant parameters.
- 2) Figure 4 shows our results for a 10-hr long observation with ALMA
- 3) we show that, as expected, the image is fainter for steeper profile slopes.
- 4) the 0.9 value (combined with 0.1 DM) results in the isothermal profile, which has been shown to be consistent with strong lensing data
- 5) if that's the case, and in the absence of massive BHs, then stellar cores as small as ~ 80 pc should allow a 5σ detection of the central images.
- 5A) we note that for much shallower density profiles, the central image is actually *magnified*, and therefore it's detection should be trivial.
- 6) the flux is significantly suppressed by the central SMBH
- 7) the dashed curves show the same for a galaxy with a $2e8 M_{\odot}$ SMBH.
- 8) we note that although for some parameter combinations (e.g., $\gamma_s = 1.2$, $M_{BH} = 2e8$, $R_{core} < 300 pc$) there is no chance of detecting the central image, over a significant fraction of allowed parameters the central image could be detected.
- 9) to calculate this plot we have used a configuration XX, because: it allows us to resolve the central image (if existent), while it is more sensitive than more extended configurations. In addition, this is the configuration that will be likely used for other projects/sciences.

with the core size: a larger core, results in a brighter image, which can be suppressed by a more massive BH.

- 6) The Fisher data show that the black hole mass cannot be measured with high significance in such observations (1.16σ)
- 7) however, interestingly, the degeneracy of the black hole mass and the size of the central core breaks at the high end. and the stellar core is measured with XX σ
- 8) this is likely because a very large core results in a fairly extended central images, which could be *resolved* (see Figure 1), in that case, the flux suppression by a crazy massive black hole will result in a distinct dip in the central image which can be used to distinguish the two.
- 9) we note that the constraints in this case will be of similar strength to those of Winn 2004.
- 10) in the case of very small core, and a massive black hole (no detection of the image) we only get an upper limits of \sim and \sim for the two.
- 11) and in case with massive core and massive black hole we measure both with high significance. 12) we note that all these parameter combinations are plausible choices, and therefore long observations such as these will be able to either measure these parameters with confidence or place strong constraints on them.

3.2. SMBHs and the core size

- 1) in this section we study how well we can constrain the mass of the central black hole, the slope of the density and the size of the stellar core.
- 2) we choose a few models where the central image is detected OR not detected and use a Fisher analysis to determine the degeneracies between the parameters and the detection significance of each parameter.
- 3) we note that in the Fisher analysis we include all nuisance parameters (e.g. source position, size, etc.) and marginalize over them
- 4) Figure 4 shows the parameters for a case with $R_{core}=180$ pc, $M_{BH} = 1e8$ and $\gamma = 1$. As seen from Figure 3, this parameter combination results in a faint ($\sim 3\sigma$) detection of the central image.
- 5) we note that the black hole mass is degenerate




FIG. 1.— illustration of the central image in 2 cases:




FIG. 2.— Model density profiles. The black dashed line shows the dark matter component, with a slope of 0.1, consistent with a projected NFW at the innermost regions. the grey dashed curves show cored-sersic five fits to stellar light profiles of massive galaxies from Ferrarese 2006. The red solid curve shows our cored-power-law model that we use to approximate this stellar component. The solid black line shows the total matter density (dark plus stellar) of our model.




FIG. 3.— Significance of detection of the central image as a function of the stellar core size for a 10-hr long ALMA observation. The colors correspond to different slopes of the stellar component. The solid curves correspond to a case without a SMBH while the dashed line show the result of a simulation which includes a $2 \times 10^8 M_{\odot}$ SMBH at its center.

4. CONCLUSION




FIG. 4.— Covariance matrix of parameters for a few different scenarios.