

Inline and Display Mathematics

Source-backed grouped formula sample

1 Expressions

Expression 1. The following expression is taken from a source-backed formula corpus.

$$I_{\epsilon}^{(1)}(a^2) = \frac{\pi}{2} \frac{1}{\cos \frac{\pi}{2} \epsilon} \left\{ \frac{1}{a^{1+\epsilon}} + 2 \sum_{n=1}^{\infty} \frac{1}{(2\pi n)^{1+\epsilon}} + 2 \sum_{n=1}^{\infty} \frac{1}{(2\pi n)^{1+\epsilon}} \left[\frac{1}{\left(1 + \frac{a^2}{4\pi^2 n^2}\right)^{\frac{1+\epsilon}{2}}} - 1 \right] \right\}.$$

Expression 2. The following expression is taken from a source-backed formula corpus.

$$\left(\frac{\det'(\Delta_v)}{\det(\Delta_v^{(0)})} \right)^{-\frac{1}{2}} \left(\frac{\det(\Delta_A)}{\det(\Delta_A^{(0)})} \right)^{-1} \left(\frac{\det(\Delta_c)}{\det(\Delta_c^{(0)})} \right)^{+1} = \left(\frac{\det'(\Delta_-)}{\det(\Delta^{(0)})} \right)^{-1}$$

Expression 3. The following expression is taken from a source-backed formula corpus.

$$P^- = \frac{1}{2} \int dx^- d^2 x^\perp : \left[-\phi \partial^{\perp 2} \phi + \sum_{i=1}^2 \left(\bar{\psi}_m^{(i)} \frac{-\partial^{\perp 2} + m^2}{i\partial^+} \gamma^+ \psi_m^{(i)} + 2g\phi \bar{\psi}_m^{(i)} \psi_m^{(i)} + g^2 \bar{\psi}_m^{(i)} \phi \frac{\gamma^+}{i\partial^+} \phi \psi_m^{(i)} \right) \right] :.$$

Expression 4. The following expression is taken from a source-backed formula corpus.

$$= \int dt d\theta d\phi dt' d\theta' d\phi' \frac{\sin \theta \sin \theta' \Phi_-(t, \theta, \phi) \Phi_-(t', \theta', \phi')}{[\cosh(\frac{t-t'}{a}) - (\sin \theta \sin \theta' \cos(\phi - \phi') + \cos \theta \cos \theta')]^h}.$$

Expression 5. The following expression is taken from a source-backed formula corpus.

$$8f_4^{(8)} \phi^i \phi^j \phi^k \phi^l v^p \left[\delta^{si} (\theta \gamma^{pm} \theta) (\theta \gamma^{mj} \theta) (\theta \gamma^{kn} \theta) (\theta \gamma^{ln} \theta) + \delta^{sp} (\theta \gamma^{mi} \theta) (\theta \gamma^{mj} \theta) (\theta \gamma^{kn} \theta) (\theta \gamma^{ln} \theta) \right],$$