

Inline and Display Mathematics

Source-backed grouped formula sample

1 Expressions

Expression 1. The following expression is taken from a source-backed formula corpus.

$$\frac{G'(\frac{1}{2} + \frac{1}{\sqrt{2}} - i\beta, \frac{1}{2} - \frac{1}{\sqrt{2}} - i\beta; \frac{iz_0}{\hbar c})}{G(\frac{1}{2} + \frac{1}{\sqrt{2}} - i\beta, \frac{1}{2} - \frac{1}{\sqrt{2}} - i\beta; \frac{iz_0}{\hbar c})} = -\frac{i}{2\hbar c} - (\frac{1}{2} + i\beta)\frac{1}{z_0}.$$

Expression 2. The following expression is taken from a source-backed formula corpus.

$$\frac{-i\langle Ta_{\mathbf{k}}^{\dagger t}(-\mathbf{q})a_{\mathbf{k}'}^{\dagger}(\mathbf{q}')\rangle}{\langle T1\rangle} = \sum_n \exp(\omega_n t) \frac{-i\langle Ta_{\mathbf{k}}^{\dagger n}(-\mathbf{q})a_{\mathbf{k}'}^{\dagger}(\mathbf{q}')\rangle}{\langle T1\rangle}$$

Expression 3. The following expression is taken from a source-backed formula corpus.

$$S = -\frac{1}{2} \int d^2x dy \left(i\bar{\psi}\gamma^\mu \frac{\partial\psi}{\partial x^\mu} - \frac{1}{2}\bar{\psi}\gamma_5 \frac{\partial\psi}{\partial y} + \frac{\partial\phi}{\partial x^\mu} \frac{\partial\phi}{\partial x_\mu} - \frac{1}{4} \frac{\partial\phi}{\partial y} \frac{\partial\phi}{\partial y} + F^2 \right)$$

Expression 4. The following expression is taken from a source-backed formula corpus.

$$\frac{\partial}{\partial q_{tA}} = \frac{\partial r}{\partial q_{tA}} \frac{\partial}{\partial r} + \frac{\partial e_B}{\partial q_{tA}} \frac{\partial}{\partial e_B} + \frac{\partial E_s}{\partial q_{tA}} \frac{\partial}{\partial E_s} + \frac{\partial y_{sB}}{\partial q_{tA}} \frac{\partial}{\partial y_{sB}}.$$

Expression 5. The following expression is taken from a source-backed formula corpus.

$$\mathcal{H}^{(0)} = -\frac{2\kappa}{\sqrt{-\gamma}} \frac{\delta I^{(0)}}{\delta \gamma_{\mu\nu}} \frac{\delta I^{(0)}}{\delta \gamma_{\lambda\rho}} \left(\gamma_{\mu\rho} \gamma_{\nu\lambda} - \frac{1}{d-1} \gamma_{\mu\nu} \gamma_{\lambda\rho} \right) + \frac{\sqrt{-\gamma}}{\kappa} \Lambda = 0,$$