

Inline and Display Mathematics

Source-backed grouped formula sample

1 Expressions

Expression 1. The following expression is taken from a source-backed formula corpus.

$$\bar{G}^{(\zeta)}(\beta) = \frac{(\beta + i\pi) \sinh\left(\frac{\pi\beta}{2\pi + \zeta}\right)}{\Gamma\left(1 - \frac{\pi - i\beta}{2\pi + \zeta}\right) \Gamma\left(1 - \frac{\pi + i\beta}{2\pi + \zeta}\right)},$$

Expression 2. The following expression is taken from a source-backed formula corpus.

$$\frac{\partial V}{\partial \Lambda} = \frac{1}{\Lambda^2} \frac{\partial V}{\partial \psi_{\alpha a}} C'(0) \frac{\partial V}{\partial \bar{\psi}_a^\alpha} - \tau \Lambda^{D-2} \frac{\partial^2 V}{\partial \psi_{\alpha a} \partial \bar{\psi}_a^\alpha},$$

Expression 3. The following expression is taken from a source-backed formula corpus.

$$f_3(r) = 2\sqrt{\frac{\bar{s}^2 - Qc}{Qc}} e^{-\tilde{c}\sqrt{\bar{s}^2 - Qc}} \frac{\left(1 - \frac{r_0^2}{r^2}\right)^{(1+\sqrt{(\bar{s}^2 - Qc)/D_2})/2}}{1 - e^{-2\tilde{c}\sqrt{\bar{s}^2 - Qc}} \left(1 - \frac{r_0^2}{r^2}\right)^{\sqrt{(\bar{s}^2 - Qc)/D_2}}}.$$

Expression 4. The following expression is taken from a source-backed formula corpus.

$$\frac{\partial \zeta^{(\lambda\mu\alpha)}}{\partial \zeta^{(\tau,\nu,\beta)}} = \psi_{(\tau,\nu,\beta)}^{(\lambda\mu\alpha)} = \frac{1}{2} \left(D_{(\tau,\nu,\beta)}^{(\lambda\mu\alpha)} + \omega_{(\tau,\nu,\beta)}^{(\rho,\omega,\gamma)} D_{(\rho,\omega,\gamma)}^{(\lambda\mu\alpha)} \right).$$

Expression 5. The following expression is taken from a source-backed formula corpus.

$$\sup_{\mathcal{T} \geq 1} \left| \frac{1}{\mathcal{T}} \int_0^{\mathcal{T}} \tilde{G}(\mu^{N,M}(t), z^{N,M}(t)) dt - \frac{1}{\mathcal{T}} \int_0^{\mathcal{T}} \tilde{G}(\bar{\mu}^{N,M}(t), \bar{z}^{N,M}(t)) dt \right| \leq \kappa_3(\epsilon), \quad \lim_{\epsilon \rightarrow 0} \kappa_3(\epsilon) = 0.$$