

Inline and Display Mathematics

Source-backed grouped formula sample

1 Expressions

Expression 1. The following expression is taken from a source-backed formula corpus.

$$\leq \frac{1}{\mathcal{T}} \sum_{l=0}^{\lfloor \frac{\mathcal{T}}{\Delta(\epsilon)} \rfloor} \int_{t_l}^{t_{l+1}} |\tilde{G}(\mu^{N,M}(t), z^{N,M}(t)) - \tilde{G}(\mu^{N,M}(t_l), z^{N,M}(t_l))| dt + L\kappa_2(\epsilon) + 2M\Delta(\epsilon) \quad \forall \mathcal{T} \geq 1,$$

Expression 2. The following expression is taken from a source-backed formula corpus.

$$\nu^{-1}(\tilde{\varepsilon}) = 4\tilde{\varepsilon}^{-1} - 8\frac{n-4}{n-2}\tilde{\varepsilon}^{-2} + 16\frac{n-4}{n-2}\tilde{\varepsilon}^{-3} - 32\left\{[52+108\zeta(3)] - [16+36\zeta(3)]n + n^2\right\} \frac{\tilde{\varepsilon}^{-4}}{(n-2)^3} + \dots$$

Expression 3. The following expression is taken from a source-backed formula corpus.

$$\Gamma_{1,fermions}[B] = \frac{1}{2} \int_0^\infty \frac{dT}{T} \int_{x(0)=x(T)} Dx(t) \exp(-\frac{1}{4} \int_0^T dt \dot{x}^2) Tr_L \Phi^{[1/2]}[\dot{x}] Tr_c P \exp(ig \int_0^T dt \dot{x} \cdot B)$$

Expression 4. The following expression is taken from a source-backed formula corpus.

$$\zeta_{P\bar{K}}(s) = \zeta_{\bar{\mathbb{V}}}(s) + \frac{\Gamma(s+\frac{1}{2})}{\sqrt{\pi}\Gamma(s)} \left[\frac{2}{3^{s+\frac{1}{2}}} {}_2F_1[\frac{1}{2}, s+\frac{1}{2}, \frac{3}{2}, -\frac{1}{3}] - \frac{1}{4^s} \frac{1}{s} \right]$$

Expression 5. The following expression is taken from a source-backed formula corpus.

$$\left(\frac{1}{2}f'+e^f\frac{1}{12\sqrt{2}}\bar{g}\left(2e^{-\varphi}+e^{2\varphi}\right)\right)\gamma_0\varepsilon_a=-\frac{1}{6}\sqrt{\frac{1}{2}}e^fe^\varphi\frac{1}{2\bar{g}}\gamma^{\nu\rho}R_{\nu\rho}{}^{ij}\gamma_{ij}\gamma_0\varepsilon_a.$$