

Inline and Display Mathematics

Source-backed grouped formula sample

1 Expressions

Expression 1. The following expression is taken from a source-backed formula corpus.

$$\delta^{ab} \delta_{xy} = \langle \frac{\delta S}{\delta \bar{c}_x^a} \bar{c}_y^b \rangle = \tilde{Z}_3 \langle i(\partial D_r c)_x^a \bar{c}_y^b \rangle - \frac{Z_\lambda \lambda}{Z_3} \langle (s_r B_x^a) \bar{c}_y^b \rangle .$$

Expression 2. The following expression is taken from a source-backed formula corpus.

$$G_D^{(+)}(\mathbf{r}, \mathbf{r}'; k) = \mathcal{G}_D^{(+)}(\mathbf{R}; k) \stackrel{(r \rightarrow \infty)}{\sim} -\frac{1}{4\pi} \left(\frac{k}{2\pi} \right)^{(D-3)/2} e^{i\gamma_D} \frac{e^{ikr}}{r^{(D-1)/2}} e^{-i\mathbf{k}' \cdot \mathbf{r}'} ,$$

Expression 3. The following expression is taken from a source-backed formula corpus.

$$N_Q(-i\Omega_m, \vec{p}) = -\frac{d_Q(R)}{2} \int \frac{d^3l}{(2\pi)^3} T \sum_{n=-\infty}^{\infty} \text{tr} S_Q(-i\omega_n + \mu_Q, \vec{l}) S_Q(-i\omega_n - i\Omega_m + \mu_Q, \vec{l} + \vec{p}).$$

Expression 4. The following expression is taken from a source-backed formula corpus.

$$\Pr\left(|\langle x, \varphi \rangle| \geq \xi\right) = \Pr\left(|Z| \geq \xi / \|\Sigma^{1/2} x\|_2\right) \geq \Pr\left(|Z| \geq \xi / \sigma_{\min}\right) \geq \frac{\sigma_{\min}}{\xi} \cdot \frac{1}{\sqrt{2\pi}} e^{-\xi^2/2\sigma_{\min}^2},$$

Expression 5. The following expression is taken from a source-backed formula corpus.

$$\exp\left(-ie\left[\frac{1}{\sqrt{\pi+gN}}\left(A_\mu^{(1)},\varepsilon_{\nu\mu}\partial_\nu\Phi^{(1)}\right)+\sum_{I=2}^N\frac{1}{\sqrt{\pi}}\left(A_\mu^{(I)},\varepsilon_{\nu\mu}\partial_\nu\Phi^{(I)}\right)\right]\right)\Bigg\rangle_{\{K^{(I)}\}}=$$