

Inline and Display Mathematics

Source-backed grouped formula sample

1 Expressions

Expression 1. The following expression is taken from a source-backed formula corpus.

$$g_{\mathbf{k},\mathbf{q}}(\mathbf{x}) = -e^{-i\mathbf{q}\cdot\mathbf{x}}(\frac{1}{2N\epsilon_{\mathbf{q}}})\Lambda_{\mathbf{k}}(-\mathbf{q})\omega_{\mathbf{k}}(\mathbf{q}) + iU_{\mathbf{q}}(\mathbf{x})\Lambda_{\mathbf{k}}(-\mathbf{q}) = -f_{\mathbf{k},\mathbf{q}}^*(\mathbf{x})$$

Expression 2. The following expression is taken from a source-backed formula corpus.

$$\leq \frac{1}{\mathcal{T}} \sum_{l=0}^{\lfloor \frac{\mathcal{T}}{\Delta(\epsilon)} \rfloor} \int_{t_l}^{t_{l+1}} |\tilde{G}(\mu^{N,M}(t), z^{N,M}(t)) - \tilde{G}(\mu^{N,M}(t_l), z^{N,M}(t_l))| dt + L\kappa_2(\epsilon) + 2M\Delta(\epsilon) \quad \forall \mathcal{T} \geq 1,$$

Expression 3. The following expression is taken from a source-backed formula corpus.

$$\mathcal{L}_G^{(1)}| = \frac{1}{\gamma^2} \frac{\partial_z v_1 \partial_{\bar{z}} v_1 + \partial_z v_2 \partial_{\bar{z}} v_2}{1 + v_1^2 + v_2^2}, \quad \mathcal{L}_G^{(2)}| = \frac{1}{\gamma^2} \frac{\partial_z c \partial_{\bar{z}} c + \partial_z v_0 \partial_{\bar{z}} v_0}{c^2 + v_0^2 - 1}.$$

Expression 4. The following expression is taken from a source-backed formula corpus.

$$\left(I_{0+}^{\alpha,\alpha',\beta,\beta',\gamma}t^{\rho-1}\right)(x)=\frac{\Gamma(\rho)\Gamma(-\alpha'+\beta'+\rho)\Gamma(-\alpha-\alpha'-\beta+\gamma+\rho)}{\Gamma(\beta'+\rho)\Gamma(-\alpha-\alpha'+\gamma+\rho)\Gamma(-\alpha'-\beta+\gamma+\rho)}x^{-\alpha-\alpha'+\gamma+\rho-1}.$$

Expression 5. The following expression is taken from a source-backed formula corpus.

$$|\lim_{\mathcal{T}\rightarrow\infty}\frac{1}{\mathcal{T}}\int_0^{\mathcal{T}}\tilde{h}(\mu^{N,M}(t),z^{N,M}(t))dt-\int_F\tilde{h}(\mu,z)\lambda^{N,M}(d\mu,dz)|\leq\phi_{\tilde{h}}(\mathcal{T}),\quad\lim_{\mathcal{T}\rightarrow\infty}\phi_{\tilde{h}}(\mathcal{T})=0.$$