

Inline and Display Mathematics

Source-backed grouped formula sample

1 Expressions

Expression 1. The following expression is taken from a source-backed formula corpus.

$$\left({}^c D_-^{\alpha,\alpha',\beta,\beta',\gamma} f\right)(x) = (-1)^{[\text{Re}(\gamma)]+1} \left(I_-^{-\alpha',-\alpha,-\beta',-\beta+[\text{Re}(\gamma)]+1,-\gamma+[\text{Re}(\gamma)]+1} f^{([\text{Re}(\gamma)]+1)}\right)(x),$$

Expression 2. The following expression is taken from a source-backed formula corpus.

$$\int_{\mathcal{V}\times\mathcal{V}} \varphi_\lambda \, d\widetilde{\pi} = \int_{\mathcal{V}} g \, d\gamma - \int_{\mathcal{S}_1\times\mathcal{S}_1} \lambda_1 c_1 \, d\nu_1 - \int_{\mathcal{S}_2\times\mathcal{S}_2} \lambda_2 c_2 \, d\nu_2 < \infty,$$

Expression 3. The following expression is taken from a source-backed formula corpus.

$$\exp\left(-ie\left[\frac{1}{\sqrt{\pi+gN}}\left(A_\mu^{(1)},\varepsilon_{\nu\mu}\partial_\nu\Phi^{(1)}\right)+\sum_{I=2}^N\frac{1}{\sqrt{\pi}}\left(A_\mu^{(I)},\varepsilon_{\nu\mu}\partial_\nu\Phi^{(I)}\right)\right]\right)\Bigg\rangle_{\{K^{(I)}\}}=$$

Expression 4. The following expression is taken from a source-backed formula corpus.

$$L=i\overline{\psi}'_0\gamma^\mu\partial_\mu\psi'_0+i\overline{\psi}'_0e^{it^a\omega_a}e^{-it^b\theta_b}\gamma^\mu\partial_\mu\left(e^{it^c\theta_c}e^{-it^l\omega_l}\right)\psi'_0-m\overline{\psi}'_0\psi'_0.$$

Expression 5. The following expression is taken from a source-backed formula corpus.

$$\left(I_-^{\alpha,\alpha',\beta,\beta',\gamma}t^{-\rho}\right)(x)=\frac{\Gamma(-\beta+\rho)\Gamma(\alpha+\alpha'-\gamma+\rho)\Gamma(\alpha+\beta'-\gamma+\rho)}{\Gamma(\rho)\Gamma(\alpha-\beta+\rho)\Gamma(\alpha+\alpha'+\beta'-\gamma+\rho)}x^{-\alpha-\alpha'+\gamma-\rho}.$$