

Inline and Display Mathematics

Source-backed grouped formula sample

1 Expressions

Expression 1. The following expression is taken from a source-backed formula corpus.

$$\tilde{\gamma}^a p_{0a} = 0 = \tilde{\gamma}^a f^\mu{}_a p_{0\mu} = (\tilde{\gamma}^m f^\alpha{}_m + \tilde{\gamma}^h f^\alpha{}_h) p_{0\alpha} + (\tilde{\gamma}^m f^\sigma{}_m + \tilde{\gamma}^h f^\sigma{}_h) p_{0\sigma},$$

Expression 2. The following expression is taken from a source-backed formula corpus.

$$\psi_{-}(\vec{x}) = \frac{1}{\sqrt{2}} \frac{1}{i\partial_{-} - eV_{-}} * \xi(\vec{x}) - \frac{e}{\sqrt{2}} \frac{1}{\Delta_{\perp}} * (J^i_{pzm}(x_{\perp}) - q_i) \alpha^i \frac{1}{i\partial_{-} - eV_{-}} * \psi_{+}(\vec{x}) ,$$

Expression 3. The following expression is taken from a source-backed formula corpus.

$$B_{\mu\nu}(t) = \sum_{\rho=0}^N \frac{\eta_{\mu\rho}\eta_{\nu\rho}}{\sqrt{8\omega_{\mu}}} \left[\left(1 + \frac{\omega_{\mu}}{\Omega_{\rho}}\right) e^{i\Omega_{\rho}t} + \left(1 - \frac{\omega_{\mu}}{\Omega_{\rho}}\right) e^{-i\Omega_{\rho}t} \right] .$$

Expression 4. The following expression is taken from a source-backed formula corpus.

$$\tilde{R}^{ab} = R^{ab} - K_c^a \wedge K^{cb} + \frac{1}{2} C^{ab}{}_{\cdot c} \tilde{T}^c + \frac{1}{4} \gamma_J^I \gamma_L^K \theta_I^a \theta_K^b A^J \wedge A^L + i \left(DK^{ab} - \frac{1}{4} \gamma_J^I (\theta_I^a e^b + \theta_I^b e^a) \wedge A^J \right) .$$

Expression 5. The following expression is taken from a source-backed formula corpus.

$$\|\lambda(\phi)\|_{K,m+n} = \sup_{\substack{|p|\leq m+n \\ z\in K}} |D^p(\lambda(\phi))(z)| \; , \qquad D^p = \frac{\partial^{|q|+|r|}}{\partial x_1^{q_1} \cdots \partial x_m^{q_m} \partial \theta_1^{r_1} \cdots \partial \theta_n^{r_n}} \; .$$