

In this sample we use Lato font [?] as the body font. For the math we use the same input as in The L^AT_EX Companion [?, § 12.5]).
In all examples we use Iwona scaled 1.15

Iwona regular

First some large operators both in text: $\iiint f(x, y, z) dx dy dz$ and $\prod_{\gamma \in \Gamma_{\tilde{c}}} \partial(\tilde{X}_{\gamma})$;
and also on display:

$$\begin{aligned} \iiint\limits_{\mathbb{Q}} f(w, x, y, z) dw dx dy dz &\leq \oint_{\partial Q} f' \left(\max \left\{ \frac{\|w\|}{|w^2 + x^2|}, \frac{\|z\|}{|y^2 + z^2|}, \frac{\|w \oplus z\|}{\|x \oplus y\|} \right\} \right) \\ &\approx \bigcup_{\mathbb{Q} \in \mathbb{Q}} \left[f^* \left(\frac{\int \mathbb{Q}(t)}{\sqrt{1 - t^2}} \right) \right]_{t=\alpha}^{t=\vartheta} - (\Delta + \nu - \nu)^3 \end{aligned} \quad (1)$$

For x in the open interval $] -1, 1[$ the infinite sum in Equation (8) is convergent; however, this does not hold throughout the closed interval $[-1, 1]$.

$$(1 - x)^{-k} = 1 + \sum_{j=1}^{\infty} (-1)^j \left\{ \begin{matrix} k \\ j \end{matrix} \right\} x^j \quad \text{for } k \in \mathbb{N}; k \neq 0. \quad (2)$$

Iwona condensed

First some large operators both in text: $\iiint f(x, y, z) dx dy dz$ and $\prod_{\gamma \in \Gamma_{\tilde{c}}} \partial(\tilde{X}_{\gamma})$;
and also on display:

$$\begin{aligned} \iiint\limits_{\mathbb{Q}} f(w, x, y, z) dw dx dy dz &\leq \oint_{\partial Q} f' \left(\max \left\{ \frac{\|w\|}{|w^2 + x^2|}, \frac{\|z\|}{|y^2 + z^2|}, \frac{\|w \oplus z\|}{\|x \oplus y\|} \right\} \right) \\ &\approx \bigcup_{\mathbb{Q} \in \mathbb{Q}} \left[f^* \left(\frac{\int \mathbb{Q}(t)}{\sqrt{1 - t^2}} \right) \right]_{t=\alpha}^{t=\vartheta} - (\Delta + \nu - \nu)^3 \end{aligned} \quad (3)$$

For x in the open interval $] -1, 1[$ the infinite sum in Equation (8) is conver-

gent; however, this does not hold throughout the closed interval $[-1, 1]$.

$$(1 - x)^{-k} = 1 + \sum_{j=1}^{\infty} (-1)^j \left\{ \begin{matrix} k \\ j \end{matrix} \right\} x^j \quad \text{for } k \in \mathbb{N}; k \neq 0. \quad (4)$$

Iwona light

First some large operators both in text: $\iiint_{\mathbb{Q}} f(x, y, z) dx dy dz$ and $\prod_{\gamma \in \Gamma_{\tilde{c}}} \partial(\tilde{X}_{\gamma})$; and also on display:

$$\begin{aligned} \iiint_{\mathbb{Q}} f(w, x, y, z) dw dx dy dz &\leq \oint_{\partial \mathbb{Q}} f' \left(\max \left\{ \frac{\|w\|}{|w^2 + x^2|}; \frac{\|z\|}{|y^2 + z^2|}; \frac{\|w \oplus z\|}{\|x \oplus y\|} \right\} \right) \\ &\approx \bigcup_{\mathbb{Q} \in \mathbb{Q}} \left[f^* \left(\frac{\lfloor \mathbb{Q}(t) \rfloor}{\sqrt{1 - t^2}} \right) \right]_{t=\alpha}^{t=\vartheta} - (\Delta + \nu - \nu)^3 \end{aligned} \quad (5)$$

For x in the open interval $] -1, 1[$ the infinite sum in Equation (8) is convergent; however, this does not hold throughout the closed interval $[-1, 1]$.

$$(1 - x)^{-k} = 1 + \sum_{j=1}^{\infty} (-1)^j \left\{ \begin{matrix} k \\ j \end{matrix} \right\} x^j \quad \text{for } k \in \mathbb{N}; k \neq 0. \quad (6)$$

Iwona light condensed

First some large operators both in text: $\iiint_{\mathbb{Q}} f(x, y, z) dx dy dz$ and $\prod_{\gamma \in \Gamma_{\tilde{c}}} \partial(\tilde{X}_{\gamma})$; and also on display:

$$\begin{aligned} \iiint_{\mathbb{Q}} f(w, x, y, z) dw dx dy dz &\leq \oint_{\partial \mathbb{Q}} f' \left(\max \left\{ \frac{\|w\|}{|w^2 + x^2|}; \frac{\|z\|}{|y^2 + z^2|}; \frac{\|w \oplus z\|}{\|x \oplus y\|} \right\} \right) \\ &\approx \bigcup_{\mathbb{Q} \in \tilde{\mathbb{Q}}} \left[f^* \left(\frac{\lfloor \mathbb{Q}(t) \rfloor}{\sqrt{1 - t^2}} \right) \right]_{t=\alpha}^{t=\vartheta} - (\Delta + \nu - \nu)^3 \end{aligned} \quad (7)$$

For x in the open interval $] -1, 1[$ the infinite sum in Equation (8) is convergent; however, this does not hold throughout the closed interval $[-1, 1]$.

$$(1 - x)^{-k} = 1 + \sum_{j=1}^{\infty} (-1)^j \left\{ \begin{matrix} k \\ j \end{matrix} \right\} x^j \quad \text{for } k \in \mathbb{N}; k \neq 0. \quad (8)$$