

Inline and Display Mathematics

Source-backed grouped formula sample

1 Expressions

Expression 1. The following expression is taken from a source-backed formula corpus.

$$\lambda^{a,\alpha\beta}\lambda_a^\gamma{}_\delta - \lambda^{a,\gamma\alpha}\lambda_a^\beta{}_\delta = \frac{2}{\sigma_0}\left(\Omega^{\alpha\beta}\delta^\gamma{}_\delta - 2\Omega^{\beta\gamma}\delta^\alpha{}_\delta + \Omega^{\gamma\alpha}\delta^\beta{}_\delta\right),$$

Expression 2. The following expression is taken from a source-backed formula corpus.

$$\mathcal{I}_D(\delta_1, 0) = \widehat{\mathcal{I}}_D(\delta_1, \delta_2) = \inf_{\lambda \in \mathbb{R}_+^2} \left[\langle \lambda, \delta \rangle + \sup_{\varpi \in \Pi(\mu_1, \mu_2)} \int_{\mathcal{V}} g_\lambda(s_1, s_2) d\varpi(s_1, s_2) \right],$$

Expression 3. The following expression is taken from a source-backed formula corpus.

$$\rho_{j_k,j_l}^{[n]} = \frac{G_R^2 P_T}{G_R^2 \sigma_R^2 \left[\mathbf{F}_m^H \tilde{\mathbf{W}}^{(J)} \left(\tilde{\mathbf{W}}^{(J)} \right)^H \mathbf{F}_m \right]_{n,n} + \sigma_S^2 \left[\mathbf{V}_{j_l,j_k}^H \mathbf{V}_{j_l,j_k} \right]_{n,n}},$$

Expression 4. The following expression is taken from a source-backed formula corpus.

$$\mathcal{H}^{(0)} = -\frac{2\kappa}{\sqrt{-\gamma}}\frac{\delta I^{(0)}}{\delta\gamma_{\mu\nu}}\frac{\delta I^{(0)}}{\delta\gamma_{\lambda\rho}}\left(\gamma_{\mu\rho}\gamma_{\nu\lambda}-\frac{1}{d-1}\gamma_{\mu\nu}\gamma_{\lambda\rho}\right)+\frac{\sqrt{-\gamma}}{\kappa}\Lambda=0,$$

Expression 5. The following expression is taken from a source-backed formula corpus.

$$Z = V_{p+1} \left(\frac{T}{2\pi} \right)^{(p+1)/2} \int d\tau \frac{\det(\psi, B)}{\sqrt{\det(B, B)}} \frac{\sqrt{\text{Det}' \hat{P}^\dagger \hat{P}}}{\text{Vol}(\text{CKV})} \left(\frac{\text{Det}_{N'} \hat{\Delta}}{\int d^2 \xi \sqrt{\hat{g}}} \right)^{-(p+1)/2}$$