

Differential Equations for Periods and Flat Coordinates in Two Dimensional Topological Matter Theories

W.Lerche, D.Smit, and N. Warner

Abstract

We derive directly from the N=2 LG superpotential the differential equations that determine the flat coordinates of arbitrary topological CFT's.

1 Introduction

===== FILE: hep-th9108013.tex
===== '@=11 =cmr10
scaled 3 =cmr10 scaled 2 =cmr10 scaled 1 =cmr9 =cmr6 =cmbx10 scaled 3 =cmbx10 scaled 2
=cmbx10 scaled 1 =cmbx9 =cmbx6 =cmmi10 scaled 3 ='177 =cmmi10 scaled 2 ='177 =cmmi10
scaled 1 ='177 =cmmi9 ='177 =cmmi6 ='177 =cmsy10 scaled 3 ='60 =cmsy10 scaled 2 ='60
=cmsy10 scaled 1 ='60 =cmsy9 ='60 =cmsy6 ='60 =cmex10 scaled 3 =cmex10 scaled 2 =cmex10
scaled 1 =cmex10 scaled =cmsl10 scaled 3 =cmsl10 scaled 2 =cmsl10 scaled 1 =cmsl9 =cmti10
scaled 3 =cmti10 scaled 2 =cmti10 scaled 1 =cmtt10 scaled 1 =cmcsc10 scaled 1 =cmcsc10

2 Method

2= 2= 2= 3= 3= 3= = = = = = = = @nt @big @big=8.5 @ @Big @Big=11.5 @ @bigg
@bigg=14.5 @ @Bigg @Bigg=17.5 @ #1 #1 #1 #1 = 20pt plus 0.2pt minus 0.1pt = 1.5pt plus
0.1pt minus 0.1pt = 1.5pt

$$L(\theta) = \sum_{i=1}^n \ell(x_i, \theta) \quad (1)$$

Equation 1 is included to keep the sample paper-like while remaining small.

3 Conclusion

@t @ @ @t3:4; @t1:1; @ @t3:5; @t4:5; @ by 5 by 4 @t#1:#2; = by #1 by #2 = by #1 by #2 =
by #1 by #2 = by #1 by #2 = by #1 by #2 = = by #1 by #2 = by by 2 = by #1 by #2 = by
2 = by 4