

# CS 577- Intro to Algorithms

## Dynamic Programming (Part 4)

Dieter van Melkebeek

October 1, 2020

# Outline

# Outline

## Paradigm

Recursive approach such that:

1. The number of distinct subproblems in the recursion tree is small.
2. Each of those subproblems is solved only once.

# Outline

## Paradigm

Recursive approach such that:

1. The number of distinct subproblems in the recursion tree is small.
2. Each of those subproblems is solved only once.

## Examples

- ▶ Computing Fibonacci numbers
- ▶ Weighted interval scheduling
- ▶ Knapsack problem
- ▶ RNA secondary structure
- ▶ Sequence alignment / Edit distance

# Outline

## Paradigm

Recursive approach such that:

1. The number of distinct subproblems in the recursion tree is small.
2. Each of those subproblems is solved only once.

## Examples

- ▶ Computing Fibonacci numbers
- ▶ Weighted interval scheduling
- ▶ Knapsack problem
- ▶ RNA secondary structure
- ▶ Sequence alignment / Edit distance
- ▶ Shortest paths

# Shortest Paths Problem

# Shortest Paths Problem

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

# Shortest Paths Problem

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$



# Shortest Paths Problem

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

Variants based on source/target

# Shortest Paths Problem

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

Variants based on source/target

- ▶ single pair

# Shortest Paths Problem

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

Variants based on source/target

- ▶ single pair
- ▶ single source

# Shortest Paths Problem

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

Variants based on source/target

- ▶ single pair
- ▶ single source
- ▶ single target

# Shortest Paths Problem

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

Variants based on source/target

- ▶ single pair
- ▶ single source
- ▶ single target
- ▶ all pairs

# Shortest Paths Problem

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

## Variants based on source/target

- ▶ single pair
- ▶ single source
- ▶ single target
- ▶ all pairs

## Variants based on edge lengths

# Shortest Paths Problem

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

## Variants based on source/target

- ▶ single pair
- ▶ single source
- ▶ single target
- ▶ all pairs

## Variants based on edge lengths

- ▶ unit

# Shortest Paths Problem

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

## Variants based on source/target

- ▶ single pair
- ▶ single source
- ▶ single target
- ▶ all pairs

## Variants based on edge lengths

- ▶ unit
- ▶ nonnegative



# Shortest Paths Problem

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

## Variants based on source/target

- ▶ single pair
- ▶ single source
- ▶ single target
- ▶ all pairs

## Variants based on edge lengths

- ▶ unit
- ▶ nonnegative
- ▶ arbitrary

# Shortest Paths Problem

## Specification

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

# Shortest Paths Problem

## Specification

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

**Distance**  $d(s, t)$

# Shortest Paths Problem

## Specification

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length

$$\ell(P) \doteq \sum_{e \in P} \ell(e)$$

**Distance**  $d(s, t)$

$$= \min\{\ell(P) \mid P \text{ path from } s \text{ to } t\}$$

# Shortest Paths Problem

## Specification

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

**Distance**  $d(s, t)$

$= \min\{\ell(P) \mid P \text{ path from } s \text{ to } t\}$

$= \infty$  if there is no path from  $s$  to  $t$

# Shortest Paths Problem

## Specification

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

**Distance**  $d(s, t)$

$= \min\{\ell(P) \mid P \text{ path from } s \text{ to } t\}$

$= \infty$  if there is no path from  $s$  to  $t$

$= -\infty$  if there is a path from  $s$  to  $t$  but no shortest one

# Shortest Paths Problem

## Specification

**Input:** (di)graph  $G = (V, E)$ ; lengths  $\ell : E \rightarrow \mathbb{R}$ ;  $s, t \in V$

**Output:** path  $P$  from  $s$  to  $t$  with minimum length  
 $\ell(P) \doteq \sum_{e \in P} \ell(e)$

## Distance $d(s, t)$

$= \min\{\ell(P) \mid P \text{ path from } s \text{ to } t\}$

$= \infty$  if there is no path from  $s$  to  $t$

$= -\infty$  if there is a path from  $s$  to  $t$  but no shortest one

## Proposition

$d(s, t) = -\infty \Leftrightarrow$

there exists cycle  $C$  with  $\ell(C) < 0$  such that  $s \rightsquigarrow C$  and  $C \rightsquigarrow t$ .

# Principle of optimality



# Single Source – subproblems and recurrence

## Subproblems

$\text{OPT}(k, v) =$

length of a shortest path from  $s$  to  $v$  using  $\leq k$  edges

$\infty$  if no such path exists

# Single Source – subproblems and recurrence

## Subproblems

$\text{OPT}(k, v) =$

length of a shortest path from  $s$  to  $v$  using  $\leq k$  edges

$\infty$  if no such path exists

$(k \in \mathbb{N} \text{ and } v \in V)$

# Single Source – subproblems and recurrence

## Subproblems

$\text{OPT}(k, v) =$

length of a shortest path from  $s$  to  $v$  using  $\leq k$  edges

$\infty$  if no such path exists

$(k \in \mathbb{N} \text{ and } v \in V)$

Base case ( $k = 0$ )

# Single Source – subproblems and recurrence

## Subproblems

$\text{OPT}(k, v) =$

length of a shortest path from  $s$  to  $v$  using  $\leq k$  edges

$\infty$  if no such path exists

$(k \in \mathbb{N} \text{ and } v \in V)$

Base case ( $k = 0$ )

$\text{OPT}(0, s) = 0$  and  $\text{OPT}(0, v) = \infty$  for  $v \neq s$

# Single Source – subproblems and recurrence

## Subproblems

$\text{OPT}(k, v) =$

length of a shortest path from  $s$  to  $v$  using  $\leq k$  edges

$\infty$  if no such path exists

$(k \in \mathbb{N} \text{ and } v \in V)$

Base case ( $k = 0$ )

$\text{OPT}(0, s) = 0$  and  $\text{OPT}(0, v) = \infty$  for  $v \neq s$

Recursive case ( $k \geq 1$ )

# Single Source – subproblems and recurrence

## Subproblems

$$\text{OPT}(k, v) =$$

length of a shortest path from  $s$  to  $v$  using  $\leq k$  edges

$\infty$  if no such path exists

$$(k \in \mathbb{N} \text{ and } v \in V)$$

## Base case ( $k = 0$ )

$$\text{OPT}(0, s) = 0 \text{ and } \text{OPT}(0, v) = \infty \text{ for } v \neq s$$

## Recursive case ( $k \geq 1$ )

$$\text{OPT}(k, v) =$$

$$\min_{(u,v) \in E} (\text{OPT}(k-1, u) + \ell(u, v))$$

# Single Source – subproblems and recurrence

## Subproblems

$$\text{OPT}(k, v) =$$

length of a shortest path from  $s$  to  $v$  using  $\leq k$  edges

$\infty$  if no such path exists

$$(k \in \mathbb{N} \text{ and } v \in V)$$

## Base case ( $k = 0$ )

$$\text{OPT}(0, s) = 0 \text{ and } \text{OPT}(0, v) = \infty \text{ for } v \neq s$$

## Recursive case ( $k \geq 1$ )

$$\text{OPT}(k, v) =$$

$$\min_{(u,v) \in E} (\text{OPT}(k-1, u) + \ell(u, v))$$

$$\text{OPT}(k-1, v)$$

# Single Source – subproblems and recurrence

## Subproblems

$$\text{OPT}(k, v) =$$

length of a shortest path from  $s$  to  $v$  using  $\leq k$  edges

$\infty$  if no such path exists

$$(k \in \mathbb{N} \text{ and } v \in V)$$

## Base case ( $k = 0$ )

$$\text{OPT}(0, s) = 0 \text{ and } \text{OPT}(0, v) = \infty \text{ for } v \neq s$$

## Recursive case ( $k \geq 1$ )

$$\text{OPT}(k, v) =$$

$$\min_{(u,v) \in E} (\text{OPT}(k-1, u) + \ell(u, v))$$

$$\text{OPT}(k-1, v)$$

## Answer

$$d(s, v)$$



# Single Source – subproblems and recurrence

## Subproblems

$$\text{OPT}(k, v) =$$

length of a shortest path from  $s$  to  $v$  using  $\leq k$  edges

$\infty$  if no such path exists

$$(k \in \mathbb{N} \text{ and } v \in V)$$

## Base case ( $k = 0$ )

$$\text{OPT}(0, s) = 0 \text{ and } \text{OPT}(0, v) = \infty \text{ for } v \neq s$$

## Recursive case ( $k \geq 1$ )

$$\text{OPT}(k, v) =$$

$$\min_{(u,v) \in E} (\text{OPT}(k-1, u) + \ell(u, v))$$

$$\text{OPT}(k-1, v)$$

## Answer

$$d(s, v) = \lim_{k \rightarrow \infty} \text{OPT}(k, v)$$

# Single Source – number of iterations

# Single Source – number of iterations

## Observation 1

If  $(\forall v) \text{OPT}(k, v) = \text{OPT}(k - 1, v)$   
then  $(\forall v) \text{OPT}(k + 1, v) = \text{OPT}(k, v)$ .

# Single Source – number of iterations

## Observation 1

If  $(\forall v) \text{OPT}(k, v) = \text{OPT}(k - 1, v)$   
then  $(\forall v) \text{OPT}(k + 1, v) = \text{OPT}(k, v)$ .

## Note

If  $\ell \geq 0$ , the hypothesis of Observation 1 holds for  $k = n$ .

# Single Source – number of Iterations

## Observation 2

If  $\text{OPT}(n, v) < \text{OPT}(n - 1, v)$  then

there exists cycle  $C$  with  $\ell(C) < 0$  such that  $s \rightsquigarrow C$  and  $C \rightsquigarrow v$ .

# Single Source – number of Iterations

## Observation 2

If  $\text{OPT}(n, v) < \text{OPT}(n-1, v)$  then

there exists cycle  $C$  with  $\ell(C) < 0$  such that  $s \rightsquigarrow C$  and  $C \rightsquigarrow v$ .

## Proof

$$\begin{aligned}\text{OPT}(n, v) &= \ell(P) \\ &= \ell(P - C) + \ell(C) \\ &\geq \text{OPT}(n-1, v) + \ell(C) \\ &> \text{OPT}(n, v) + \ell(C) \\ \Rightarrow \ell(C) &< 0\end{aligned}$$

# Single Source – number of Iterations

## Observation 1

If  $(\forall v) \text{OPT}(k, v) = \text{OPT}(k - 1, v)$   
then  $(\forall v) \text{OPT}(k + 1, v) = \text{OPT}(k, v)$ .

## Observation 2

If  $\text{OPT}(n, u) < \text{OPT}(n - 1, u)$  then  
there exists cycle  $C$  with  $\ell(C) < 0$  such that  $s \rightsquigarrow C$  and  $C \rightsquigarrow u$ .

## Criterion

$d(s, v) = -\infty \Leftrightarrow$   
there exist  $u \in V$  s.t.  $u \rightsquigarrow v$  and  $\text{OPT}(n, u) < \text{OPT}(n - 1, u)$ .

# Single Source – algorithm



# Single Source – algorithm

## Recurrence

- ▶  $\text{OPT}(0, s) = 0$  and  $\text{OPT}(0, v) = \infty$  for  $v \neq s$
- ▶  $\text{OPT}(k, v) =$   
 $\min(\text{OPT}(k-1, v), \min_{(u,v) \in E} (\text{OPT}(k-1, u) + \ell(u, v)))$

# Single Source – complexity analysis

## Recurrence

- ▶  $\text{OPT}(0, s) = 0$  and  $\text{OPT}(0, v) = \infty$  for  $v \neq s$
- ▶  $\text{OPT}(k, v) = \min(\text{OPT}(k-1, v), \min_{(u,v) \in E} (\text{OPT}(k-1, u) + \ell(u, v)))$

## Time

- ▶  $O(n + m)$  per row
- ▶  $\leq n + 1$  rows
- ▶  $O(n(n + m))$  total

## Space

- ▶  $O(n)$  for computing distances
- ▶  $O(n^2) \rightarrow O(n)$  for finding shortest paths

# Principle of optimality

# Principle of optimality

$$V = [n]$$

# All Pairs – subproblems

## Subproblems

$\text{OPT}(s, t, k) =$

length of a shortest path from  $s$  to  $t$

that only uses  $\{k, \dots, n\}$  as intermediate vertices

# All Pairs – subproblems

## Subproblems

$\text{OPT}(s, t, k) =$

length of a shortest path from  $s$  to  $t$

that only uses  $\{k, \dots, n\}$  as intermediate vertices

$\infty$  if no such path exists

# All Pairs – subproblems

## Subproblems

$\text{OPT}(s, t, k) =$

length of a shortest path from  $s$  to  $t$

that only uses  $\{k, \dots, n\}$  as intermediate vertices

$\infty$  if no such path exists

$-\infty$  if there is such a path but no shortest one

# All Pairs – subproblems

## Subproblems

$\text{OPT}(s, t, k) =$

length of a shortest path from  $s$  to  $t$

that only uses  $\{k, \dots, n\}$  as intermediate vertices

$\infty$  if no such path exists

$-\infty$  if there is such a path but no shortest one

$(s, t \in V \text{ and } k \in [n + 1])$



# All Pairs – subproblems

## Subproblems

$\text{OPT}(s, t, k) =$

length of a shortest path from  $s$  to  $t$

that only uses  $\{k, \dots, n\}$  as intermediate vertices

$\infty$  if no such path exists

$-\infty$  if there is such a path but no shortest one

$(s, t \in V \text{ and } k \in [n + 1])$

Base case ( $k = n + 1$ )

# All Pairs – subproblems

## Subproblems

$\text{OPT}(s, t, k) =$

length of a shortest path from  $s$  to  $t$

that only uses  $\{k, \dots, n\}$  as intermediate vertices

$\infty$  if no such path exists

$-\infty$  if there is such a path but no shortest one

$(s, t \in V \text{ and } k \in [n + 1])$

Base case ( $k = n + 1$ )

►  $\text{OPT}(s, t, n + 1) = 0$  if  $s = t$

# All Pairs – subproblems

## Subproblems

$\text{OPT}(s, t, k) =$

length of a shortest path from  $s$  to  $t$

that only uses  $\{k, \dots, n\}$  as intermediate vertices

$\infty$  if no such path exists

$-\infty$  if there is such a path but no shortest one

$(s, t \in V \text{ and } k \in [n + 1])$

Base case ( $k = n + 1$ )

- ▶  $\text{OPT}(s, t, n + 1) = 0$  if  $s = t$
- ▶  $\text{OPT}(s, t, n + 1) = \ell(s, t)$  if  $(s, t) \in E$

# All Pairs – subproblems

## Subproblems

$\text{OPT}(s, t, k) =$

length of a shortest path from  $s$  to  $t$

that only uses  $\{k, \dots, n\}$  as intermediate vertices

$\infty$  if no such path exists

$-\infty$  if there is such a path but no shortest one

$(s, t \in V \text{ and } k \in [n + 1])$

Base case ( $k = n + 1$ )

- ▶  $\text{OPT}(s, t, n + 1) = 0$  if  $s = t$
- ▶  $\text{OPT}(s, t, n + 1) = \ell(s, t)$  if  $(s, t) \in E$
- ▶  $\text{OPT}(s, t, n + 1) = \infty$  otherwise

# All Pairs – subproblems

## Subproblems

$\text{OPT}(s, t, k) =$

length of a shortest path from  $s$  to  $t$

that only uses  $\{k, \dots, n\}$  as intermediate vertices

$\infty$  if no such path exists

$-\infty$  if there is such a path but no shortest one

$(s, t \in V \text{ and } k \in [n + 1])$

Base case ( $k = n + 1$ )

- ▶  $\text{OPT}(s, t, n + 1) = 0$  if  $s = t$
- ▶  $\text{OPT}(s, t, n + 1) = \ell(s, t)$  if  $(s, t) \in E$
- ▶  $\text{OPT}(s, t, n + 1) = \infty$  otherwise

Answer

$d(s, t) = \text{OPT}(s, t, 1)$

# All Pairs – recurrence and analysis

# All Pairs – recurrence and analysis

Recursive case ( $k \leq n$ )

# All Pairs – recurrence and analysis

Recursive case ( $k \leq n$ )

If  $\text{OPT}(s, k, k + 1) < \infty$  and  $\text{OPT}(k, t, k + 1) < \infty$

and  $\text{OPT}(k, k, k + 1) < 0$

then  $\text{OPT}(s, t, k) = -\infty$

else  $\text{OPT}(s, t, k) =$

$\min(\text{OPT}(s, t, k + 1), \text{OPT}(s, k, k + 1) + \text{OPT}(k, t, k + 1))$



# All Pairs – recurrence and analysis

## Recursive case ( $k \leq n$ )

If  $\text{OPT}(s, k, k+1) < \infty$  and  $\text{OPT}(k, t, k+1) < \infty$

and  $\text{OPT}(k, k, k+1) < 0$

then  $\text{OPT}(s, t, k) = -\infty$

else  $\text{OPT}(s, t, k) =$

$\min(\text{OPT}(s, t, k+1), \text{OPT}(s, k, k+1) + \text{OPT}(k, t, k+1))$

## Time

- ▶  $O(1)$  per entry
- ▶  $O(n^3)$  entries
- ▶  $O(n^3)$  total

# All Pairs – recurrence and analysis

## Recursive case ( $k \leq n$ )

If  $\text{OPT}(s, k, k+1) < \infty$  and  $\text{OPT}(k, t, k+1) < \infty$

and  $\text{OPT}(k, k, k+1) < 0$

then  $\text{OPT}(s, t, k) = -\infty$

else  $\text{OPT}(s, t, k) =$

$\min(\text{OPT}(s, t, k+1), \text{OPT}(s, k, k+1) + \text{OPT}(k, t, k+1))$

## Time

- ▶  $O(1)$  per entry
- ▶  $O(n^3)$  entries
- ▶  $O(n^3)$  total

## Space

$O(n^2)$  for distances and shortest paths