

# CS 577- Intro to Algorithms

## Network Flow (Part 3)

Dieter van Melkebeek

November 3, 2020

# Outline

# Outline

## Recap

- ▶ Notions: network, flow, cut
- ▶ Computational problems: max flow vs min cut

# Outline

## Recap

- ▶ Notions: network, flow, cut
- ▶ Computational problems: max flow vs min cut

## Applications of max flow

- ▶ Bipartite matching
- ▶ Edge-disjoint paths
- ▶ Survey design

# Recap - notions

# Recap - notions

## Network

- ▶ a digraph  $(V, E)$
- ▶ edge capacities  $c : E \rightarrow [0, \infty)$
- ▶ the source  $s \in V$ , which has indegree 0, and
- ▶ the sink  $t \in V$ , which has outdegree 0.

# Recap - notions

## Network

- ▶ a digraph  $(V, E)$
- ▶ edge capacities  $c : E \rightarrow [0, \infty)$
- ▶ the source  $s \in V$ , which has indegree 0, and
- ▶ the sink  $t \in V$ , which has outdegree 0.

## Flow

A mapping  $f : E \rightarrow [0, \infty)$  satisfying

- ▶ [capacity constraints]  $(\forall e \in E) f(e) \leq c(e)$
- ▶ [conservation constraints]  $(\forall v \in V \setminus \{s, t\}) f_{\text{in}}(v) = f_{\text{out}}(v)$   
where  $f_{\text{in}}(v) \doteq \sum_{u \in V: e \doteq (u, v) \in E} f(e)$  and  
 $f_{\text{out}}(v) \doteq \sum_{u \in V: e \doteq (v, u) \in E} f(e)$ .

# Recap - notions

## Network

- ▶ a digraph  $(V, E)$
- ▶ edge capacities  $c : E \rightarrow [0, \infty)$
- ▶ the source  $s \in V$ , which has indegree 0, and
- ▶ the sink  $t \in V$ , which has outdegree 0.

## Flow

A mapping  $f : E \rightarrow [0, \infty)$  satisfying

- ▶ [capacity constraints]  $(\forall e \in E) f(e) \leq c(e)$
- ▶ [conservation constraints]  $(\forall v \in V \setminus \{s, t\}) f_{\text{in}}(v) = f_{\text{out}}(v)$   
where  $f_{\text{in}}(v) \doteq \sum_{u \in V: e \doteq (u, v) \in E} f(e)$  and  
 $f_{\text{out}}(v) \doteq \sum_{u \in V: e \doteq (v, u) \in E} f(e)$ .

## $st$ -Cut

A partition  $(S, T)$  of  $V$  such that  $s \in S$  and  $t \in T$ .



# Recap - notions

## Integral network

- ▶ a digraph  $(V, E)$
- ▶ edge capacities  $c : E \rightarrow \mathbb{N}$
- ▶ the source  $s \in V$ , which has indegree 0, and
- ▶ the sink  $t \in V$ , which has outdegree 0.

## Integral flow

A mapping  $f : E \rightarrow \mathbb{N}$  satisfying

- ▶ [capacity constraints]  $(\forall e \in E) f(e) \leq c(e)$
- ▶ [conservation constraints]  $(\forall v \in V \setminus \{s, t\}) f_{\text{in}}(v) = f_{\text{out}}(v)$   
where  $f_{\text{in}}(v) \doteq \sum_{u \in V: e \doteq (u, v) \in E} f(e)$  and  
 $f_{\text{out}}(v) \doteq \sum_{u \in V: e \doteq (v, u) \in E} f(e)$ .

## st-Cut

A partition  $(S, T)$  of  $V$  such that  $s \in S$  and  $t \in T$ .

# Recap - computational problems

# Recap - computational problems

## Max flow

**Input:** network  $N = (V, E, c, s, t)$

**Output:** flow  $f$  such that  $\nu(f) \doteq f_{\text{out}}(s)$  is maximized

# Recap - computational problems

## Max flow

**Input:** network  $N = (V, E, c, s, t)$

**Output:** flow  $f$  such that  $\nu(f) \doteq f_{\text{out}}(s)$  is maximized

**Complexity:** time  $O(nm)$

# Recap - computational problems

## Max flow

**Input:** network  $N = (V, E, c, s, t)$

**Output:** flow  $f$  such that  $\nu(f) \doteq f_{\text{out}}(s)$  is maximized

**Complexity:** time  $O(nm)$

**Note:** integrality preserved

# Recap - computational problems

## Max flow

**Input:** network  $N = (V, E, c, s, t)$

**Output:** flow  $f$  such that  $\nu(f) \doteq f_{\text{out}}(s)$  is maximized

**Complexity:** time  $O(nm)$

**Note:** integrality preserved

## Min cut

**Input:** network  $N = (V, E, c, s, t)$

**Output:**  $st$ -cut  $(S, T)$  such that  $c(S, T) \doteq \sum_{e \in S \times T} c(e)$  is minimized

# Recap - computational problems

## Max flow

**Input:** network  $N = (V, E, c, s, t)$

**Output:** flow  $f$  such that  $\nu(f) \doteq f_{\text{out}}(s)$  is maximized

**Complexity:** time  $O(nm)$

**Note:** integrality preserved

## Min cut

**Input:** network  $N = (V, E, c, s, t)$

**Output:**  $st$ -cut  $(S, T)$  such that  $c(S, T) \doteq \sum_{e \in S \times T} c(e)$  is minimized

**Complexity:** time  $O(n + m)$  when given max flow

# Recap - computational problems

## Max flow

**Input:** network  $N = (V, E, c, s, t)$

**Output:** flow  $f$  such that  $\nu(f) \doteq f_{\text{out}}(s)$  is maximized

**Complexity:** time  $O(nm)$

**Note:** integrality preserved

## Min cut

**Input:** network  $N = (V, E, c, s, t)$

**Output:**  $st$ -cut  $(S, T)$  such that  $c(S, T) \doteq \sum_{e \in S \times T} c(e)$  is minimized

**Complexity:** time  $O(n + m)$  when given max flow

## Duality

$$\max_{\text{flow } f} (\nu(f)) = \min_{st\text{-cut } (S, T)} (c(S, T))$$



# Bipartite Matching

# Bipartite Matching

## Definition

A matching  $M$  in a graph  $G = (V, E)$  is a subset  $M \subseteq E$  such that each  $v \in V$  appears in at most one  $e \in M$ .

# Bipartite Matching

## Definition

A **perfect** matching  $M$  in a graph  $G = (V, E)$  is a subset  $M \subseteq E$  such that each  $v \in V$  appears in **exactly** one  $e \in M$ .

# Bipartite Matching

## Definition

A **perfect** matching  $M$  in a graph  $G = (V, E)$  is a subset  $M \subseteq E$  such that each  $v \in V$  appears in **exactly** one  $e \in M$ .

## Computational problem

**Input:** bipartite graph  $G$  with bipartition  $(L, R)$

**Output:** matching  $M$  such that  $|M|$  is maximized

# Bipartite Matching

## Definition

A **perfect** matching  $M$  in a graph  $G = (V, E)$  is a subset  $M \subseteq E$  such that each  $v \in V$  appears in **exactly** one  $e \in M$ .

## Computational problem

**Input:** bipartite graph  $G$  with bipartition  $(L, R)$

**Output:** matching  $M$  such that  $|M|$  is maximized

## Reduction to max flow

# Bipartite Matching

## Definition

A **perfect** matching  $M$  in a graph  $G = (V, E)$  is a subset  $M \subseteq E$  such that each  $v \in V$  appears in **exactly** one  $e \in M$ .

## Computational problem

**Input:** bipartite graph  $G$  with bipartition  $(L, R)$

**Output:** matching  $M$  such that  $|M|$  is maximized

## Reduction to max flow

- Construction of integral network  $N$  such that

$$\text{matching } M \text{ in } G \xleftrightarrow{\text{bijection}} \text{integral flow } f \text{ in } N$$

# Bipartite Matching

## Definition

A **perfect** matching  $M$  in a graph  $G = (V, E)$  is a subset  $M \subseteq E$  such that each  $v \in V$  appears in **exactly** one  $e \in M$ .

## Computational problem

**Input:** bipartite graph  $G$  with bipartition  $(L, R)$

**Output:** matching  $M$  such that  $|M|$  is maximized

## Reduction to max flow

- Construction of integral network  $N$  such that

$$\begin{array}{ccc} \text{matching } M \text{ in } G & \xleftrightarrow{\text{bijection}} & \text{integral flow } f \text{ in } N \\ |M| & = & \nu(f) \end{array}$$

# Bipartite Matching

## Definition

A **perfect** matching  $M$  in a graph  $G = (V, E)$  is a subset  $M \subseteq E$  such that each  $v \in V$  appears in **exactly** one  $e \in M$ .

## Computational problem

**Input:** bipartite graph  $G$  with bipartition  $(L, R)$

**Output:** matching  $M$  such that  $|M|$  is maximized

## Reduction to max flow

- Construction of integral network  $N$  such that

$$\begin{array}{ccc} \text{matching } M \text{ in } G & \xleftrightarrow{\text{bijection}} & \text{integral flow } f \text{ in } N \\ |M| & = & \nu(f) \end{array}$$

- Resulting algorithm runs in time  $O(nm)$ .



# Bipartite Matching – duality

# Bipartite Matching – duality

Capacity of a cut in  $N$

# Bipartite Matching – duality

## Capacity of a cut in $N$

- ▶ For any  $A \subseteq V$ , let  $G(A)$  denote all the neighbors of  $A$  in  $G$ :  
 $G(A) \doteq \{v : (\exists u \in A) (u, v) \in E\}.$

# Bipartite Matching – duality

## Capacity of a cut in $N$

- ▶ For any  $A \subseteq V$ , let  $G(A)$  denote all the neighbors of  $A$  in  $G$ :  
 $G(A) \doteq \{v : (\exists u \in A) (u, v) \in E\}.$
- ▶  $c(S, T)$  is finite iff  $G(L \cap S) \subseteq S$ .

# Bipartite Matching – duality

## Capacity of a cut in $N$

- ▶ For any  $A \subseteq V$ , let  $G(A)$  denote all the neighbors of  $A$  in  $G$ :  
 $G(A) \doteq \{v : (\exists u \in A) (u, v) \in E\}.$
- ▶  $c(S, T)$  is finite iff  $G(L \cap S) \subseteq S$ .
- ▶ If  $c(S, T)$  is finite then  $c(S, T) = |L \cap T| + |R \cap S|.$

# Bipartite Matching – duality

## Capacity of a cut in $N$

- ▶ For any  $A \subseteq V$ , let  $G(A)$  denote all the neighbors of  $A$  in  $G$ :  
 $G(A) \doteq \{v : (\exists u \in A) (u, v) \in E\}.$
- ▶  $c(S, T)$  is finite iff  $G(L \cap S) \subseteq S$ .
- ▶ If  $c(S, T)$  is finite then  $c(S, T) = |L \cap T| + |R \cap S|.$

## Marriage theorem

A bipartite graph  $G = (V, E)$  with bipartition  $(L, R)$  and  $|L| = |R| = n$  has a perfect matching iff

$$(\forall A \subseteq L) |G(A)| \geq |A|.$$

# Bipartite Matching – duality

## Capacity of a cut in $N$

- ▶ For any  $A \subseteq V$ , let  $G(A)$  denote all the neighbors of  $A$  in  $G$ :  
 $G(A) \doteq \{v : (\exists u \in A) (u, v) \in E\}.$
- ▶  $c(S, T)$  is finite iff  $G(L \cap S) \subseteq S$ .
- ▶ If  $c(S, T)$  is finite then  $c(S, T) = |L \cap T| + |R \cap S|.$

## Marriage theorem

A bipartite graph  $G = (V, E)$  with bipartition  $(L, R)$  and  $|L| = |R| = n$  has a perfect matching iff

$$(\forall A \subseteq L) |G(A)| \geq |A|.$$

## Proof

# Bipartite Matching – duality

## Capacity of a cut in $N$

- ▶ For any  $A \subseteq V$ , let  $G(A)$  denote all the neighbors of  $A$  in  $G$ :  
 $G(A) \doteq \{v : (\exists u \in A) (u, v) \in E\}.$
- ▶  $c(S, T)$  is finite iff  $G(L \cap S) \subseteq S$ .
- ▶ If  $c(S, T)$  is finite then  $c(S, T) = |L \cap T| + |R \cap S|.$

## Marriage theorem

A bipartite graph  $G = (V, E)$  with bipartition  $(L, R)$  and  $|L| = |R| = n$  has a perfect matching iff

$$(\forall A \subseteq L) |G(A)| \geq |A|.$$

## Proof

$\Rightarrow$ : Contrapositive follows from definition of perfect matching.



# Bipartite Matching – duality

## Capacity of a cut in $N$

- ▶ For any  $A \subseteq V$ , let  $G(A)$  denote all the neighbors of  $A$  in  $G$ :  
 $G(A) \doteq \{v : (\exists u \in A) (u, v) \in E\}$ .
- ▶  $c(S, T)$  is finite iff  $G(L \cap S) \subseteq S$ .
- ▶ If  $c(S, T)$  is finite then  $c(S, T) = |L \cap T| + |R \cap S|$ .

## Marriage theorem

A bipartite graph  $G = (V, E)$  with bipartition  $(L, R)$  and  $|L| = |R| = n$  has a perfect matching iff

$$(\forall A \subseteq L) |G(A)| \geq |A|.$$

## Proof

$\Rightarrow$ : Contrapositive follows from definition of perfect matching.

$\Leftarrow$ : Proof of contrapositive on next slide.

# Proof of Marriage Theorem

- ▶ Suppose  $G = (V, E)$  with bipartition  $(L, R)$  and  $|L| = |R| = n$  has no perfect matching.

# Proof of Marriage Theorem

- ▶ Suppose  $G = (V, E)$  with bipartition  $(L, R)$  and  $|L| = |R| = n$  has no perfect matching.
- ▶  $\nu(f) < n$  for every flow  $f$  in  $N$ .

# Proof of Marriage Theorem

- ▶ Suppose  $G = (V, E)$  with bipartition  $(L, R)$  and  $|L| = |R| = n$  has no perfect matching.
- ▶  $\nu(f) < n$  for every flow  $f$  in  $N$ .
- ▶ For every min cut  $(S, T)$  in  $N$ :

# Proof of Marriage Theorem

- ▶ Suppose  $G = (V, E)$  with bipartition  $(L, R)$  and  $|L| = |R| = n$  has no perfect matching.
- ▶  $\nu(f) < n$  for every flow  $f$  in  $N$ .
- ▶ For every min cut  $(S, T)$  in  $N$ :
  - (1)  $c(S, T) < n$

# Proof of Marriage Theorem

- ▶ Suppose  $G = (V, E)$  with bipartition  $(L, R)$  and  $|L| = |R| = n$  has no perfect matching.
- ▶  $\nu(f) < n$  for every flow  $f$  in  $N$ .
- ▶ For every min cut  $(S, T)$  in  $N$ :
  - (1)  $c(S, T) < n$
  - (2)  $G(L \cap S) \subseteq R \cap S$

# Proof of Marriage Theorem

- ▶ Suppose  $G = (V, E)$  with bipartition  $(L, R)$  and  $|L| = |R| = n$  has no perfect matching.
- ▶  $\nu(f) < n$  for every flow  $f$  in  $N$ .
- ▶ For every min cut  $(S, T)$  in  $N$ :
  - (1)  $c(S, T) < n$
  - (2)  $G(L \cap S) \subseteq R \cap S$
  - (3)  $c(S, T) = |L \cap T| + |R \cap S|$

# Proof of Marriage Theorem

- ▶ Suppose  $G = (V, E)$  with bipartition  $(L, R)$  and  $|L| = |R| = n$  has no perfect matching.
- ▶  $\nu(f) < n$  for every flow  $f$  in  $N$ .
- ▶ For every min cut  $(S, T)$  in  $N$ :
  - (1)  $c(S, T) < n$
  - (2)  $G(L \cap S) \subseteq R \cap S$
  - (3)  $c(S, T) = |L \cap T| + |R \cap S|$
- ▶ Consider  $A \doteq L \cap S$ .



# Proof of Marriage Theorem

- ▶ Suppose  $G = (V, E)$  with bipartition  $(L, R)$  and  $|L| = |R| = n$  has no perfect matching.
- ▶  $\nu(f) < n$  for every flow  $f$  in  $N$ .
- ▶ For every min cut  $(S, T)$  in  $N$ :
  - (1)  $c(S, T) < n$
  - (2)  $G(L \cap S) \subseteq R \cap S$
  - (3)  $c(S, T) = |L \cap T| + |R \cap S|$
- ▶ Consider  $A \doteq L \cap S$ .

$$|G(A)| \stackrel{(2)}{\leq} |R \cap S| \stackrel{(3)}{=} c(S, T) - |L \cap T| \stackrel{(1)}{<} n - |L \cap T| = |L \cap S| = |A|$$

# Edge-Disjoint Paths

# Edge-Disjoint Paths

## Computational problem

**Input:** digraph  $G = (V, E)$ ;  $s, t \in V$

**Output:** set  $C$  of edge-disjoint  $st$ -paths in  $G$  such that  $|C|$  is maximized

# Edge-Disjoint Paths

## Computational problem

**Input:** digraph  $G = (V, E)$ ;  $s, t \in V$

**Output:** set  $C$  of edge-disjoint  $st$ -paths in  $G$  such that  $|C|$  is maximized

## Reduction to max flow

# Edge-Disjoint Paths

## Computational problem

**Input:** digraph  $G = (V, E)$ ;  $s, t \in V$

**Output:** set  $C$  of edge-disjoint  $st$ -paths in  $G$  such that  $|C|$  is maximized

## Reduction to max flow

- ▶ Integral network  $N \doteq (V, E \setminus (V \times \{s\} \cup \{t\} \times V), c \equiv 1, s, t)$

# Edge-Disjoint Paths

## Computational problem

**Input:** digraph  $G = (V, E)$ ;  $s, t \in V$

**Output:** set  $C$  of edge-disjoint  $st$ -paths in  $G$  such that  $|C|$  is maximized

## Reduction to max flow

► Integral network  $N \doteq (V, E \setminus (V \times \{s\} \cup \{t\} \times V), c \equiv 1, s, t)$

set  $C$  of edge-disjoint  $st$ -paths in  $G \iff$  integral flow  $f$  in  $N$

# Edge-Disjoint Paths

## Computational problem

**Input:** digraph  $G = (V, E)$ ;  $s, t \in V$

**Output:** set  $C$  of edge-disjoint  $st$ -paths in  $G$  such that  $|C|$  is maximized

## Reduction to max flow

► Integral network  $N \doteq (V, E \setminus (V \times \{s\} \cup \{t\} \times V), c \equiv 1, s, t)$

set  $C$  of edge-disjoint  $st$ -paths in  $G$   $\longleftrightarrow$  integral flow  $f$  in  $N$   
 $|C| = \nu(f)$

# Edge-Disjoint Paths

## Computational problem

**Input:** digraph  $G = (V, E)$ ;  $s, t \in V$

**Output:** set  $C$  of edge-disjoint  $st$ -paths in  $G$  such that  $|C|$  is maximized

## Reduction to max flow

- ▶ Integral network  $N \doteq (V, E \setminus (V \times \{s\} \cup \{t\} \times V), c \equiv 1, s, t)$

$$\begin{array}{ccc} \text{set } C \text{ of edge-disjoint } st\text{-paths in } G & \longleftrightarrow & \text{integral flow } f \text{ in } N \\ |C| & = & \nu(f) \end{array}$$

- ▶ Resulting algorithm runs in time  $O(nm)$ .



# Edge-Disjoint Paths – duality

# Edge-Disjoint Paths – duality

Capacity of a cut in  $N$

# Edge-Disjoint Paths – duality

Capacity of a cut in  $N$

►  $c(S, T) = |E \cap S \times T|$

# Edge-Disjoint Paths – duality

Capacity of a cut in  $N$

- ▶  $c(S, T) = |E \cap S \times T|$
- ▶ Removing  $E \cap S \times T$  from  $G$  ensures no  $st$ -path remains.

# Edge-Disjoint Paths – duality

## Capacity of a cut in $N$

- ▶  $c(S, T) = |E \cap S \times T|$
- ▶ Removing  $E \cap S \times T$  from  $G$  ensures no  $st$ -path remains.

## Edge connectivity duality

The maximum number of edge-disjoint  $st$ -paths equals the minimum number of edges to be removed so no  $st$ -path remains.

# Edge-Disjoint Paths – duality

## Capacity of a cut in $N$

- ▶  $c(S, T) = |E \cap S \times T|$
- ▶ Removing  $E \cap S \times T$  from  $G$  ensures no  $st$ -path remains.

## Edge connectivity duality

The maximum number of edge-disjoint  $st$ -paths equals the minimum number of edges to be removed so no  $st$ -path remains.

## Proof

# Edge-Disjoint Paths – duality

## Capacity of a cut in $N$

- ▶  $c(S, T) = |E \cap S \times T|$
- ▶ Removing  $E \cap S \times T$  from  $G$  ensures no  $st$ -path remains.

## Edge connectivity duality

The maximum number of edge-disjoint  $st$ -paths equals the minimum number of edges to be removed so no  $st$ -path remains.

## Proof

- ▶  $\ell \doteq \max$  number of edge-disjoint  $st$ -paths

# Edge-Disjoint Paths – duality

## Capacity of a cut in $N$

- ▶  $c(S, T) = |E \cap S \times T|$
- ▶ Removing  $E \cap S \times T$  from  $G$  ensures no  $st$ -path remains.

## Edge connectivity duality

The maximum number of edge-disjoint  $st$ -paths equals the minimum number of edges to be removed so no  $st$ -path remains.

## Proof

- ▶  $\ell \doteq \max$  number of edge-disjoint  $st$ -paths
- ▶  $r \doteq \min$  number of edges to be removed so no  $st$ -path



# Edge-Disjoint Paths – duality

## Capacity of a cut in $N$

- ▶  $c(S, T) = |E \cap S \times T|$
- ▶ Removing  $E \cap S \times T$  from  $G$  ensures no  $st$ -path remains.

## Edge connectivity duality

The maximum number of edge-disjoint  $st$ -paths equals the minimum number of edges to be removed so no  $st$ -path remains.

## Proof

- ▶  $\ell \doteq \max$  number of edge-disjoint  $st$ -paths
- ▶  $r \doteq \min$  number of edges to be removed so no  $st$ -path
- ▶  $\max_{\text{flow } f} f(\nu(f)) = \ell \leq r \leq \min_{st\text{-cut } (S,T)} (c(S, T))$

# Edge-Disjoint Paths – duality

## Capacity of a cut in $N$

- ▶  $c(S, T) = |E \cap S \times T|$
- ▶ Removing  $E \cap S \times T$  from  $G$  ensures no  $st$ -path remains.

## Edge connectivity duality

The maximum number of edge-disjoint  $st$ -paths equals the minimum number of edges to be removed so no  $st$ -path remains.

## Proof

- ▶  $\ell \doteq \max$  number of edge-disjoint  $st$ -paths
- ▶  $r \doteq \min$  number of edges to be removed so no  $st$ -path
- ▶  $\max_{\text{flow } f} (\nu(f)) = \ell \leq r \leq \min_{st\text{-cut } (S,T)} (c(S, T))$
- ▶ By duality, LHS = RHS so  $\ell = r$ .

# Survey Design

# Survey Design

## Computational problem

**Input:**  $n$  customers  $i \in [n]$

$m$  products  $j \in [m]$

$S_i \subseteq [m]$ : products that customer  $i \in [n]$  can survey

$c_i \in \mathbb{N}$ : max number of surveys for customer  $i \in [n]$

$p_j \in \mathbb{N}$ : min number of surveys of product  $j \in [m]$

**Output:** set  $D \subset [n] \times [m]$  such that

- $(\forall (i, j) \in D) j \in S_i$
- $(\forall i \in [n]) |\{j \in [m] : (i, j) \in D\}| \leq c_i$
- $(\forall j \in [m]) |\{i \in [n] : (i, j) \in D\}| \geq p_j$

# Survey Design

## Computational problem

**Input:**  $n$  customers  $i \in [n]$

$m$  products  $j \in [m]$

$S_i \subseteq [m]$ : products that customer  $i \in [n]$  can survey

$c_i \in \mathbb{N}$ : max number of surveys for customer  $i \in [n]$

$p_j \in \mathbb{N}$ : min number of surveys of product  $j \in [m]$

**Output:** set  $D \subset [n] \times [m]$  such that

- $(\forall (i, j) \in D) j \in S_i$
- $(\forall i \in [n]) |\{j \in [m] : (i, j) \in D\}| \leq c_i$
- $(\forall j \in [m]) |\{i \in [n] : (i, j) \in D\}| \geq p_j$

## Model

# Survey Design

## Computational problem

**Input:**  $n$  customers  $i \in [n]$

$m$  products  $j \in [m]$

$S_i \subseteq [m]$ : products that customer  $i \in [n]$  can survey

$c_i \in \mathbb{N}$ : max number of surveys for customer  $i \in [n]$

$p_j \in \mathbb{N}$ : min number of surveys of product  $j \in [m]$

**Output:** set  $D \subset [n] \times [m]$  such that


- $(\forall (i, j) \in D) j \in S_i$
- $(\forall i \in [n]) |\{j \in [m] : (i, j) \in D\}| \leq c_i$
- $(\forall j \in [m]) |\{i \in [n] : (i, j) \in D\}| \geq p_j$

## Model

Represent each  $(i, j) \in D$  as a unit of flow that passes through customer  $i$  and product  $j$ .

# Survey Design – reduction to max flow

## Survey Design – reduction to max flow



$$S_1 = \{1, 3, m\}$$



# Survey Design – analysis

# Survey Design – analysis

- ▶ Minimal survey  $D$  satisfies

$$(\forall j \in [m]) |\{i \in [m] : (i, j) \in D\}| = p_j.$$

# Survey Design – analysis

- ▶ Minimal survey  $D$  satisfies

$$(\forall j \in [m]) |\{i \in [m] : (i, j) \in D\}| = p_j.$$

- ▶ Construction of integral network  $N$  such that

$$\begin{array}{ccc} \text{minimal survey } D & \overset{\text{bijection}}{\longleftrightarrow} & \text{integral flow } f \text{ with } \nu(f) = \sum_{j \in [m]} p_j \\ D & = & \text{middle edges that carry flow} \end{array}$$

# Survey Design – analysis

- ▶ Minimal survey  $D$  satisfies

$$(\forall j \in [m]) |\{i \in [m] : (i, j) \in D\}| = p_j.$$

- ▶ Construction of integral network  $N$  such that

$$\begin{array}{ccc} \text{minimal survey } D & \xleftrightarrow{\text{bijection}} & \text{integral flow } f \text{ with } \nu(f) = \sum_{j \in [m]} p_j \\ D & = & \text{middle edges that carry flow} \end{array}$$

- ▶ Resulting algorithm runs in time  $O((n + m)(n + m + \sum_{i \in [n]} |S_i|))$ .