

Definite Integrals

page 17 of 89

PART A : A Let $f(x)$ be a continuous function defined on $[a, b]$,

$\int f(x) dx = F(x) + c$. Then $\int_a^b f(x) dx = F(b) - F(a)$ is called definite integral. This formula is known as Newton-Leibnitz formula.

Note : 1. The indefinite integral $\int f(x) dx$ is a function of x , where as definite integral $\int_a^b f(x) dx$ is a number.

2. Given $\int f(x) dx$ we can find $\int_a^b f(x) dx$, but given $\int_a^b f(x) dx$ we cannot find $\int f(x) dx$

Illustration. 1 Evaluate $\int_1^2 \frac{dx}{(x+1)(x+2)}$

Sol. $\therefore \frac{1}{(x+1)(x+2)} = \frac{1}{x+1} - \frac{1}{x+2}$ (by partial fractions)

$$\int_1^2 \frac{dx}{(x+1)(x+2)} = [\log_e(x+1) - \log_e(x+2)]_1^2$$

$$= \log_e^3 - \log_e^4 - \log_e^2 + \log_e^3 = \log_e\left(\frac{9}{8}\right)$$

Self Practice Problems

Evaluate the following

1. $\int_1^2 \frac{5x^2}{x^2 + 4x + 3} dx$ **Ans.** $5 - \frac{5}{2} \left(9\log_e \frac{5}{4} - \log_e \frac{3}{2} \right)$

2. $\int_0^{\frac{\pi}{2}} (2\sec^2 x + x^3 + 2) dx$ **Ans.** $\frac{\pi^4}{1024} + \frac{\pi}{2} + 2$

3. $\int_0^{\frac{\pi}{3}} \frac{x}{1 + \sec x} dx$ **Ans.** $\frac{\pi^2}{18} - \frac{\pi}{3\sqrt{3}} + 2\log_e\left(\frac{2}{\sqrt{3}}\right)$

PART B : Properties of definite integral

P - 1 $\int_a^b f(x) dx = \int_a^b f(t) dt$

i.e. definite integral is independent of variable of integration.

P - 2 $\int_a^b f(x) dx = - \int_b^a f(x) dx$

P - 3 $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$, where c may lie inside or outside the interval $[a, b]$.

Illustration 2 If $f(x) = \begin{cases} x+3 & : x < 3 \\ 3x^2+1 & : x \geq 3 \end{cases}$, then find $\int_2^5 f(x) dx$

Sol. $\int_2^5 f(x) dx = \int_2^3 f(x) dx + \int_3^5 f(x) dx$

$$= \int_2^3 (x+3) dx + \int_3^5 (3x^2+1) dx$$

$$= \frac{9-4}{2} + 3(3-2) + 5^3 - 3^3 + 5 - 3 = \frac{211}{2}$$

Illustration 3 Evaluate $\int_2^8 |x-5| dx$

Sol. $\int_2^8 |x-5| dx = \int_2^5 (-x+5) dx + \int_5^8 (x+5) dx = 9$

Illustration 4 Show that $\int_0^2 (2x+1) dx = \int_0^5 (2x+1) + \int_5^2 (2x+1)$

Sol. L.H.S. = $x^2 + x \Big|_0^2 = 4 + 2 = 6$
R.H.S. = $25 + 5 - 0 + (4 + 2) - (25 + 5) = 6 \quad \therefore \text{L.H.S.} = \text{R.H.S.}$

Self Practice Problems Evaluate the following

1. $\int_0^2 |x^2 + 2x - 3| dx$ **Ans.** 4

2. $\int_0^3 [x] dx$, where $[x]$ is integral part of x . **Ans.** 3

3. $\int_0^9 [\sqrt{t}] dt$ **Ans.** 13

PART C : $P - 4 \int_{-a}^a f(x) dx = \int_0^a (f(x) + f(-x)) dx$
 $= 2 \int_0^a f(x) dx$ if $f(-x) = f(x)$ i.e. $f(x)$ is even
 $= 0$ if $f(-x) = -f(x)$ i.e. $f(x)$ is odd

Illustration 5 Evaluate $\int_{-1}^1 \frac{e^x + e^{-x}}{1 + e^x} dx$

Sol. $\int_{-1}^1 \frac{e^x + e^{-x}}{1 + e^x} dx = \int_0^1 \left(\frac{e^x + e^{-x}}{1 + e^x} + \frac{e^{-x} + e^x}{1 + e^{-x}} \right) dx$
 $= \int_0^1 \left(\frac{e^x + e^{-x}}{1 + e^x} + \frac{e^x(e^{-x} + e^x)}{e^x + 1} \right) dx = \int_0^1 (e^x + e^{-x}) dx = e - 1 + \frac{(e^{-1} - 1)}{-1} = \frac{e^2 - 1}{e}$

Illustration 6 Evaluate $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos x dx$

Sol. $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \cos x dx = 2 \int_0^{\frac{\pi}{2}} \cos x dx = 2$ ($\because \cos x$ is even function)

Illustration 7 Evaluate $\int_{-1}^1 \log_e \left(\frac{2-x}{2+x} \right) dx$

Sol. Let $f(x) = \log_e \left(\frac{2-x}{2+x} \right)$
 $\Rightarrow f(-x) = \log_e \left(\frac{2+x}{2-x} \right) = -\log_e \left(\frac{2-x}{2+x} \right) = -f(x)$
i.e. $f(x)$ is odd function
 $\therefore \int_{-1}^1 \log_e \left(\frac{2-x}{2+x} \right) dx = 0$

Self Practice Problems : Evaluate the following

1. $\int_{-1}^1 |x| dx$ **Ans.** 1

2. $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^7 x dx$ **Ans.** 0

Successful People Replace the words like; "wish", "try" & "should" with "I Will". **Ineffective People** don't.

3. $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{\cos x}{1+e^x} dx$ **Ans.** 1

PART D : $P-5 \quad \int_a^b f(x) dx = \int_a^b f(a+b-x) dx$

Further $\int_0^a f(x) dx = \int_0^a f(a-x) dx$

Illustration 8 Prove that $\int_0^{\frac{\pi}{2}} \frac{g(\sin x)}{g(\sin x) + g(\cos x)} dx = \int_0^{\frac{\pi}{2}} \frac{g(\cos x)}{g(\sin x) + g(\cos x)} dx = \frac{\pi}{4}$

Sol. Let $I = \int_0^{\frac{\pi}{2}} \frac{g(\sin x)}{g(\sin x) + g(\cos x)} dx$

$\Rightarrow I = \int_0^{\frac{\pi}{2}} \frac{g\left(\sin\left(\frac{\pi}{2}-x\right)\right)}{g\left(\sin\left(\frac{\pi}{2}-x\right)\right) + g\left(\cos\left(\frac{\pi}{2}-x\right)\right)} dx = \int_0^{\frac{\pi}{2}} \frac{g(\cos x)}{g(\cos x) + g(\sin x)} dx$

on adding, we obtain

$2I = \int_0^{\frac{\pi}{2}} \left(\frac{g(\sin x)}{g(\sin x) + g(\cos x)} + \frac{g(\cos x)}{g(\cos x) + g(\sin x)} \right) dx = \int_0^{\frac{\pi}{2}} dx \Rightarrow I = \frac{\pi}{4}$

Note : 1. The above illustration can be remembered as a formula
2. Other similar formulae are

$\int_0^{\frac{\pi}{2}} \frac{g(\tan x)}{g(\tan x) + g(\cot x)} dx = \int_0^{\frac{\pi}{2}} \frac{g(\cot x)}{g(\tan x) + g(\cot x)} dx = \frac{\pi}{4}$

$\int_0^{\frac{\pi}{2}} \frac{g(\operatorname{cosec} x)}{g(\operatorname{cosec} x) + g(\sec x)} dx = \int_0^{\frac{\pi}{2}} \frac{g(\sec x)}{g(\operatorname{cosec} x) + g(\sec x)} dx = \frac{\pi}{4}$

$\int_0^a \frac{g(x)}{g(x) + g(a-x)} dx = \frac{a}{2}$

Self Practice Problems : Evaluate the following

1. $\int_0^{\frac{\pi}{2}} \frac{x}{1+\sin x} dx$ **Ans.** π

2. $\int_0^{\frac{\pi}{2}} \frac{x}{\sin x + \cos x} dx$ **Ans.** $\frac{\pi}{2\sqrt{2}} \log_e (1 + \sqrt{2})$

3. $\int_0^{\frac{\pi}{2}} \frac{x \sin x \cos x}{\sin^4 x + \cos^4 x} dx$ **Ans.** $\frac{\pi^2}{16}$

4. $\int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{dx}{1 + \sqrt{\tan x}}$ **Ans.** $\frac{\pi}{12}$

PART E : $P-6 \quad \int_0^{2a} f(x) dx = \int_0^a (f(x) + f(2a-x)) dx$

$= 2 \int_0^a f(x) dx \quad \text{if } f(2a-x) = f(x)$

$= 0 \quad \text{if } f(2a-x) = -f(x)$

Illustration 9 Evaluate $\int_0^{\pi} \sin^3 x \cos^3 x \, dx$

Sol. Let $f(x) = \sin^3 x \cos^3 x \Rightarrow f(\pi - x) = -f(x) \therefore \int_0^{\pi} \sin^3 x \cos^3 x \, dx = 0$

Illustration 10 Evaluate $\int_0^{\pi} \frac{dx}{1+2\sin^2 x}$

Sol. Let $f(x) = \frac{1}{1+2\sin^2 x} \Rightarrow f(\pi - x) = f(x)$

$$\Rightarrow \int_0^{\pi} \frac{dx}{1+2\sin^2 x} = 2 \int_0^{\frac{\pi}{2}} \frac{dx}{1+2\sin^2 x} = 2 \int_0^{\frac{\pi}{2}} \frac{\sec^2 x \, dx}{1+\tan^2 x + 2\tan^2 x}$$

$$= 2 \int_0^{\frac{\pi}{2}} \frac{\sec^2 x \, dx}{1+3\tan^2 x} = \frac{2}{\sqrt{3}} \left[\tan^{-1}(\sqrt{3} \tan x) \right]_0^{\frac{\pi}{2}}$$

$\therefore \tan \frac{\pi}{2}$ is undefined, we take limit

$$= \frac{2}{\sqrt{3}} \left[\lim_{x \rightarrow \frac{\pi}{2}^-} \tan^{-1}(\sqrt{3} \tan x) - \tan^{-1}(\sqrt{3} \tan 0) \right]$$

$$= \frac{2}{\sqrt{3}} \cdot \frac{\pi}{2} = \frac{\pi}{\sqrt{3}}$$

Note : We can evaluate the integral without using this property

Alternatively : $\int_0^{\pi} \frac{dx}{1+2\sin^2 x} = \int_0^{\pi} \frac{\operatorname{cosec}^2 x}{\operatorname{cosec}^2 x + 2} \, dx = \int_0^{\pi} \frac{\operatorname{cosec}^2 x \, dx}{\cot^2 x + 3}$

Observe that we are not converting in terms of $\tan x$ as it is not continuous in $(0, \pi)$

$$= -\frac{1}{\sqrt{3}} \left[\tan^{-1}\left(\frac{\cot x}{\sqrt{3}}\right) \right]_0^{\pi} = -\frac{1}{\sqrt{3}} \left[\lim_{x \rightarrow \pi^-} \tan^{-1}\left(\frac{\cot x}{\sqrt{3}}\right) - \lim_{x \rightarrow 0^+} \tan^{-1}\left(\frac{\cot x}{\sqrt{3}}\right) \right]$$

$$= -\frac{1}{\sqrt{3}} \left[-\frac{\pi}{2} - \frac{\pi}{2} \right] = \frac{\pi}{\sqrt{3}}$$

Note : If we convert in terms of $\tan x$, then we have to break integral using property P – 3.

Illustration 11 Prove that $\int_0^{\frac{\pi}{2}} \log_e \sin x \, dx = \int_0^{\frac{\pi}{2}} \log_e \cos x \, dx = \int_0^{\frac{\pi}{2}} \log_e (\sin 2x) \, dx = -\frac{\pi}{2} \log_e 2$.

Sol. Let $I = \int_0^{\frac{\pi}{2}} \log_e \sin x \, dx$ (i)

$$\Rightarrow I = \int_0^{\frac{\pi}{2}} \log_e \left(\sin \left(\frac{\pi}{2} - x \right) \right) \, dx \quad (\text{by property P – 5})$$

$$I = \int_0^{\frac{\pi}{2}} \log_e (\cos x) \, dx \quad \text{.....(ii)}$$

Adding (i) and (ii)

$$2I = \int_0^{\frac{\pi}{2}} \log_e (\sin x \cdot \cos x) \, dx = \int_0^{\frac{\pi}{2}} \log_e \left(\frac{\sin 2x}{2} \right) \, dx$$

$$2I = \int_0^{\frac{\pi}{2}} \log_e (\sin 2x) \, dx - \int_0^{\frac{\pi}{2}} \log_e 2 \, dx$$

$$2I = I_1 - \frac{\pi}{2} \log_e^2 \quad \dots\dots\dots(iii)$$

where $I_1 = \int_0^{\frac{\pi}{2}} \log_e(\sin 2x) dx$

put $2x = t \Rightarrow dx = \frac{1}{2} dt$
 L . L : $x = 0 \Rightarrow t = 0$

U . L : $x = \frac{\pi}{2} \Rightarrow t = \pi \Rightarrow I_1 = \int_0^{\pi} \log_e(\sin t) \frac{1}{2} dt$
 $= \frac{1}{2} \times 2 \int_0^{\frac{\pi}{2}} \log_e(\sin t) dt$ (by using property P – 6)

$\Rightarrow I_1 = I \therefore (iii) \text{ gives } I = -\frac{\pi}{2} \log_e^2$

Self Practice Problems

Evaluate the following

1. $\int_0^{\infty} \left(\frac{\log_e \left(x + \frac{1}{x} \right)}{1+x^2} \right) dx$: **Ans :** $\pi \log_e^2$

2. $\int_0^1 \frac{\sin^{-1} x}{x} dx$: **Ans :** $\frac{\pi}{2} \log_e^2$

3. $\int_0^{\pi} x \log_e \sin x dx$ **Ans :** $-\frac{\pi^2}{2} \log_e^2$

PART F : P – 7 If $f(x)$ is a periodic function with period T , then

(i) $\int_0^{nT} f(x) dx = n \int_0^T f(x) dx, n \in \mathbb{Z}$ (ii) $\int_a^{a+nT} f(x) dx = n \int_0^T f(x) dx, n \in \mathbb{Z}, a \in \mathbb{R}$

(iii) $\int_{mT}^{nT} f(x) dx = (n-m) \int_0^T f(x) dx, m, n \in \mathbb{Z}$ (iv) $\int_{nT}^{a+nT} f(x) dx = \int_0^a f(x) dx, n \in \mathbb{Z}, a \in \mathbb{R}$

(v) $\int_{a+nT}^{b+nT} f(x) dx = \int_a^b f(x) dx, n \in \mathbb{Z}, a, b \in \mathbb{R}$

Illustration 12 Evaluate $\int_{-1}^2 e^{\{x\}} dx$

Sol. $\int_{-1}^2 e^{\{x\}} dx = \int_{-1}^{-1+3} e^{\{x\}} dx = 3 \int_0^1 e^{\{x\}} dx = 3 \int_0^1 e^{\{x\}} dx = 3(e-1)$

Illustration 13 Evaluate $\int_0^{n\pi+v} |\cos x| dx, \frac{\pi}{2} < v < \pi$ and $n \in \mathbb{Z}$

Sol. $\int_0^{n\pi+v} |\cos x| dx = \int_0^v |\cos x| dx + \int_v^{n\pi+v} |\cos x| dx$

$= \int_0^{\frac{\pi}{2}} \cos x - \int_{\pi}^v \cos x dx + n \int_0^{\pi} |\cos x| dx$

$= (1-0) - (\sin v - 1) + 2n \int_0^{\frac{\pi}{2}} \cos x dx = 2 - \sin v + 2n(1-0) = 2n+2 - \sin v$

Self Practice Problem : Evaluate the following

1. $\int_{-1}^2 e^{\{3x\}} dx$ **Ans.** $3(e-1)$

2. $\int_0^{2000\pi} \frac{dx}{1+e^{\sin x}} dx$ **Ans.** 1000π

3. $\int_{\frac{5\pi}{4}}^{\frac{\pi}{4}} \frac{\sin 2x}{\sin^4 x + \cos^4 x} dx$ **Ans.** $\frac{\pi}{4}$

PART G : P-8 If $\psi(x) \leq f(x) \leq \phi(x)$ for $a \leq x \leq b$, then

$$\int_a^b \psi(x) dx \leq \int_a^b f(x) dx \leq \int_a^b \phi(x) dx$$

P-9 If $m \leq f(x) \leq M$ for $a \leq x \leq b$, then $m(b-a) \leq \int_a^b f(x) dx \leq M(b-a)$

Further if $f(x)$ is monotonically decreasing in (a, b) then $f(b)(b-a) < \int_a^b f(x) dx < f(a)(b-a)$ and if $f(x)$

is monotonically increasing in (a, b) then $f(a)(b-a) < \int_a^b f(x) dx < f(b)(b-a)$

P-10 $\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx$

P-11 If $f(x) \geq 0$ on $[a, b]$ then $\int_a^b f(x) dx \geq 0$

Illustration 14 For $x \in (0, 1)$ arrange $f_1(x) = \frac{1}{\sqrt{4-x^2}}$, $f_2(x) = \frac{1}{\sqrt{4-2x^2}}$ and $f_3(x) = \frac{1}{\sqrt{4-x^2-x^3}}$ in ascending

order and hence prove that $\frac{\pi}{6} < \int_0^1 \frac{dx}{\sqrt{4-x^2-x^3}} < \frac{\pi}{4\sqrt{2}}$

Sol.

$$\Rightarrow \because 0 < x^3 < x^2 \Rightarrow \frac{x^2}{4-2x^2} < \frac{x^2}{4-x^2-x^3} < \frac{x^2}{4-x^2} \Rightarrow -2x^2 < -x^2-x^3 < -x^2$$

$$\Rightarrow \sqrt{4-2x^2} < \sqrt{4-x^2-x^3} < \sqrt{4-x^2} \Rightarrow f_1(x) < f_3(x) < f_2(x) \text{ for } x \in (0, 1)$$

$$\Rightarrow \int_0^1 f_1(x) dx < \int_0^1 f_3(x) dx < \int_0^1 f_2(x) dx$$

$$\sin^{-1} \left(\frac{x}{2} \right) \Big|_0^1 < \int_0^1 \frac{dx}{\sqrt{4-x^2-x^3}} < \frac{1}{\sqrt{2}} \sin^{-1} \frac{x}{\sqrt{2}} \Big|_0^1$$

$$\frac{\pi}{6} < \int_0^1 \frac{dx}{\sqrt{4-x^2-x^3}} < \frac{\pi}{4\sqrt{2}}$$

Illustration 15 Estimate the value of $\int_0^{\frac{\pi}{2}} \frac{\sin x}{x} dx$

Sol.

$$\text{Let } f(x) = \frac{\sin x}{x}$$

$$f'(x) = \frac{x \cos x - \sin x}{x^2} = \frac{(\cos x)(x - \tan x)}{x^2} < 0$$

\Rightarrow $f(x)$ is monotonically decreasing function.
 $f(0)$ is not defined, so we evaluate

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \frac{\sin x}{x} = 1. \text{ Take } f(0) = \lim_{x \rightarrow 0^+} f(x) = 1$$

$$f\left(\frac{\pi}{2}\right) = \frac{2}{\pi}$$

$$\frac{2}{\pi} \cdot \left(\frac{\pi}{2} - 0 \right) < \int_0^{\frac{\pi}{2}} \frac{\sin x}{x} dx < 1 \cdot \left(\frac{\pi}{2} - 0 \right)$$

$$1 < \int_0^{\frac{\pi}{2}} \frac{\sin x}{x} dx < \frac{\pi}{2}$$

Note : Here by making the use of graph we can make more appropriate approximation as in next illustration.


Illustration 16 Estimate the value of $\int_0^1 e^{x^2} dx$ using (i) rectangle, (ii) triangle

Sol. (i) By using rectangle

$$\text{Area OAED} < \int_0^1 e^{x^2} dx < \text{Area OABC}$$

$$1 < \int_0^1 e^{x^2} dx < 1 \cdot e$$

$$1 < \int_0^1 e^{x^2} dx < e$$



(ii) By using triangle

$$\text{Area OAED} < \int_0^1 e^{x^2} dx < \text{Area OAED} + \text{Area of triangle DEB}$$


$$1 < \int_0^1 e^{x^2} dx < 1 + \frac{1}{2} \cdot 1 \cdot (e - 1) \quad 1 < \int_0^1 e^{x^2} dx < \frac{e+1}{2}$$

Illustration 17 Estimate the value of $\int_0^1 e^{x^2} dx$ by using $\int_0^1 e^x dx$

Sol. For $x \in (0, 1)$, $e^{x^2} < e^x$

$$\Rightarrow 1 \times 1 < \int_0^1 e^{x^2} dx < \int_0^1 e^x dx$$

$$1 < \int_0^1 e^{x^2} dx < e - 1$$



Exercise : Prove the following :

- $\int_0^1 e^{-x} \cos^2 x dx < \int_0^1 e^{-x^2} \cos^2 x dx$
- $0 < \int_0^{\frac{\pi}{2}} \sin^{n+1} x dx < \int_0^{\frac{\pi}{2}} \sin^2 x dx$
- $e^{-\frac{1}{4}} < \int_0^1 e^{x^2-x} dx < 1$
- $-\frac{1}{2} \leq \int_0^1 \frac{x^3 \cos x}{2+x^2} dx < \frac{1}{2}$
- $1 < \int_0^{\frac{\pi}{2}} \sqrt{\sin x} dx < \sqrt{\frac{\pi}{2}}$
- $0 < \int_0^2 \frac{x dx}{16+x^3} < \frac{1}{6}$

PART - H: Leibnitz Theorem : If $F(x) = \int_{g(x)}^{h(x)} f(t) dt$, then

Successful People Replace the words like; "wish", "try" & "should" with "I Will". Ineffective People don't.

$$\frac{dF(x)}{dx} = h'(x) f(h(x)) - g'(x) f(g(x))$$

Proof :

$$\text{Let } P(t) = \int f(t) dt$$

$$\Rightarrow F(x) = \int_{g(x)}^{h(x)} f(t) dt = P(h(x)) - P(g(x)) \Rightarrow \frac{dF(x)}{dx} = P'(h(x)) h'(x) - P'(g(x)) g'(x) = f(h(x)) h'(x) - f(g(x)) g'(x)$$

Illustration 18 If $F(x) = \int_x^{x^2} \sqrt{\sin t} dt$, then find $F'(x)$

Sol. $F'(x) = 2x \cdot \sqrt{\sin x^2} - 1 \cdot \sqrt{\sin x}$

Illustration 19 If $F(x) = \int_{e^{2x}}^{e^{3x}} \frac{t}{\log_e t} dt$, then find first and second derivative of $F(x)$ with respect to $\log_e x$.

at $x = \log_e^2$

Sol. $\frac{dF(x)}{d(\log_e^x)} = \frac{dF(x)}{dx} \cdot \frac{dx}{d(\log_e^x)} = \left[3 \cdot e^{3x} \cdot \frac{e^{3x}}{\log_e^{3x}} - 2 \cdot e^{2x} \cdot \frac{e^{2x}}{\log_e^{2x}} \right] x = e^{6x} - e^{4x}.$

$$\frac{d^2 F(x)}{d(\log_e^x)^2} = \frac{d}{d(\log_e^x)} (e^{6x} - e^{4x}) = \frac{d}{dx} (e^{6x} - e^{4x}) \times \frac{1}{\frac{dx}{d(\log_e^x)}} = (6e^{6x} - 4e^{4x}) x$$

First derivative of $F(x)$ at $x = \log_e^2$ (i.e. $e^x = 2$) is $2^6 - 2^4 = 48$
Second derivative of $F(x)$ at $x = \log_e^2$ (i.e. $e^x = 2$) is $(6 \cdot 2^6 - 4 \cdot 2^4) \cdot \log_e^2 = 5 \cdot 2^6 \cdot \log_e^2$.

Illustration 20 Evaluate $\lim_{x \rightarrow \infty} \frac{\left(\int_0^x e^{t^2} dt \right)^2}{\int_0^x e^{2t^2} dt}$

Sol. $\lim_{x \rightarrow \infty} \frac{\left(\int_0^x e^{t^2} dt \right)^2}{\int_0^x e^{2t^2} dt} \quad \left(\frac{\infty}{\infty} \text{ form} \right)$

Applying L' Hospital rule

$$= \lim_{x \rightarrow \infty} \frac{2 \cdot \int_0^x e^{t^2} dt \cdot e^{x^2}}{1 \cdot e^{2x^2}} = \lim_{x \rightarrow \infty} \frac{2 \cdot \int_0^x e^{t^2} dt}{e^{x^2}} = \lim_{x \rightarrow \infty} \frac{2 \cdot e^{x^2}}{2x \cdot e^{x^2}} = 0$$

Modified Leibnitz Theorem :

If $F(x) = \int_{g(x)}^{h(x)} f(x, t) dt$, then

$$F'(x) = \int_{g(x)}^{h(x)} \frac{\partial f(x, t)}{\partial x} dt + f(x, h(x)) h'(x) - f(x, g(x)) \cdot g'(x)$$

Illustration 21 If $f(x) = \int_{\log_e^x}^x \frac{dt}{x+t}$, then find $f'(x)$

Sol. $f'(x) = \int_{\log_e^x}^x \frac{-1}{(x+t)^2} dt + 1 \cdot \frac{1}{2x} - \frac{1}{x} \cdot \frac{1}{(x+\log_e^x)} = \frac{1}{(x+t)} \Big|_{\log_e^x}^x + \frac{1}{2x} - \frac{1}{x(x+\log_e^x)}$
 $= \frac{1}{2x} - \frac{1}{x+\log_e^x} + \frac{1}{2x} - \frac{1}{x(x+\log_e^x)} = \frac{1}{x} - \frac{x+1}{x(x+\log_e^x)} = \frac{\log_e^x - 1}{x(x+\log_e^x)}$

Alternatively : $f(x) = \int_{\log_e^x}^x \frac{dt}{x+t} = \log_e(x+t) \Big|_{\log_e^x}^x$ (treating 't' as constant)

$$f(x) = \log_e^{2x} - \log_e(x + \log_e^x)$$

$$f'(x) = \frac{1}{x} - \frac{1}{(x + \log_e^x)} \left(1 + \frac{1}{x}\right) = \frac{\log_e^x - 1}{x(x + \log_e^x)}$$

Definite Integrals dependent on parameters :

Illustration 23 Evaluate $\int_0^1 \frac{x^b - 1}{\log_e^x} dx$, 'b' being parameter

Sol. Let $I(b) = \int_0^1 \frac{x^b - 1}{\log_e^x} dx$

$$\frac{dI(b)}{db} = \int_0^1 \frac{x^b \log_e^x}{\log_e^x} dx + 0 - 0$$

(using modified Leibnitz Theorem)

$$= \int_0^1 x^b dx = \left[\frac{x^{b+1}}{b+1} \right]_0^1 = \frac{1}{b+1}$$

$$I(b) = \log_e(b+1) + c$$

$$b=0 \Rightarrow I(0) = 0$$

$$\therefore c = 0 \therefore I(b) = \log_e(b+1)$$

Illustration 24 Evaluate $\int_0^1 \frac{\tan^{-1}(ax)}{x\sqrt{1-x^2}} dx$, 'a' being parameter

Sol. Let $I(a) = \int_0^1 \frac{\tan^{-1}(ax)}{x\sqrt{1-x^2}} dx$

$$\frac{dI(a)}{da} = \int_0^1 \frac{x}{(1+a^2x^2)} \cdot \frac{1}{x\sqrt{1-x^2}} dx = \int_0^1 \frac{dx}{(1+a^2x^2)\sqrt{1-x^2}}$$

$$\text{Put } x = \sin t \Rightarrow dx = \cos t dt$$

$$\text{L.L. : } x=0 \Rightarrow t=0$$

$$\text{U.L. : } x=1 \Rightarrow t = \frac{\pi}{2}$$

$$\frac{dI(a)}{da} = \int_0^{\frac{\pi}{2}} \frac{1}{1+a^2 \sin^2 t} \cdot \frac{1}{\cos t} \cos t dt = \int_0^{\frac{\pi}{2}} \frac{dt}{1+a^2 \sin^2 t}$$

$$= \int_0^{\frac{\pi}{2}} \frac{\sec^2 t dt}{1+(1+a^2) \tan^2 t} = \frac{1}{\sqrt{1+a^2}} \tan^{-1} \left(\sqrt{1+a^2} \tan t \right) \Big|_0^{\frac{\pi}{2}}$$

$$= \frac{1}{\sqrt{1+a^2}} \cdot \frac{\pi}{2}$$

$$\Rightarrow I(a) = \frac{\pi}{2} \log_e \left(a + \sqrt{1+a^2} \right) + c$$

$$\text{But } I(0) = 0 \Rightarrow c = 0 \Rightarrow I(a) = \frac{\pi}{2} \log_e \left(a + \sqrt{1+a^2} \right)$$

Self Practice Problems :

1. If $f(x) = \int_0^{x^3} \sqrt{\cos t} dt$, find $f'(x)$. **Ans.** $3x^2 \sqrt{\cos x^3}$

2. If $f(x) = e^{g(x)}$ and $g(x) = \int_2^x \frac{t}{1+t^4} dt$ then find the value of $f'(2)$. **Ans.** $\frac{2}{17}$

3. If $x = \int_0^y \frac{dt}{\sqrt{1+4t^2}}$ and $\frac{d^2y}{dx^2} = Ry$ then find R **Ans.** 4

4. If $f(x) = \int_x^{x^2} \sin t \, dt$ then find $f'(x)$. **Ans.** $x^2 (2x \sin x^2 - \sin x) + (\cos x - \cos x^2) x$

5. If $\phi(x) = \cos x - \int_0^x (x-t) \phi(t) \, dt$, then find the value of $\phi''(x) + \phi(x)$. **Ans.** $-\cos x$


6. Find the value of the function $f(x) = 1 + x + \int_1^x ((\log_e t)^2 + 2 \log_e t) \, dt$ where $f'(x)$ vanishes. **Ans.** $1 + \frac{2}{e}$

7. Evaluate $\lim_{x \rightarrow 0} \frac{\int_0^{x^2} \cos t^2 \, dt}{x \sin x}$. **Ans.** 1

8. Evaluate $\int_0^\pi \log_e (1 + b \cos x) \, dx$, 'b' being parameter. **Ans.** $\pi \log_e \left(\frac{1 + \sqrt{1-b^2}}{2} \right)$

PART - I: Definite Integral as a Limit of Sum.

Let $f(x)$ be a continuous real valued function defined on the closed interval $[a, b]$ which is divided into n parts as shown in figure.



The point of division on x-axis are $a, a + h, a + 2h, \dots, a + (n-1)h, a + nh$, where $\frac{b-a}{n} = h$.

Let S_n denotes the area of these n rectangles.

Then, $S_n = hf(a) + hf(a+h) + hf(a+2h) + \dots + hf(a+(n-1)h)$

Clearly, S_n is area very close to the area of the region bounded by curve $y = f(x)$, x-axis and the ordinates $x = a, x = b$.

Hence $\int_a^b f(x) \, dx = \lim_{n \rightarrow \infty} S_n$

$$\int_a^b f(x) \, dx = \lim_{n \rightarrow \infty} \sum_{r=0}^{n-1} h f(a+rh) = \lim_{n \rightarrow \infty} \sum_{r=0}^{n-1} \left(\frac{b-a}{n} \right) f \left(a + \frac{(b-a)r}{n} \right)$$

Note : 1. We can also write

$$S_n = hf(a+h) + hf(a+2h) + \dots + hf(a+nh) \text{ and } \int_a^b f(x) \, dx = \lim_{n \rightarrow \infty} \sum_{r=1}^n \left(\frac{b-a}{n} \right) f \left(a + \left(\frac{b-a}{n} \right) r \right)$$

2. If $a = 0, b = 1, \int_0^1 f(x) \, dx = \lim_{n \rightarrow \infty} \sum_{r=0}^{n-1} \frac{1}{n} f \left(\frac{r}{n} \right)$

Steps to express the limit of sum as definite integral

Step 1. Replace $\frac{r}{n}$ by $x, \frac{1}{n}$ by dx and $\lim_{n \rightarrow \infty} \sum$ by \int

Step 2. Evaluate $\lim_{n \rightarrow \infty} \left(\frac{r}{n} \right)$ by putting least and greatest values of r as lower and upper limits respectively.

For example $\lim_{n \rightarrow \infty} \sum_{r=1}^{pn} \frac{1}{n} f \left(\frac{r}{n} \right) = \int_0^p f(x) \, dx \quad \left(\because \lim_{n \rightarrow \infty} \left(\frac{r}{n} \right) \Big|_{r=1} = 0, \lim_{n \rightarrow \infty} \left(\frac{r}{n} \right) \Big|_{r=np} = p \right)$

Illustration 25 : Evaluate

$$\lim_{n \rightarrow \infty} \left[\frac{1}{1+n} + \frac{1}{2+n} + \frac{1}{3+n} + \dots + \frac{1}{2n} \right]$$

Sol. $\lim_{n \rightarrow \infty} \left[\frac{1}{1+n} + \frac{1}{2+n} + \frac{1}{3+n} + \dots + \frac{1}{2n} \right] = \lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{1}{r+n}$

$$= \lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{1}{n} \frac{1}{\left(\frac{r}{n}\right)+1} = \int_0^1 \frac{dx}{x+1} = [\log_e(x+1)]_0^1 = \log_e 2.$$

Illustration 26 : Evaluate $\lim_{n \rightarrow \infty} \left[\frac{n+1}{n^2+1^2} + \frac{n+2}{n^2+2^2} + \frac{n+3}{n^2+3^2} + \dots + \frac{3}{5n} \right]$

Sol. $\lim_{n \rightarrow \infty} \sum_{r=1}^{2n} \frac{n+r}{n^2+r^2} = \lim_{n \rightarrow \infty} \sum_{r=1}^{2n} \frac{1}{n} \frac{1+\frac{r}{n}}{1+\left(\frac{r}{n}\right)^2}$

$\therefore \lim_{n \rightarrow \infty} \left(\frac{r}{n}\right) = 0$, when $r = 1$, lower limit = 0

and $\lim_{n \rightarrow \infty} \left(\frac{r}{n}\right) = \lim_{n \rightarrow \infty} \left(\frac{2n}{n}\right) = 2$, when $r = 2n$, upper limit = 2

$$\int_0^2 \frac{1+x}{1+x^2} dx = \int_0^2 \frac{1}{1+x^2} dx + \frac{1}{2} \int_0^2 \frac{2x}{1+x^2} dx$$

$$= \tan^{-1}x \Big|_0^2 + \frac{1}{2} \log_e(1+x^2) \Big|_0^2 = \tan^{-1} 2 + \frac{1}{2} \log_e 5$$

Illustration 27 : Evaluate

$$\lim_{n \rightarrow \infty} \left(\frac{n!}{n^n}\right)^{\frac{1}{n}}$$

Sol. Let $y = \lim_{n \rightarrow \infty} \left(\frac{n!}{n^n}\right)^{\frac{1}{n}}$

$$\begin{aligned} \log_e y &= \lim_{n \rightarrow \infty} \frac{1}{n} \log_e \left(\frac{n!}{n^n}\right) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \log_e \left(\frac{1 \cdot 2 \cdot 3 \dots n}{n^n}\right) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \left[\log_e \left(\frac{1}{n}\right) + \log_e \left(\frac{2}{n}\right) + \log_e \left(\frac{3}{n}\right) + \dots + \log_e \left(\frac{n}{n}\right) \right] \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=1}^n \log_e \left(\frac{r}{n}\right) \\ &= \int_0^1 \log_e x dx = x \log_e x - x \Big|_0^1 \\ &= (0-1) - \lim_{x \rightarrow 0^+} x \log_e x + 0 \\ &= -1 - 0 = -1 \Rightarrow y = \frac{1}{e} \end{aligned}$$

Self Practice Problems :

Evaluate the following limits

1. $\lim_{n \rightarrow \infty} \left[\frac{1}{\sqrt{n^2}} + \frac{1}{\sqrt{n^2+n}} + \frac{1}{\sqrt{n^2+2n}} + \dots + \frac{1}{\sqrt{n^2+n^2}} \right]$

Ans. $2(\sqrt{2}-1)$

2. $\lim_{n \rightarrow \infty} \left[\frac{1}{1+n} + \frac{1}{2+n} + \frac{1}{3+n} + \dots + \frac{1}{5n} \right]$

Ans. $\log_e 5$

3. $\lim_{n \rightarrow \infty} \frac{1}{n^2} \left[\sin^3 \frac{\pi}{4n} + 2 \sin^3 \frac{2\pi}{4n} + 3 \sin^3 \frac{3\pi}{4n} + \dots + n \sin^3 \frac{n\pi}{4n} \right]$

Ans. $\frac{\sqrt{2}}{9\pi^2} (52 - 15\pi)$

4. $\lim_{n \rightarrow \infty} \sum_{r=0}^{n-1} \frac{1}{\sqrt{n^2-r^2}}$

Ans. $\frac{\pi}{2}$

5. $\lim_{n \rightarrow \infty} \frac{3}{n} \left[1 + \sqrt{\frac{n}{n+3}} + \sqrt{\frac{n}{n+6}} + \sqrt{\frac{n}{n+9}} + \dots + \sqrt{\frac{n}{n+3(n-1)}} \right]$ Ans. 2

PART – J: Reduction Formulae in Definite Integrals

1. If $I_n = \int_0^{\frac{\pi}{2}} \sin^n x \, dx$, then show that $I_n = \left(\frac{n-1}{n} \right) I_{n-2}$

Proof : $I_n = \int_0^{\frac{\pi}{2}} \sin^n x \, dx$

$$I_n = \left[-\sin^{n-1} x \cos x \right]_0^{\frac{\pi}{2}} + \int_0^{\frac{\pi}{2}} (n-1) \sin^{n-2} x \cdot \cos^2 x \, dx$$

$$= (n-1) \int_0^{\frac{\pi}{2}} \sin^{n-2} x \cdot (1 - \sin^2 x) \, dx$$

$$= (n-1) \int_0^{\frac{\pi}{2}} \sin^{n-2} x \, dx - (n-1) \int_0^{\frac{\pi}{2}} \sin^n x \, dx$$

$$I_n + (n-1) I_n = (n-1) I_{n-2}$$

$$I_n = \left(\frac{n-1}{n} \right) I_{n-2}$$

Note : 1. $\int_0^{\frac{\pi}{2}} \sin^n x \, dx = \int_0^{\frac{\pi}{2}} \cos^n x \, dx$

2. $I_n = \left(\frac{n-1}{n} \right) \left(\frac{n-3}{n-2} \right) \left(\frac{n-5}{n-4} \right) \dots I_0$ or I_1

according as n is even or odd. $I_0 = \frac{\pi}{2}$, $I_1 = 1$

Hence $I_n = \begin{cases} \left(\frac{n-1}{n} \right) \left(\frac{n-3}{n-2} \right) \left(\frac{n-5}{n-4} \right) \dots \left(\frac{1}{2} \right) \cdot \frac{\pi}{2} & \text{if } n \text{ is even} \\ \left(\frac{n-1}{n} \right) \left(\frac{n-3}{n-2} \right) \left(\frac{n-5}{n-4} \right) \dots \left(\frac{2}{3} \right) \cdot 1 & \text{if } n \text{ is odd} \end{cases}$

2. If $I_n = \int_0^{\frac{\pi}{4}} \tan^n x \, dx$, then show that $I_n + I_{n-2} = \frac{1}{n-1}$

Sol. $I_n = \int_0^{\frac{\pi}{4}} (\tan x)^{n-2} \cdot \tan^2 x \, dx$

$$= \int_0^{\frac{\pi}{4}} (\tan x)^{n-2} (\sec^2 x - 1) \, dx = \int_0^{\frac{\pi}{4}} (\tan x)^{n-2} \sec^2 x \, dx - \int_0^{\frac{\pi}{4}} (\tan x)^{n-2} \, dx$$

$$= \left[\frac{(\tan x)^{n-1}}{n-1} \right]_0^{\frac{\pi}{4}} - I_{n-2}$$

$$I_n = \frac{1}{n-1} - I_{n-2} \quad \therefore \quad I_n + I_{n-2} = \frac{1}{n-1}$$

3. If $I_{m,n} = \int_0^{\frac{\pi}{2}} \sin^m x \cdot \cos^n x \, dx$, then show that $I_{m,n} = \frac{m-1}{m+n} I_{m-2, n}$, n

Sol.
$$I_{m,n} = \int_0^{\frac{\pi}{2}} \sin^{m-1} x (\sin x \cos^n x) dx$$

$$= \left[-\frac{\sin^{m-1} x \cdot \cos^{n+1} x}{n+1} \right]_0^{\frac{\pi}{2}} + \int_0^{\frac{\pi}{2}} \frac{\cos^{n+1} x}{n+1} (m-1) \sin^{m-2} x \cos x dx$$

$$= \left(\frac{m-1}{n+1} \right) \int_0^{\frac{\pi}{2}} \sin^{m-2} x \cdot \cos^n x \cdot \cos^2 x dx$$

$$= \left(\frac{m-1}{n+1} \right) \int_0^{\frac{\pi}{2}} (\sin^{m-2} x \cdot \cos^n x - \sin^m x \cdot \cos^n x) dx$$

$$= \left(\frac{m-1}{n+1} \right) I_{m-2,n} - \left(\frac{m-1}{n+1} \right) I_{m,n}$$

$$\Rightarrow \left(1 + \frac{m-1}{n+1} \right) I_{m,n} = \left(\frac{m-1}{n+1} \right) I_{m-2,n}$$

$$I_{m,n} = \left(\frac{m-1}{m+n} \right) I_{m-2,n}$$

Note : 1. $I_{m,n} = \left(\frac{m-1}{m+n} \right) \left(\frac{m-3}{m+n-2} \right) \left(\frac{m-5}{m+n-4} \right) \dots \dots I_{0,n}$ or $I_{1,n}$ according as m is even or odd.

$$I_{0,n} = \int_0^{\frac{\pi}{2}} \cos^n x dx \text{ and } I_{1,n} = \int_0^{\frac{\pi}{2}} \sin x \cdot \cos^n x dx = \frac{1}{n+1}$$

2. Walli's Formula

$$I_{m,n} = \begin{cases} \frac{(n-1)(n-3)(m-5) \dots (n-1)(n-3)(n-5) \dots \pi}{(m+n)(m+n-2)(m+n-4) \dots} \cdot \frac{\pi}{2} & \text{when both } m, n \text{ are even} \\ \frac{(m-1)(m-3)(m-5) \dots (n-1)(n-3)(n-5) \dots}{(m+n)(m+n-2)(m+n-4) \dots} & \text{otherwise} \end{cases}$$

Illustration 28 : Evaluate $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^2 x \cos^2 x (\sin x + \cos x) dx$

Sol. Given integral = $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^3 x \cos^2 x dx + \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^2 x \cos^3 x dx$

$$= 0 + 2 \int_0^{\frac{\pi}{2}} \sin^2 x \cos^3 x dx \quad (\because \sin^3 x \cos^2 x \text{ is odd and } \sin^2 x \cos^3 x \text{ is even})$$

$$= 2 \cdot \frac{1 \cdot 2}{5 \cdot 3 \cdot 1} = \frac{4}{15}$$

Illustration 29 : Evaluate $\int_0^{\pi} x \sin^5 x \cos^6 x dx$

Sol. Let $I = \int_0^{\pi} x \sin^5 x \cos^6 x dx$

$$I = \int_0^{\pi} (\pi - x) \sin^5 (\pi - x) \cos^6 (\pi - x) dx$$

$$= \pi \int_0^{\pi} \sin^5 x \cdot \cos^6 x dx - \int_0^{\pi} x \sin^5 x \cdot \cos^6 x dx$$

$$\Rightarrow 2I = \pi \cdot 2 \int_0^{\frac{\pi}{2}} \sin^5 x \cdot \cos^6 x \, dx$$

$$I = \pi \frac{4 \cdot 2 \cdot 5 \cdot 3 \cdot 1}{11 \cdot 9 \cdot 7 \cdot 5 \cdot 3 \cdot 1}$$

$$I = \frac{8\pi}{693}$$

Illustration 30 : Evaluate $\int_0^1 x^3(1-x)^5 \, dx$

Sol. Put $x = \sin^2 \theta \Rightarrow dx = 2 \sin \theta \cos \theta \, d\theta$
 L.L. : $x = 0 \Rightarrow \theta = 0$

U.L. : $x = 1 \Rightarrow \theta = \frac{\pi}{2}$

$$\therefore \int_0^1 x^3(1-x)^5 \, dx = \int_0^{\frac{\pi}{2}} \sin^6 \theta (\cos^2 \theta)^5 \cdot 2 \sin \theta \cdot \cos \theta \, d\theta$$

$$= 2 \cdot \int_0^{\frac{\pi}{2}} \sin^7 \theta \cos^{11} \theta \, d\theta$$

$$= 2 \cdot \frac{6 \cdot 4 \cdot 2 \cdot 10 \cdot 8 \cdot 6 \cdot 4 \cdot 2}{18 \cdot 16 \cdot 14 \cdot 12 \cdot 10 \cdot 8 \cdot 6 \cdot 4 \cdot 2} = \frac{1}{504}$$

Self Practice Problems:

Evaluate the following

1. $\int_0^{\frac{\pi}{2}} \sin^5 x \, dx$

Ans. $\frac{18}{15}$

2. $\int_0^{\frac{\pi}{2}} \sin^5 x \cos^4 x \, dx$

Ans. $\frac{8}{315}$

3. $\int_0^1 x^6 \sin^{-1} x \, dx$

Ans. $\frac{\pi}{14} - \frac{16}{245}$

4. $\int_0^a x (a^2 - x^2)^{\frac{7}{2}} \, dx$

Ans. $\frac{a^9}{9}$

5. $\int_0^2 x^{3/2} \sqrt{2-x} \, dx$

Ans. $\frac{\pi}{2}$