

11. Linear Combinations:

Given a finite set of vectors $\vec{a}, \vec{b}, \vec{c}, \dots$ then the vector $\vec{r} = x\vec{a} + y\vec{b} + z\vec{c} + \dots$ is called a linear combination of $\vec{a}, \vec{b}, \vec{c}, \dots$ for any $x, y, z, \dots \in \mathbb{R}$. We have the following results:

- (a) If \vec{a}, \vec{b} are non zero, non-collinear vectors then $x\vec{a} + y\vec{b} = x'\vec{a} + y'\vec{b} \Rightarrow x = x'; y = y'$
- (b) **Fundamental Theorem:** Let \vec{a}, \vec{b} be non zero, non collinear vectors. Then any vector \vec{r} coplanar with \vec{a}, \vec{b} can be expressed uniquely as a linear combination of \vec{a}, \vec{b}
i.e. There exist some uniquely $x, y \in \mathbb{R}$ such that $x\vec{a} + y\vec{b} = \vec{r}$.
- (c) If $\vec{a}, \vec{b}, \vec{c}$ are non-zero, non-coplanar vectors then:

$$x\vec{a} + y\vec{b} + z\vec{c} = x'\vec{a} + y'\vec{b} + z'\vec{c} \Rightarrow x = x', y = y', z = z'$$

- (d) **Fundamental Theorem In Space:** Let $\vec{a}, \vec{b}, \vec{c}$ be non-zero, non-coplanar vectors in space. Then any vector \vec{r} , can be uniquely expressed as a linear combination of $\vec{a}, \vec{b}, \vec{c}$ i.e. There exist some unique $x, y \in \mathbb{R}$ such that $x\vec{a} + y\vec{b} + z\vec{c} = \vec{r}$.

- (e) If $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n$ are n non zero vectors, & k_1, k_2, \dots, k_n are n scalars & if the linear combination $k_1\vec{x}_1 + k_2\vec{x}_2 + \dots + k_n\vec{x}_n = 0 \Rightarrow k_1 = 0, k_2 = 0, \dots, k_n = 0$ then we say that vectors $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n$ are **LINEARLY INDEPENDENT VECTORS**.

- (f) If $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n$ are not **LINEARLY INDEPENDENT** then they are said to be **LINEARLY DEPENDENT** vectors. i.e. if $k_1\vec{x}_1 + k_2\vec{x}_2 + \dots + k_n\vec{x}_n = 0$ & if there exists at least one $k_i \neq 0$ then $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_n$ are said to be

LINEARLY DEPENDENT.

Note 1: If $k_r \neq 0$; $k_1\vec{x}_1 + k_2\vec{x}_2 + k_3\vec{x}_3 + \dots + k_r\vec{x}_r + \dots + k_n\vec{x}_n = 0$

$$-k_r\vec{x}_r = k_1\vec{x}_1 + k_2\vec{x}_2 + \dots + k_{r-1}\vec{x}_{r-1} + k_{r+1}\vec{x}_{r+1} + \dots + k_n\vec{x}_n$$

$$-k_r \frac{1}{k_r} \vec{x}_r = k_1 \frac{1}{k_r} \vec{x}_1 + k_2 \frac{1}{k_r} \vec{x}_2 + \dots + k_{r-1} \frac{1}{k_r} \vec{x}_{r-1} + \dots + k_n \frac{1}{k_r} \vec{x}_n$$

$$\vec{x}_r = c_1\vec{x}_1 + c_2\vec{x}_2 + \dots + c_{r-1}\vec{x}_{r-1} + c_{r+1}\vec{x}_{r+1} + \dots + c_n\vec{x}_n$$

i.e. \vec{x}_r is expressed as a linear combination of vectors.

$$\vec{x}_1, \vec{x}_2, \dots, \vec{x}_{r-1}, \vec{x}_{r+1}, \dots, \vec{x}_n$$

Hence \vec{x}_r with $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_{r-1}, \vec{x}_{r+1}, \dots, \vec{x}_n$ forms a linearly dependent set of vectors.

Note 2: If $\vec{a} = 3\hat{i} + 2\hat{j} + 5\hat{k}$ then \vec{a} is expressed as a **LINEAR COMBINATION** of vectors $\hat{i}, \hat{j}, \hat{k}$ Also, $\vec{a}, \hat{i}, \hat{j}, \hat{k}$ form a linearly dependent set of vectors. In general, every set of four vectors is a linearly dependent system.

$\hat{i}, \hat{j}, \hat{k}$ are **Linearly Independent** set of vectors. For $K_1\hat{i} + K_2\hat{j} + K_3\hat{k} = 0 \Rightarrow K_1 = K_2 = K_3 = 0$

Two vectors \vec{a} & \vec{b} are linearly dependent $\Rightarrow \vec{a}$ is parallel to \vec{b} i.e. $\vec{a} \times \vec{b} = 0 \Rightarrow$ linear dependence of \vec{a} & \vec{b} . Conversely if $\vec{a} \times \vec{b} \neq 0$ then \vec{a} & \vec{b} are linearly independent.

If three vectors $\vec{a}, \vec{b}, \vec{c}$ are linearly dependent, then they are coplanar i.e. $[\vec{a}, \vec{b}, \vec{c}] = 0$, conversely, if $[\vec{a}, \vec{b}, \vec{c}] \neq 0$, then the vectors are linearly independent.

Solved Example: Given A that the points $\vec{a} - 2\vec{b} + 3\vec{c}, 2\vec{a} + 3\vec{b} - 4\vec{c}, -7\vec{b} + 10\vec{c}$, A, B, C have position vector prove that vectors \vec{AB} and \vec{AC} are linearly dependent.

Solution. Let A, B, C be the given points and O be the point of reference then

$$\vec{OA} = \vec{a} - 2\vec{b} + 3\vec{c}, \vec{OB} = 2\vec{a} + 3\vec{b} - 4\vec{c} \quad \text{and} \quad \vec{OC} = -7\vec{b} + 10\vec{c}$$

Now $\vec{AB} = \text{p.v. of B} - \text{p.v. of A}$

$$= \vec{OB} - \vec{OA} = (2\vec{a} + 3\vec{b} - 4\vec{c}) - (\vec{a} - 2\vec{b} + 3\vec{c}) = \vec{a} + 5\vec{b} - 7\vec{c}$$

$\therefore \vec{AC} = \lambda \vec{AB}$ where $\lambda = -1$. Hence \vec{AB} and \vec{AC} are linearly dependent

Solved Example: Prove that the vectors $5\vec{a} + 6\vec{b} + 7\vec{c}, 7\vec{a} - 8\vec{b} + 9\vec{c}$ and $3\vec{a} + 20\vec{b} + 5\vec{c}$ are linearly dependent $\vec{a}, \vec{b}, \vec{c}$ being linearly independent vectors.

Solution. We know that if these vectors are linearly dependent, then we can express one of them as a linear combination of the other two.

Now let us assume that the given vector are coplanar, then we can write

$$5\vec{a} + 6\vec{b} + 7\vec{c} = \ell(7\vec{a} - 8\vec{b} + 9\vec{c}) + m(3\vec{a} + 20\vec{b} + 5\vec{c})$$

where ℓ, m are scalars

Comparing the coefficients of \vec{a}, \vec{b} and \vec{c} on both sides of the equation

$$5 = 7\ell + 3m \quad \dots\dots\dots(i) \quad 6 = -8\ell + 20m \quad \dots\dots\dots(ii)$$

$$7 = 9\ell + 5m \quad \dots\dots\dots(iii)$$

From (i) and (iii) we get

$$4 = 8\ell \quad \Rightarrow \quad \ell = \frac{1}{2} = m \text{ which evidently satisfies (ii) equation too.}$$

Hence the given vectors are linearly dependent.

Self Practice Problems :

1. Does there exist scalars u, v, w such that $u\vec{e}_1 + v\vec{e}_2 + w\vec{e}_3 = \vec{i}$ where $\vec{e}_1 = \vec{k}, \vec{e}_2 = \vec{j} + \vec{k}, \vec{e}_3 = -\vec{j} + 2\vec{k}$?

Ans. No

2. Consider a base $\vec{a}, \vec{b}, \vec{c}$ and a vector $-2\vec{a} + 3\vec{b} - \vec{c}$. Compute the co-ordinates of this vector relatively to the base p, q, r where $\vec{p} = 2\vec{a} - 3\vec{b}, \vec{q} = \vec{a} - 2\vec{b} + \vec{c}, \vec{r} = -3\vec{a} + \vec{b} + 2\vec{c}$.

Ans. $(0, -7/5, 1/5)$

3. If \vec{a} and \vec{b} are non-collinear vectors and $\vec{A} = (x + 4y)\vec{a} + (2x + y + 1)\vec{b}$ and $\vec{B} = (y - 2x + 2)\vec{a} + (2x - 3y - 1)\vec{b}$, find x and y such that $3\vec{A} = 2\vec{B}$.

Ans. $x = 2, y = -1$

4. If vectors $\vec{a}, \vec{b}, \vec{c}$ be linearly independent, then show that : (i) $\vec{a} - 2\vec{b} + 3\vec{c}, -2\vec{a} + 3\vec{b} - 4\vec{c}, -\vec{b} + 2\vec{c}$ are linearly dependent (ii) $\vec{a} - 3\vec{b} + 2\vec{c}, -2\vec{a} - 4\vec{b} - \vec{c}, 3\vec{a} + 2\vec{b} - \vec{c}$ are linearly independent.

5. Given that $\hat{i} - \hat{j}, \hat{i} - 2\hat{j}$ are two vectors. Find a unit vector coplanar with these vectors and perpendicular to the first vector $\hat{i} - \hat{j}$. Find also the unit vector which is perpendicular to the plane of the two given

vectors. Do you thus obtain an orthonormal triad? **Ans.** $\frac{1}{\sqrt{2}}(\hat{i} + \hat{j}); \hat{k}$; Yes

6. If with reference to a right handed system of mutually perpendicular unit vectors $\hat{i}, \hat{j}, \hat{k}$ $\vec{\alpha} = 3\hat{i} - \hat{j}, \vec{\beta} = 2\hat{i} + \hat{j} - 3\hat{k}$ express $\vec{\beta}$ in the form $\vec{\beta} = \vec{\beta}_1 + \vec{\beta}_2$ where $\vec{\beta}_1$ is parallel to $\vec{\alpha}$ & $\vec{\beta}_2$ is perpendicular to $\vec{\alpha}$.

Ans. $\vec{\beta}_1 = \frac{3}{2}\vec{\alpha} - \frac{1}{2}\vec{j}, \vec{\beta}_2 = \frac{1}{2}\vec{i} + \frac{3}{2}\vec{j} - 3\vec{k}$

7. Prove that a vector \vec{r} in space can be expressed linearly in terms of three non-coplanar, non-null vectors $\vec{a}, \vec{b}, \vec{c}$ in the form $\vec{r} = \frac{[\vec{r} \vec{b} \vec{c}]\vec{a} + [\vec{r} \vec{c} \vec{a}]\vec{b} + [\vec{r} \vec{a} \vec{b}]\vec{c}}{[\vec{a} \vec{b} \vec{c}]}$

Note: Test Of Collinearity: Three points A, B, C with position vectors $\vec{a}, \vec{b}, \vec{c}$ respectively are collinear, if & only if there exist scalars x, y, z not all zero simultaneously such that; $x\vec{a} + y\vec{b} + z\vec{c} = \vec{0}$, where $x + y + z = 0$.

Note: Test Of Coplanarity: Four points A, B, C, D with position vectors $\vec{a}, \vec{b}, \vec{c}, \vec{d}$ respectively are coplanar if and only if there exist scalars x, y, z, w not all zero simultaneously such that $x\vec{a} + y\vec{b} + z\vec{c} + w\vec{d} = \vec{0}$ where, $x + y + z + w = 0$.

Solved Example Show that the vectors $2\vec{a} - \vec{b} + 3\vec{c}, \vec{a} + \vec{b} - 2\vec{c}$ and $\vec{a} + \vec{b} - 3\vec{c}$ are non-coplanar vectors.

Solution. Let, the given vectors be coplanar.
Then one of the given vectors is expressible in terms of the other two.

$$\text{Let } 2\vec{a} - \vec{b} + 3\vec{c} = x(\vec{a} + \vec{b} - 2\vec{c}) + y(\vec{a} + \vec{b} - 3\vec{c}), \text{ for some scalars } x \text{ and } y.$$

$$\Rightarrow 2\vec{a} - \vec{b} + 3\vec{c} = (x + y)\vec{a} + (x + y)\vec{b} + (-2x - 3y)\vec{c}$$

$$\Rightarrow 2 = x + y, -1 = x + y \text{ and } 3 = -2x - 3y.$$

Solving, first and third of these equations, we get $x = 9$ and $y = -7$.

Clearly, these values do not satisfy the third equation.

Hence, the given vectors are not coplanar.

Solved Example: Prove that four points $2\vec{a} + 3\vec{b} - \vec{c}, \vec{a} - 2\vec{b} + 3\vec{c}, 3\vec{a} + 4\vec{b} - 2\vec{c}$ and $\vec{a} - 6\vec{b} + 6\vec{c}$ are coplanar.

Solution. Let the given four points be P, Q, R and S respectively. These points are coplanar if the vectors \vec{PQ}, \vec{PR} and \vec{PS} are coplanar. These vectors are coplanar iff one of them can be expressed as a linear combination of other two. So, let

$$\vec{PQ} = x\vec{PR} + y\vec{PS}$$

$$\Rightarrow -\vec{a} - 5\vec{b} + 4\vec{c} = x(\vec{a} + \vec{b} - \vec{c}) + y(-\vec{a} - 9\vec{b} - 7\vec{c}) \Rightarrow -\vec{a} - 5\vec{b} + 4\vec{c} = (x - y)\vec{a} + (x - 9y)\vec{b} + (-x + 7y)\vec{c}$$

$$\Rightarrow x - y = -1, x - 9y = -5, -x + 7y = 4 \quad [\text{Equating coeff. of } \vec{a}, \vec{b}, \vec{c} \text{ on both sides}]$$


Solving the first of these three equations, we get $x = -\frac{1}{2}, y = \frac{1}{2}$.

These values also satisfy the third equation. Hence, the given four points are coplanar.


Self Practice Problems :

1. If, $\vec{a}, \vec{b}, \vec{c}, \vec{d}$ are any four vectors in 3-dimensional space with the same initial point and such that $3\vec{a} - 2\vec{b} + \vec{c} - 2\vec{d} = \vec{0}$, show that the terminal A, B, C, D of these vectors are coplanar. Find the point at which AC and BD meet. Find the ratio in which P divides AC and BD.
2. Show that the vector $\vec{a} - \vec{b} + \vec{c}$, $\vec{b} - \vec{c} - \vec{a}$ and $2\vec{a} - 3\vec{b} - 4\vec{c}$ are non-coplanar, where $\vec{a}, \vec{b}, \vec{c}$, are any non-coplanar vectors.
3. Find the value of λ for which the four points with position vectors $-\hat{j} - \hat{k}$, $4\hat{i} + 5\hat{j} + \lambda\hat{k}$, $3\hat{i} + 9\hat{j} + 4\hat{k}$ and $-4\hat{i} + 4\hat{j} + 4\hat{k}$ are coplanar. **Ans.** $\lambda = 1$

- 12. Application Of Vectors:** (a) Work done against a constant force \vec{F} over a displacement \vec{s} is defined as $\vec{W} = \vec{F} \cdot \vec{s}$ (b) The tangential velocity \vec{V} of a body moving in a circle is given by $\vec{V} = \vec{\omega} \times \vec{r}$ where \vec{r} is the pv of the point P.



- (c) The moment of \vec{F} about 'O' is defined as $\vec{M} = \vec{r} \times \vec{F}$ where \vec{r} is the pv of P wrt 'O'. The direction of \vec{M} is along the normal to the plane OPN such that \vec{r}, \vec{F} & \vec{M} form a right handed system.
- (d) Moment of the couple $= (\vec{r}_1 - \vec{r}_2) \times \vec{F}$ where \vec{r}_1 & \vec{r}_2 are pv's of the point of the application of the forces \vec{F} & $-\vec{F}$.



Solved Example: Forces of magnitudes 5 and 3 units acting in the directions $6\hat{i} + 2\hat{j} + 3\hat{k}$ and $3\hat{i} + 2\hat{j} + 6\hat{k}$ respectively act on a particle which is displaced from the point (2, 2, -1) to (4, 3, 1). Find the work done by the forces.

Solution. Let \vec{F} be the resultant force and \vec{d} be the displacement vector. Then,

$$\vec{F} = 5 \frac{(6\hat{i} + 2\hat{j} + 3\hat{k})}{\sqrt{36 + 4 + 9}} + 3 \frac{(3\hat{i} + 2\hat{j} + 6\hat{k})}{\sqrt{9 + 4 + 36}} = \frac{1}{7} (39\hat{i} + 4\hat{j} + 33\hat{k})$$

$$\text{and, } \vec{d} = (4\hat{i} + 3\hat{j} + \hat{k}) - (2\hat{i} + 2\hat{j} - \hat{k}) = 2\hat{i} + \hat{j} + 2\hat{k}$$

$$\therefore \text{Total work done} = \vec{F} \cdot \vec{d} = \frac{1}{7} (39\hat{i} + 4\hat{j} + 33\hat{k}) \cdot (2\hat{i} + \hat{j} + 2\hat{k})$$

$$= \frac{1}{7} (78 + 4 + 66) = \frac{148}{7} \text{ units.}$$

Self Practice Problems :1. A point describes a circle uniformly in the \hat{i}, \hat{j} plane taking 12 seconds to complete one revolution. If its initial position vector relative to the centre is \hat{i} , and the rotation is from \hat{i} to \hat{j} , find the position vector at the end of 7 seconds. Also find the velocity vector. **Ans.** $\frac{1}{2} (\hat{j} - \sqrt{3}\hat{i}), \pi/12 (\hat{i} - \sqrt{3}\hat{j})$

2. The force represented by $3\hat{i} + 2\hat{k}$ is acting through the point $5\hat{i} + 4\hat{j} - 3\hat{k}$. Find its moment about the point $\hat{i} + 3\hat{j} + \hat{k}$. **Ans.** $2\hat{i} - 20\hat{j} - 3\hat{k}$
3. Find the moment of the couple formed by the forces $5\hat{i} + \hat{k}$ and $-5\hat{i} - \hat{k}$ acting at the points (9, -1, 2) and (3, -2, 1) respectively. **Ans.** $\hat{i} - \hat{j} - 5\hat{k}$

Miscellaneous Solved Examples

Solved Example: Show that the points A, B, C with position vectors $2\hat{i} - \hat{j} + \hat{k}$, $\hat{i} - 3\hat{j} - 5\hat{k}$ and $3\hat{i} - 4\hat{j} - 4\hat{k}$ respectively, are the vertices of a right angled triangle. Also find the remaining angles of the triangle.

Solution. We have, $\vec{AB} = \text{Position vector of B} - \text{Position vector of A}$

$$= (\hat{i} - 3\hat{j} - 5\hat{k}) - (2\hat{i} - \hat{j} + \hat{k}) = -\hat{i} - 2\hat{j} - 6\hat{k}$$

$$\vec{BC} = \text{Position vector of C} - \text{Position vector of B}$$

$$= (3\hat{i} - 4\hat{j} - 4\hat{k}) - (\hat{i} - 3\hat{j} - 5\hat{k}) = 2\hat{i} - \hat{j} + \hat{k}$$

and, \vec{CA} = Position vector of A – Position vector of C

$$= (2\hat{i} - \hat{j} + \hat{k}) - (3\hat{i} - 4\hat{j} - 4\hat{k}) = -\hat{i} + 3\hat{j} + 5\hat{k}$$

Since $\vec{AB} + \vec{BC} + \vec{CA} = (-\hat{i} - 2\hat{j} - 6\hat{k}) + (2\hat{i} - \hat{j} + \hat{k}) + (-\hat{i} + 3\hat{j} + 5\hat{k}) = \vec{0}$
 So, A, B and C are the vertices of a triangle.

Now, $\vec{BC} \cdot \vec{CA} = (2\hat{i} - \hat{j} + \hat{k}) \cdot (-\hat{i} + 3\hat{j} + 5\hat{k}) = -2 - 3 + 5 = 0$

$$\Rightarrow \vec{BC} \perp \vec{CA} \Rightarrow \angle BCA = \frac{\pi}{2} \quad \text{Hence, ABC is a right angled triangle.}$$

Since a is the angle between the vectors \vec{AB} and \vec{AC} . Therefore

$$\cos A = \frac{|\vec{AB} \cdot \vec{AC}|}{|\vec{AB}| |\vec{AC}|} = \frac{(-\hat{i} - 2\hat{j} - 6\hat{k}) \cdot (\hat{i} - 3\hat{j} - 5\hat{k})}{\sqrt{(-1)^2 + (-2)^2 + (-6)^2} \sqrt{1^2 + (-3)^2 + (-5)^2}}$$

$$= \frac{-1 + 6 + 30}{\sqrt{1 + 4 + 36} \sqrt{1 + 9 + 25}} = \frac{35}{\sqrt{41} \sqrt{35}} = \sqrt{\frac{35}{41}}$$

$$A = \cos^{-1} \sqrt{\frac{35}{41}}$$

$$\cos B = \frac{|\vec{BA} \cdot \vec{BC}|}{|\vec{BA}| |\vec{BC}|} = \frac{(\hat{i} + 2\hat{j} + 6\hat{k}) \cdot (2\hat{i} - \hat{j} + \hat{k})}{\sqrt{1^2 + 2^2 + 6^2} \sqrt{2^2 + (-1)^2 + 1^2}} \Rightarrow \cos B = \frac{2 - 2 + 6}{\sqrt{41} \sqrt{6}} = \sqrt{\frac{6}{41}} \Rightarrow B = \cos^{-1} \sqrt{\frac{6}{41}}$$

Solved Example: If $\vec{a}, \vec{b}, \vec{c}$ are three mutually perpendicular vectors of equal magnitude, prove that $\vec{a} + \vec{b} + \vec{c}$ is equally inclined with vectors \vec{a}, \vec{b} and \vec{c} .

Solution.: Let $|\vec{a}| = |\vec{b}| = |\vec{c}| = \lambda$ (say). Since $\vec{a}, \vec{b}, \vec{c}$ are mutually perpendicular vectors, therefore $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{c} = \vec{c} \cdot \vec{a} = 0$

Now, $|\vec{a} + \vec{b} + \vec{c}|^2$

$$= \vec{a} \cdot \vec{a} + \vec{b} \cdot \vec{b} + \vec{c} \cdot \vec{c} + 2\vec{a} \cdot \vec{b} + 2\vec{b} \cdot \vec{c} + 2\vec{c} \cdot \vec{a}$$

$$= |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 \quad [\text{Using (i)}]$$

$$= 3\lambda^2 \quad [\because |\vec{a}| = |\vec{b}| = |\vec{c}| = \lambda]$$

$$\therefore |\vec{a} + \vec{b} + \vec{c}| = \sqrt{3}\lambda \quad \dots\dots\dots(ii)$$

Suppose $\vec{a} + \vec{b} + \vec{c}$ makes angles $\theta_1, \theta_2, \theta_3$ with \vec{a}, \vec{b} and \vec{c} respectively. Then,

$$\cos \theta_1 = \frac{\vec{a} \cdot (\vec{a} + \vec{b} + \vec{c})}{|\vec{a}| |\vec{a} + \vec{b} + \vec{c}|} = \frac{\vec{a} \cdot \vec{a} + \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}}{|\vec{a}| |\vec{a} + \vec{b} + \vec{c}|}$$

$$= \frac{|\vec{a}|^2}{|\vec{a}| |\vec{a} + \vec{b} + \vec{c}|} = \frac{|\vec{a}|}{|\vec{a} + \vec{b} + \vec{c}|} = \frac{\lambda}{\sqrt{3}\lambda} = \frac{1}{\sqrt{3}} \quad [\text{Using (ii)}]$$

$$\therefore \theta_1 = \cos^{-1} \left(\frac{1}{\sqrt{3}} \right)$$

Similarly, $\theta_2 = \cos^{-1} \left(\frac{1}{\sqrt{3}} \right)$ and $\theta_3 = \cos^{-1} \left(\frac{1}{\sqrt{3}} \right) \therefore \theta_1 = \theta_2 = \theta_3$.

Hence, $\vec{a} + \vec{b} + \vec{c}$ is equally inclined with \vec{a}, \vec{b} and \vec{c}

Solved Example: Prove using vectors : If two medians of a triangle are equal, then it is isosceles.

Solution.: Let ABC be a triangle and let BE and CF be two equal medians. Taking A as the origin, let the position vectors of B and C be \vec{b} and \vec{c} respectively. Then,

P.V. of E = $\frac{1}{2} \vec{c}$ and, P.V. of F = $\frac{1}{2} \vec{b}$ $\therefore \vec{BE} = \frac{1}{2} (\vec{c} - 2\vec{b})$


$$\vec{CF} = \frac{1}{2} (\vec{b} - 2\vec{c})$$

Now, $BE = CF \Rightarrow |\vec{BE}| = |\vec{CF}|$

$$\Rightarrow |\vec{BE}|^2 = |\vec{CF}|^2 \Rightarrow \left| \frac{1}{2} (\vec{c} - 2\vec{b}) \right|^2 = \left| \frac{1}{2} (\vec{b} - 2\vec{c}) \right|^2$$

$$\Rightarrow \frac{1}{4} |\vec{c} - 2\vec{b}|^2 = \frac{1}{4} |\vec{b} - 2\vec{c}|^2 \Rightarrow |\vec{c} - 2\vec{b}|^2 = |\vec{b} - 2\vec{c}|^2$$

$$\Rightarrow (\vec{c} - 2\vec{b}) \cdot (\vec{c} - 2\vec{b}) = (\vec{b} - 2\vec{c}) \cdot (\vec{b} - 2\vec{c})$$



$$\Rightarrow \vec{c} \cdot \vec{c} - 4\vec{b} \cdot \vec{c} + 4\vec{b} \cdot \vec{b} = \vec{b} \cdot \vec{b} - 4\vec{b} \cdot \vec{c} + 4\vec{c} \cdot \vec{c}$$

$$\Rightarrow |\vec{c}|^2 - 4\vec{b} \cdot \vec{c} + 4|\vec{b}|^2 = |\vec{b}|^2 - 4\vec{b} \cdot \vec{c} + 4|\vec{c}|^2$$

$$\Rightarrow 3|\vec{b}|^2 = 3|\vec{c}|^2 \Rightarrow |\vec{b}|^2 = |\vec{c}|^2$$

$$\Rightarrow AB = AC \quad \text{Hence, triangle ABC is an isosceles triangle.}$$

Solved Example: Using vectors : Prove that $\cos(A + B) = \cos A \cos B - \sin A \sin B$

Solution. Let OX and OY be the coordinate axes and let \hat{i} and \hat{j} be unit vectors along OX and OY respectively. Let $\angle XOP = A$ and $\angle XOQ = B$. Drawn $PL \perp OX$ and $QM \perp OX$.

Clearly angle between \vec{OP} and \vec{OQ} is $A + B$

In $\triangle OLP$, $OL = OP \cos A$ and $LP = OP \sin A$. Therefore $\vec{OL} = (OP \cos A) \hat{i}$ and $\vec{LP} = (OP \sin A) (-\hat{j})$

Now, $\vec{OL} + \vec{LP} = \vec{OP}$

$$\Rightarrow \vec{OP} = OP [(\cos A) \hat{i} - (\sin A) \hat{j}]$$

In $\triangle OMQ$, $OM = OQ \cos B$ and $MQ = OQ \sin B$.

Therefore,

$$\vec{OM} = (OQ \cos B) \hat{i}, \quad \vec{MQ} = (OQ \sin B) \hat{j}$$

Now, $\vec{OM} + \vec{MQ} = \vec{OQ}$

From (i) and (ii), we get

$$\vec{OP} \cdot \vec{OQ} = OP [(\cos A) \hat{i} - (\sin A) \hat{j}] \cdot OQ [(\cos B) \hat{i} + (\sin B) \hat{j}]$$

$$= OP \cdot OQ [\cos A \cos B - \sin A \sin B]$$

But, $\vec{OP} \cdot \vec{OQ} = |\vec{OP}| |\vec{OQ}| \cos(A + B) = OP \cdot OQ \cos(A + B)$

$$\therefore OP \cdot OQ \cos(A + B) = OP \cdot OQ [\cos A \cos B - \sin A \sin B]$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

Solved Example: Prove that in any triangle ABC

$$(i) \quad c^2 = a^2 + b^2 - 2ab \cos C \quad (ii) \quad c = b \cos A + a \cos B.$$

Solution. (i) In $\triangle ABC$, $\vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$

or, $\vec{BC} + \vec{CA} = -\vec{AB}$ (i)

Squaring both sides

$$(\vec{BC})^2 + (\vec{CA})^2 + 2(\vec{BC} \cdot \vec{CA}) = (\vec{AB})^2$$

$$\Rightarrow a^2 + b^2 + 2(\vec{BC} \cdot \vec{CA}) = c^2 \Rightarrow c^2 = a^2 + b^2 - 2ab \cos C$$

$$\Rightarrow c^2 = a^2 + b^2 - 2ab \cos C$$

$$(ii) \quad (\vec{BC} + \vec{CA}) \cdot \vec{AB} = -\vec{AB} \cdot \vec{AB}$$

$$\vec{BC} \cdot \vec{AB} + \vec{CA} \cdot \vec{AB} = -c^2$$

$$-ac \cos B - bc \cos A = -c^2$$

$$a \cos B + b \cos A = c.$$

Solved Ex.: If D, E, F are the mid-points of the sides of a triangle ABC, prove by vector method that area of

$$\triangle DEF = \frac{1}{4} (\text{area of } \triangle ABC)$$

Solution. Taking A as the origin, let the position vectors of B and C be \vec{b} and \vec{c} respectively. Then, the

position vectors of D, E and F are $\frac{1}{2}(\vec{b} + \vec{c})$, $\frac{1}{2}\vec{c}$ and $\frac{1}{2}\vec{b}$ respectively.

$$\text{Now, } \vec{DE} = \frac{1}{2}\vec{c} - \frac{1}{2}(\vec{b} + \vec{c}) = \frac{-\vec{b}}{2}$$

$$\text{and } \vec{DF} = \frac{1}{2}\vec{b} - \frac{1}{2}(\vec{b} + \vec{c}) = \frac{-\vec{c}}{2}$$

$$\therefore \text{Vector area of } \triangle DEF = \frac{1}{2} (\vec{DE} \times \vec{DF}) = \frac{1}{2} \left(\frac{-\vec{b}}{2} \times \frac{-\vec{c}}{2} \right)$$

$$= \frac{1}{8} (\vec{b} \times \vec{c}) = \frac{1}{4} \left\{ \frac{1}{2} (\vec{AB} \times \vec{AC}) \right\}$$




$$= \frac{1}{4} (\text{vector area of } \triangle ABC) \quad \text{Hence, area of } \triangle DEF = \frac{1}{4} \text{ area of } \triangle ABC.$$

Solved Example: P, Q are the mid-points of the non-parallel sides BC and AD of a trapezium ABCD. Show that $\triangle APD = \triangle CQB$.

Solution. Let $\vec{AB} = \vec{b}$ and $\vec{AD} = \vec{d}$

Now DC is parallel to AB \Rightarrow there exists a scalar t such that $\vec{DC} = t \vec{AB} = t \vec{b}$

$$\therefore \vec{AC} = \vec{AD} + \vec{DC} = \vec{d} + t \vec{b}$$




The position vectors of P and Q are $\frac{1}{2}(\vec{b} + \vec{d} + t\vec{b})$ and $\frac{1}{2}\vec{d}$ respectively.

$$\begin{aligned}\text{Now } 2\Delta \overrightarrow{APD} &= \overrightarrow{AP} \times \overrightarrow{AD} \\ &= \frac{1}{2}(\vec{b} + \vec{d} + t\vec{b}) \times \vec{d} = \frac{1}{2}(1+t)(\vec{b} \times \vec{d})\end{aligned}$$

$$\text{Also } 2\Delta \overrightarrow{CQB} = \overrightarrow{CQ} \times \overrightarrow{CB} = \left[\frac{1}{2}\vec{d} - (\vec{d} + t\vec{b}) \right] \times [\vec{b} - (\vec{d} + t\vec{b})]$$

$$= \left[-\frac{1}{2}\vec{d} - t\vec{b} \right] \times [-\vec{d} + (1-t)\vec{b}] = -\frac{1}{2}(1-t)\vec{d} \times \vec{b} + t\vec{b} \times \vec{d}$$

$$= \frac{1}{2}(1-t+2t)\vec{b} \times \vec{d} = \frac{1}{2}(1+t)\vec{b} \times \vec{d} = 2\Delta \overrightarrow{APD} \quad \text{Hence the result.}$$



Solved Example: Let \vec{u} and \vec{v} are unit vectors and \vec{w} is a vector such that $\vec{u} \times \vec{v} + \vec{u} = \vec{w}$ and $\vec{w} \times \vec{u} = \vec{v}$ then find the value of $[\vec{u} \vec{v} \vec{w}]$.

Solution. Given $\vec{u} \times \vec{v} + \vec{u} = \vec{w}$ and $\vec{w} \times \vec{u} = \vec{v}$

$$\begin{aligned}\Rightarrow (\vec{u} \times \vec{v} + \vec{u}) \times \vec{u} &= \vec{w} \times \vec{u} \Rightarrow (\vec{u} \times \vec{v}) \times \vec{u} + \vec{u} \times \vec{u} = \vec{v} \quad (\text{as, } \vec{w} \times \vec{u} = \vec{v}) \\ \Rightarrow (\vec{u} \cdot \vec{u}) \vec{v} - (\vec{v} \cdot \vec{u}) \vec{u} + \vec{u} \times \vec{u} &= \vec{v} \quad (\text{using } \vec{u} \cdot \vec{u} = 1 \text{ and } \vec{u} \times \vec{u} = 0, \text{ since unit vector}) \\ \Rightarrow \vec{v} - (\vec{v} \cdot \vec{u}) \vec{u} &= \vec{v} \Rightarrow (\vec{u} \cdot \vec{v}) \vec{u} = \vec{0} \\ \Rightarrow \vec{u} \cdot \vec{v} &= 0 \quad (\text{as; } \vec{u} \neq \vec{0}) \quad \dots\dots\dots(i) \\ \Rightarrow \vec{u} \cdot (\vec{v} \times \vec{w}) &= \vec{u} \cdot (\vec{v} \times (\vec{u} \times \vec{v} + \vec{u}))\end{aligned}$$

$$\begin{aligned}&= \vec{u} \cdot (\vec{v} \times (\vec{u} \times \vec{v} + \vec{u})) \quad (\text{given } \vec{w} = \vec{u} \times \vec{v} + \vec{u}) \\ &= \vec{u} \cdot (\vec{v} \times (\vec{u} \times \vec{v}) + \vec{v} \times \vec{u}) = \vec{u} \cdot ((\vec{v} \cdot \vec{v}) \vec{u} - (\vec{v} \cdot \vec{u}) \vec{v} + \vec{v} \times \vec{u}) \\ &= \vec{u} \cdot (|\vec{v}|^2 \vec{u} - 0 + \vec{v} \times \vec{u}) \quad (\text{as; } \vec{u} \cdot \vec{v} = 0 \text{ from (i)}) \\ &= |\vec{v}|^2 (\vec{u} \cdot \vec{u}) - \vec{u} \cdot (\vec{v} \times \vec{u}) = |\vec{v}|^2 |\vec{u}|^2 - 0 \quad (\text{as, } [\vec{u} \vec{v} \vec{u}] = 0) \\ &= 1 \quad (\text{as; } |\vec{u}| = |\vec{v}| = 1) \quad \therefore [\vec{u} \vec{v} \vec{w}] = 1\end{aligned}$$

Sol. Ex.: In any triangle, show that the perpendicular bisectors of the sides are concurrent.

Solution. Let ABC be the triangle and D, E and F are respectively middle points of sides BC, CA and AB. Let the perpendicular of D and E meet at O join OF. We are required to prove that OF is \perp to AB. Let the position vectors of A, B, C with O as origin of reference be \vec{a} , \vec{b} and \vec{c} respectively.

$$\therefore \overrightarrow{OD} = \frac{1}{2}(\vec{b} + \vec{c}), \overrightarrow{OE} = \frac{1}{2}(\vec{c} + \vec{a}) \text{ and } \overrightarrow{OF} = \frac{1}{2}(\vec{a} + \vec{b})$$

$$\text{Also } \overrightarrow{BC} = \vec{c} - \vec{b}, \overrightarrow{CA} = \vec{a} - \vec{c} \text{ and } \overrightarrow{AB} = \vec{b} - \vec{a}$$

$$\begin{aligned}\text{Since } OD \perp BC, \frac{1}{2}(\vec{b} + \vec{c}) \cdot (\vec{c} - \vec{b}) &= 0 \\ \Rightarrow b^2 &= c^2 \quad \dots\dots\dots(i)\end{aligned}$$

$$\begin{aligned}\text{Similarly } \frac{1}{2}(\vec{c} + \vec{a}) \cdot (\vec{a} - \vec{c}) &= 0 \\ \Rightarrow a^2 &= c^2 \quad \dots\dots\dots(ii) \\ \text{from (i) and (ii) we have } a^2 - b^2 &= 0\end{aligned}$$

$$\Rightarrow (\vec{a} + \vec{b}) \cdot (\vec{b} + \vec{a}) = 0 \Rightarrow \frac{1}{2}(\vec{b} + \vec{a}) \cdot (\vec{b} - \vec{a}) = 0$$

Solved Example: A, B, C, D are four points in space. using vector methods, prove that $AC^2 + BD^2 + AD^2 + BC^2 \geq AB^2 + CD^2$ what is the implication of the sign of equality.

Solution.: Let the position vector of A, B, C, D be $\vec{a}, \vec{b}, \vec{c}$ and \vec{d} respectively then

$$\begin{aligned}AC^2 + BD^2 + AD^2 + BC^2 &= (\vec{c} - \vec{a}) \cdot (\vec{c} - \vec{a}) + (\vec{d} - \vec{b}) \cdot (\vec{d} - \vec{b}) + (\vec{d} - \vec{a}) \cdot (\vec{d} - \vec{a}) + (\vec{c} - \vec{b}) \cdot (\vec{c} - \vec{b}) \\ &= |\vec{c}|^2 + |\vec{a}|^2 - 2\vec{a} \cdot \vec{c} + |\vec{d}|^2 + |\vec{b}|^2 - 2\vec{d} \cdot \vec{b} + |\vec{d}|^2 + |\vec{a}|^2 - 2\vec{a} \cdot \vec{d} + |\vec{c}|^2 + |\vec{b}|^2 - 2\vec{b} \cdot \vec{c} \\ &= |\vec{a}|^2 + |\vec{b}|^2 - 2\vec{a} \cdot \vec{b} + |\vec{c}|^2 + |\vec{d}|^2 - 2\vec{c} \cdot \vec{d} + |\vec{a}|^2 + |\vec{b}|^2 + |\vec{c}|^2 + |\vec{d}|^2 \\ &\quad + 2\vec{a} \cdot \vec{b} + 2\vec{c} \cdot \vec{d} - 2\vec{a} \cdot \vec{c} - 2\vec{b} \cdot \vec{d} - 2\vec{a} \cdot \vec{d} - 2\vec{b} \cdot \vec{c} \\ &= (\vec{a} - \vec{b}) \cdot (\vec{a} - \vec{b}) + (\vec{c} - \vec{d}) \cdot (\vec{c} - \vec{d}) + (\vec{a} - \vec{b} - \vec{c} + \vec{d}) \cdot (\vec{a} - \vec{b} - \vec{c} + \vec{d}) \geq AB^2 + CD^2 \\ &= AB^2 + CD^2 + (\vec{a} + \vec{b} - \vec{c} - \vec{d}) \cdot (\vec{a} + \vec{b} - \vec{c} - \vec{d}) \leq AB^2 + CD^2 \\ \therefore AC^2 + BD^2 + AD^2 + BC^2 &\geq AB^2 + CD^2\end{aligned}$$

for the sign of equality to hold, $\vec{a} + \vec{b} - \vec{c} - \vec{d} = \vec{0}$

$$\vec{a} - \vec{c} = \vec{d} - \vec{b}$$

$\Rightarrow \overrightarrow{AC}$ and \overrightarrow{BD} are collinear the four points A, B, C, D are collinear

