

Recap: Stationary Distribution

- ▶ π is said to be a stationary distribution for the Markov chain with transition probabilities P if

$$\pi(y) = \sum_{x \in S} \pi(x)P(x, y), \quad \forall y \in S$$

Recap: Stationary Distribution

- ▶ π is said to be a stationary distribution for the Markov chain with transition probabilities P if

$$\pi(y) = \sum_{x \in S} \pi(x)P(x, y), \quad \forall y \in S$$

- ▶ When π is stationary distribution,
 $\pi_0 = \pi \Rightarrow \pi_n = \pi, \forall n$

Recap: Stationary Distribution

- ▶ π is said to be a stationary distribution for the Markov chain with transition probabilities P if

$$\pi(y) = \sum_{x \in S} \pi(x)P(x, y), \quad \forall y \in S$$

- ▶ When π is stationary distribution,
 $\pi_0 = \pi \Rightarrow \pi_n = \pi, \forall n$
- ▶ If $\pi_n = \pi, \forall n$ then π is a stationary distribution

Recap: Stationary Distribution

- ▶ π is said to be a stationary distribution for the Markov chain with transition probabilities P if

$$\pi(y) = \sum_{x \in S} \pi(x)P(x, y), \quad \forall y \in S$$

- ▶ When π is stationary distribution,
 $\pi_0 = \pi \Rightarrow \pi_n = \pi, \forall n$
- ▶ If $\pi_n = \pi, \forall n$ then π is a stationary distribution
- ▶ For a finite chain: $P^T \pi = \pi$

Recap: Stationary Distribution

- ▶ π is said to be a stationary distribution for the Markov chain with transition probabilities P if

$$\pi(y) = \sum_{x \in S} \pi(x)P(x, y), \quad \forall y \in S$$

- ▶ When π is stationary distribution,
 $\pi_0 = \pi \Rightarrow \pi_n = \pi, \forall n$
- ▶ If $\pi_n = \pi, \forall n$ then π is a stationary distribution
- ▶ For a finite chain: $P^T \pi = \pi$
- ▶ A stationary distribution always exists for a finite chain

Recap

- ▶ $N_n(y)$ – number of visits to y till n

Recap

- ▶ $N_n(y)$ – number of visits to y till n
- ▶ $G_n(x, y) = E_x[N_n(y)] = \sum_{m=1}^n P^m(x, y)$
– expected number of visits to y till n

Recap

- ▶ $N_n(y)$ – number of visits to y till n
- ▶ $G_n(x, y) = E_x[N_n(y)] = \sum_{m=1}^n P^m(x, y)$
– expected number of visits to y till n
- ▶ $m_y = E_y[T_y]$ – mean return time to y

Recap

- ▶ $N_n(y)$ – number of visits to y till n
- ▶ $G_n(x, y) = E_x[N_n(y)] = \sum_{m=1}^n P^m(x, y)$
– expected number of visits to y till n
- ▶ $m_y = E_y[T_y]$ – mean return time to y

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{I_{[T_y < \infty]}}{m_y}, \quad w.p.1$$

Recap

- ▶ $N_n(y)$ – number of visits to y till n
- ▶ $G_n(x, y) = E_x[N_n(y)] = \sum_{m=1}^n P^m(x, y)$
– expected number of visits to y till n
- ▶ $m_y = E_y[T_y]$ – mean return time to y

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{I_{[T_y < \infty]}}{m_y}, \quad w.p.1$$

$$\lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n P^m(x, y) = \frac{\rho_{xy}}{m_y}$$

Recap: positive and null recurrent states

- ▶ y is positive recurrent if $m_y < \infty$

Recap: positive and null recurrent states

- ▶ y is positive recurrent if $m_y < \infty$
- ▶ y is null recurrent if $m_y = \infty$

Recap: positive and null recurrent states

- ▶ y is positive recurrent if $m_y < \infty$
- ▶ y is null recurrent if $m_y = \infty$
- ▶ If x is positive recurrent and x leads to y , then y is positive recurrent

Recap: positive and null recurrent states

- ▶ y is positive recurrent if $m_y < \infty$
- ▶ y is null recurrent if $m_y = \infty$
- ▶ If x is positive recurrent and x leads to y , then y is positive recurrent
- ▶ In a closed irreducible set of recurrent states either all states are positive recurrent or all states are null recurrent

Recap: positive and null recurrent states

- ▶ y is positive recurrent if $m_y < \infty$
- ▶ y is null recurrent if $m_y = \infty$
- ▶ If x is positive recurrent and x leads to y , then y is positive recurrent
- ▶ In a closed irreducible set of recurrent states either all states are positive recurrent or all states are null recurrent
- ▶ A finite closed set has to have at least one positive recurrent state

Recap: positive and null recurrent states

- ▶ y is positive recurrent if $m_y < \infty$
- ▶ y is null recurrent if $m_y = \infty$
- ▶ If x is positive recurrent and x leads to y , then y is positive recurrent
- ▶ In a closed irreducible set of recurrent states either all states are positive recurrent or all states are null recurrent
- ▶ A finite closed set has to have at least one positive recurrent state
- ▶ A finite chain cannot have null recurrent states

Recap: Existence of stationary distribution

- ▶ In any stationary distribution π , $\pi(y) = 0$ if y is transient or null recurrent

Recap: Existence of stationary distribution

- ▶ In any stationary distribution π , $\pi(y) = 0$ if y is transient or null recurrent
- ▶ An irreducible transient or null recurrent chain does not have a stationary distribution

Recap: Existence of stationary distribution

- ▶ In any stationary distribution π , $\pi(y) = 0$ if y is transient or null recurrent
- ▶ An irreducible transient or null recurrent chain does not have a stationary distribution
- ▶ An irreducible positive recurrent chain has a unique stationary distribution: $\pi(y) = \frac{1}{m_y}$

Recap: Existence of stationary distribution

- ▶ In any stationary distribution π , $\pi(y) = 0$ if y is transient or null recurrent
- ▶ An irreducible transient or null recurrent chain does not have a stationary distribution
- ▶ An irreducible positive recurrent chain has a unique stationary distribution: $\pi(y) = \frac{1}{m_y}$
- ▶ An irreducible chain has a stationary distribution iff it is positive recurrent

Recap: Existence of stationary distribution

- ▶ In any stationary distribution π , $\pi(y) = 0$ if y is transient or null recurrent
- ▶ An irreducible transient or null recurrent chain does not have a stationary distribution
- ▶ An irreducible positive recurrent chain has a unique stationary distribution: $\pi(y) = \frac{1}{m_y}$
- ▶ An irreducible chain has a stationary distribution iff it is positive recurrent
- ▶ For a non-irreducible chain, for each closed irreducible set of positive recurrent states, there is a unique stationary distribution concentrated on that set.

Recap: Existence of stationary distribution

- ▶ In any stationary distribution π , $\pi(y) = 0$ if y is transient or null recurrent
- ▶ An irreducible transient or null recurrent chain does not have a stationary distribution
- ▶ An irreducible positive recurrent chain has a unique stationary distribution: $\pi(y) = \frac{1}{m_y}$
- ▶ An irreducible chain has a stationary distribution iff it is positive recurrent
- ▶ For a non-irreducible chain, for each closed irreducible set of positive recurrent states, there is a unique stationary distribution concentrated on that set.
- ▶ All stationary distributions of the chain are convex combinations of these

Recap: Periodic chains

- ▶ The period of a state x is
$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$

Recap: Periodic chains

- ▶ The period of a state x is
$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$
- ▶ If x and y lead to each other, $d_x = d_y$

Recap: Periodic chains

- ▶ The period of a state x is
$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$
- ▶ If x and y lead to each other, $d_x = d_y$
- ▶ In an irreducible chain, all states have the same period

Recap: Periodic chains

- ▶ The period of a state x is
$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$
- ▶ If x and y lead to each other, $d_x = d_y$
- ▶ In an irreducible chain, all states have the same period
- ▶ An irreducible chain is called aperiodic if the period is 1

Recap: Periodic chains


- ▶ The period of a state x is
$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$
- ▶ If x and y lead to each other, $d_x = d_y$
- ▶ In an irreducible chain, all states have the same period
- ▶ An irreducible chain is called aperiodic if the period is 1
- ▶ For an irreducible aperiodic positive recurrent chain, π_n converges to π , the unique stationary distribution, irrespective of what π_0 is.

Recap: Periodic chains

- ▶ The period of a state x is
$$d_x = \gcd\{n \geq 1 : P^n(x, x) > 0\}$$
- ▶ If x and y lead to each other, $d_x = d_y$
- ▶ In an irreducible chain, all states have the same period
- ▶ An irreducible chain is called aperiodic if the period is 1
- ▶ For an irreducible aperiodic positive recurrent chain, π_n converges to π , the unique stationary distribution, irrespective of what π_0 is.
- ▶ Also, for an irreducible, aperiodic, positive recurrent chain, $P^n(x, y)$ converges to $\frac{1}{m_y}$


Recall: Birth-Death chains - stationary distributions

- The following is a finite irreducible birth-death chain



Recall: Birth-Death chains - stationary distributions

- ▶ The following is a finite irreducible birth-death chain




- ▶ The stationary distribution is given by

$$\pi(0) = \frac{1}{\sum_{j=0}^N \eta_j} \quad \text{and} \quad \pi(n) = \eta_n \pi(0), \quad n = 1, \dots, N$$

where $\eta_n = \frac{p_0 p_1 \cdots p_{n-1}}{q_1 q_2 \cdots q_n}, \quad n = 1, 2, \dots, N.$

Recall: Birth-Death chains - stationary distributions

- ▶ The following is a finite irreducible birth-death chain



- ▶ The stationary distribution is given by


$$\pi(0) = \frac{1}{\sum_{j=0}^N \eta_j} \quad \text{and} \quad \pi(n) = \eta_n \pi(0), \quad n = 1, \dots, N$$

where $\eta_n = \frac{p_0 p_1 \cdots p_{n-1}}{q_1 q_2 \cdots q_n}$, $n = 1, 2, \dots, N$.

- ▶ This is applicable to an infinite chain also
However, we need $\sum_{j=0}^{\infty} \eta_j < \infty$ for the stationary distribution to exist.


Recap: Birth-Death chains

- Consider a finite or infinite birth-death chain



Recap: Birth-Death chains

- Consider a finite or infinite birth-death chain




- Define

$$U(y) = P_y[T_a < T_b], \quad a < y < b, \quad U(a) = 1, \quad U(b) = 0$$

Recap: Birth-Death chains

- Consider a finite or infinite birth-death chain



- Define

$$U(y) = P_y[T_a < T_b], \quad a < y < b, \quad U(a) = 1, \quad U(b) = 0$$

- Then,


$$U(y) = \frac{\sum_{x=y}^{b-1} \gamma_x}{\sum_{x=a}^{b-1} \gamma_x}, \quad \gamma_x = \frac{q_x q_{x-1} \cdots q_{a+1}}{p_x p_{x-1} \cdots p_{a+1}}$$

$$P_y[T_b < T_a] = \frac{\sum_{x=a}^{y-1} \gamma_x}{\sum_{x=a}^{b-1} \gamma_x}$$


- Suppose this is a Gambler's ruin chain:

$$p_x = p, q_x = q, \forall x$$

► Suppose this is a Gambler's ruin chain:




- Suppose this is a Gambler's ruin chain:



- Then, $\gamma_x = \left(\frac{q}{p}\right)^x$


- Suppose this is a Gambler's ruin chain:



- Then, $\gamma_x = \left(\frac{q}{p}\right)^x$
- Hence, for a Gambler's ruin chain we get, e.g.,

$$P_i[T_N < T_0] = \frac{\sum_{x=0}^{i-1} \gamma_x}{\sum_{x=0}^{N-1} \gamma_x} = \frac{\left(\frac{q}{p}\right)^i - 1}{\left(\frac{q}{p}\right)^N - 1}$$

- Suppose this is a Gambler's ruin chain:




- Then, $\gamma_x = \left(\frac{q}{p}\right)^x$
- Hence, for a Gambler's ruin chain we get, e.g.,

$$P_i[T_N < T_0] = \frac{\sum_{x=0}^{i-1} \gamma_x}{\sum_{x=0}^{N-1} \gamma_x} = \frac{\left(\frac{q}{p}\right)^i - 1}{\left(\frac{q}{p}\right)^N - 1}$$


- This is the probability of gambler being successful

- ▶ Consider the following chain over $\{0, 1, \dots\}$

► Consider the following chain over $\{0, 1, \dots\}$




- Consider the following chain over $\{0, 1, \dots\}$




- This is an infinite irreducible birth-death chain

- Consider the following chain over $\{0, 1, \dots\}$




- This is an infinite irreducible birth-death chain
- We want to know whether the chain is transient or recurrent etc.

- Consider the following chain over $\{0, 1, \dots\}$



- This is an infinite irreducible birth-death chain
- We want to know whether the chain is transient or recurrent etc.
- We can use the earlier analysis for this too.


- Consider the following chain over $\{0, 1, \dots\}$



- This is an infinite irreducible birth-death chain
- We want to know whether the chain is transient or recurrent etc.
- We can use the earlier analysis for this too.

$$P_1[T_0 < T_n] = \frac{\sum_{x=1}^{n-1} \gamma_x}{\sum_{x=0}^{n-1} \gamma_x}, \quad \forall n > 1$$

- Consider the following chain over $\{0, 1, \dots\}$




- This is an infinite irreducible birth-death chain
- We want to know whether the chain is transient or recurrent etc.
- We can use the earlier analysis for this too.


$$\begin{aligned}
 P_1[T_0 < T_n] &= \frac{\sum_{x=1}^{n-1} \gamma_x}{\sum_{x=0}^{n-1} \gamma_x}, \quad \forall n > 1 \\
 &= \frac{\sum_{x=0}^{n-1} \gamma_x - \gamma_0}{\sum_{x=0}^{n-1} \gamma_x} = 1 - \frac{1}{\sum_{x=0}^{n-1} \gamma_x}
 \end{aligned}$$

- ▶ Consider this chain started in state 1.

- Consider this chain started in state 1.




- Consider this chain started in state 1.



$$[T_0 < T_n] \subset [T_0 < T_{n+1}], \quad n = 2, 3, \dots$$

- Consider this chain started in state 1.




$$[T_0 < T_n] \subset [T_0 < T_{n+1}], \quad n = 2, 3, \dots$$

since the chain cannot hit $n + 1$ without hitting n .

- Also, $1 \leq T_2 < T_3 < \dots < T_n$

- Consider this chain started in state 1.



$$[T_0 < T_n] \subset [T_0 < T_{n+1}], \quad n = 2, 3, \dots$$

since the chain cannot hit $n + 1$ without hitting n .

- Also, $1 \leq T_2 < T_3 < \dots < T_n$
- Hence $[T_0 < \infty]$ is same as $[T_0 < T_n, \text{ for some } n]$

- ▶ Consider this chain started in state 1.

$$[T_0 < T_n] \subset [T_0 < T_{n+1}], \quad n = 2, 3, \dots$$

since the chain cannot hit $n + 1$ without hitting n .

- ▶ Also, $1 \leq T_2 < T_3 < \dots < T_n$
- ▶ Hence $[T_0 < \infty]$ is same as $[T_0 < T_n, \text{ for some } n]$

- ▶ Consider this chain started in state 1.

$$[T_0 < T_n] \subset [T_0 < T_{n+1}], \quad n = 2, 3, \dots$$

since the chain cannot hit $n + 1$ without hitting n .

- ▶ Also, $1 \leq T_2 < T_3 < \dots < T_n$
- ▶ Hence $[T_0 < \infty]$ is same as $[T_0 < T_n, \text{ for some } n]$

$$P_1[T_0 < T_n, \text{ for some } n] = P_1(\cup_{n \geq 1} [T_0 < T_n])$$

- Consider this chain started in state 1.

$$[T_0 < T_n] \subset [T_0 < T_{n+1}], \quad n = 2, 3, \dots$$

since the chain cannot hit $n + 1$ without hitting n .

- Also, $1 \leq T_2 < T_3 < \dots < T_n$
- Hence $[T_0 < \infty]$ is same as $[T_0 < T_n, \text{ for some } n]$

$$\begin{aligned} P_1[T_0 < T_n, \text{ for some } n] &= P_1(\cup_{n \geq 1} [T_0 < T_n]) \\ &= P_1\left(\lim_{n \rightarrow \infty} [T_0 < T_n]\right) \end{aligned}$$

- Consider this chain started in state 1.

$$[T_0 < T_n] \subset [T_0 < T_{n+1}], \quad n = 2, 3, \dots$$

since the chain cannot hit $n + 1$ without hitting n .

- Also, $1 \leq T_2 < T_3 < \dots < T_n$
- Hence $[T_0 < \infty]$ is same as $[T_0 < T_n, \text{ for some } n]$

$$\begin{aligned} P_1[T_0 < T_n, \text{ for some } n] &= P_1(\cup_{n \geq 1} [T_0 < T_n]) \\ &= P_1\left(\lim_{n \rightarrow \infty} [T_0 < T_n]\right) \\ &= \lim_{n \rightarrow \infty} P_1([T_0 < T_n]) \end{aligned}$$

- Consider this chain started in state 1.

$$[T_0 < T_n] \subset [T_0 < T_{n+1}], \quad n = 2, 3, \dots$$

since the chain cannot hit $n + 1$ without hitting n .

- Also, $1 \leq T_2 < T_3 < \dots < T_n$
- Hence $[T_0 < \infty]$ is same as $[T_0 < T_n, \text{ for some } n]$

$$\begin{aligned} P_1[T_0 < T_n, \text{ for some } n] &= P_1(\cup_{n \geq 1} [T_0 < T_n]) \\ &= P_1\left(\lim_{n \rightarrow \infty} [T_0 < T_n]\right) \\ &= \lim_{n \rightarrow \infty} P_1([T_0 < T_n]) \\ &= \lim_{n \rightarrow \infty} 1 - \frac{1}{\sum_{x=0}^{n-1} \gamma_x} \end{aligned}$$

- Consider this chain started in state 1.

$$[T_0 < T_n] \subset [T_0 < T_{n+1}], \quad n = 2, 3, \dots$$

since the chain cannot hit $n + 1$ without hitting n .

- Also, $1 \leq T_2 < T_3 < \dots < T_n$
- Hence $[T_0 < \infty]$ is same as $[T_0 < T_n, \text{ for some } n]$

$$\begin{aligned} P_1[T_0 < T_n, \text{ for some } n] &= P_1(\cup_{n>1} [T_0 < T_n]) \\ &= P_1\left(\lim_{n \rightarrow \infty} [T_0 < T_n]\right) \\ &= \lim_{n \rightarrow \infty} P_1([T_0 < T_n]) \\ &= \lim_{n \rightarrow \infty} 1 - \frac{1}{\sum_{x=0}^{n-1} \gamma_x} \\ \Rightarrow P_1[T_0 < \infty] &= 1 - \frac{1}{\sum_{x=0}^{\infty} \gamma_x} \end{aligned}$$

► **Theorem:** The chain is recurrent iff $\sum_{x=0}^{\infty} \gamma_x = \infty$

► **Theorem:** The chain is recurrent iff $\sum_{x=0}^{\infty} \gamma_x = \infty$

Proof

- ▶ **Theorem:** The chain is recurrent iff $\sum_{x=0}^{\infty} \gamma_x = \infty$

Proof

- ▶ Suppose chain is recurrent.

- **Theorem:** The chain is recurrent iff $\sum_{x=0}^{\infty} \gamma_x = \infty$

Proof

- Suppose chain is recurrent. Since it is irreducible,

$$P_1[T_0 < \infty] = 1$$

- **Theorem:** The chain is recurrent iff $\sum_{x=0}^{\infty} \gamma_x = \infty$

Proof

- Suppose chain is recurrent. Since it is irreducible,

$$P_1[T_0 < \infty] = 1 \Rightarrow \sum_{x=0}^{\infty} \gamma_x = \infty$$

- **Theorem:** The chain is recurrent iff $\sum_{x=0}^{\infty} \gamma_x = \infty$

Proof

- Suppose chain is recurrent. Since it is irreducible,

$$P_1[T_0 < \infty] = 1 \Rightarrow \sum_{x=0}^{\infty} \gamma_x = \infty$$

- Suppose $\sum_{x=0}^{\infty} \gamma_x = \infty$

- **Theorem:** The chain is recurrent iff $\sum_{x=0}^{\infty} \gamma_x = \infty$

Proof

- Suppose chain is recurrent. Since it is irreducible,

$$P_1[T_0 < \infty] = 1 \Rightarrow \sum_{x=0}^{\infty} \gamma_x = \infty$$

- Suppose $\sum_{x=0}^{\infty} \gamma_x = \infty \Rightarrow P_1[T_0 < \infty] = 1$

- **Theorem:** The chain is recurrent iff $\sum_{x=0}^{\infty} \gamma_x = \infty$

Proof

- Suppose chain is recurrent. Since it is irreducible,

$$P_1[T_0 < \infty] = 1 \Rightarrow \sum_{x=0}^{\infty} \gamma_x = \infty$$

- Suppose $\sum_{x=0}^{\infty} \gamma_x = \infty \Rightarrow P_1[T_0 < \infty] = 1$

$$P_0[T_0 < \infty] = P(0,0) + P(0,1) P_1[T_0 < \infty]$$

- **Theorem:** The chain is recurrent iff $\sum_{x=0}^{\infty} \gamma_x = \infty$

Proof

- Suppose chain is recurrent. Since it is irreducible,

$$P_1[T_0 < \infty] = 1 \Rightarrow \sum_{x=0}^{\infty} \gamma_x = \infty$$

- Suppose $\sum_{x=0}^{\infty} \gamma_x = \infty \Rightarrow P_1[T_0 < \infty] = 1$

$$\begin{aligned} P_0[T_0 < \infty] &= P(0,0) + P(0,1) P_1[T_0 < \infty] \\ &= P(0,0) + P(0,1) = 1 \end{aligned}$$

- **Theorem:** The chain is recurrent iff $\sum_{x=0}^{\infty} \gamma_x = \infty$

Proof

- Suppose chain is recurrent. Since it is irreducible,

$$P_1[T_0 < \infty] = 1 \Rightarrow \sum_{x=0}^{\infty} \gamma_x = \infty$$

- Suppose $\sum_{x=0}^{\infty} \gamma_x = \infty \Rightarrow P_1[T_0 < \infty] = 1$

$$\begin{aligned} P_0[T_0 < \infty] &= P(0,0) + P(0,1) P_1[T_0 < \infty] \\ &= P(0,0) + P(0,1) = 1 \end{aligned}$$

- Implies state 0 is recurrent and hence the chain is recurrent because it is irreducible.

- **Theorem:** The chain is recurrent iff $\sum_{x=0}^{\infty} \gamma_x = \infty$

Proof

- Suppose chain is recurrent. Since it is irreducible,

$$P_1[T_0 < \infty] = 1 \Rightarrow \sum_{x=0}^{\infty} \gamma_x = \infty$$

- Suppose $\sum_{x=0}^{\infty} \gamma_x = \infty \Rightarrow P_1[T_0 < \infty] = 1$

$$\begin{aligned} P_0[T_0 < \infty] &= P(0,0) + P(0,1) P_1[T_0 < \infty] \\ &= P(0,0) + P(0,1) = 1 \end{aligned}$$

- Implies state 0 is recurrent and hence the chain is recurrent because it is irreducible.
- Note that we have used the fact that the chain is infinite only to the right.

- ▶ The chain is transient if $\sum_{x=0}^{\infty} \gamma_x < \infty$

- ▶ The chain is transient if $\sum_{x=0}^{\infty} \gamma_x < \infty$
- ▶ Let $p_x = p, q_x = q \Rightarrow \gamma_x = \left(\frac{q}{p}\right)^x$

- ▶ The chain is transient if $\sum_{x=0}^{\infty} \gamma_x < \infty$
- ▶ Let $p_x = p, q_x = q \Rightarrow \gamma_x = \left(\frac{q}{p}\right)^x$

$$\text{Transient if } \sum_{x=0}^{\infty} \left(\frac{q}{p}\right)^x < \infty \Leftrightarrow q < p$$

- ▶ The chain is transient if $\sum_{x=0}^{\infty} \gamma_x < \infty$
- ▶ Let $p_x = p, q_x = q \Rightarrow \gamma_x = \left(\frac{q}{p}\right)^x$

$$\text{Transient if } \sum_{x=0}^{\infty} \left(\frac{q}{p}\right)^x < \infty \Leftrightarrow q < p$$

$$\text{Recurrent if } \sum_{x=0}^{\infty} \left(\frac{q}{p}\right)^x = \infty \Leftrightarrow q \geq p$$

- ▶ The chain is transient if $\sum_{x=0}^{\infty} \gamma_x < \infty$
- ▶ Let $p_x = p, q_x = q \Rightarrow \gamma_x = \left(\frac{q}{p}\right)^x$

$$\text{Transient if } \sum_{x=0}^{\infty} \left(\frac{q}{p}\right)^x < \infty \Leftrightarrow q < p$$

$$\text{Recurrent if } \sum_{x=0}^{\infty} \left(\frac{q}{p}\right)^x = \infty \Leftrightarrow q \geq p$$

- ▶ Intuitively clear

- ▶ The chain is transient if $\sum_{x=0}^{\infty} \gamma_x < \infty$
- ▶ Let $p_x = p, q_x = q \Rightarrow \gamma_x = \left(\frac{q}{p}\right)^x$

$$\text{Transient if } \sum_{x=0}^{\infty} \left(\frac{q}{p}\right)^x < \infty \Leftrightarrow q < p$$

$$\text{Recurrent if } \sum_{x=0}^{\infty} \left(\frac{q}{p}\right)^x = \infty \Leftrightarrow q \geq p$$

- ▶ Intuitively clear
- ▶ This chain with $q < p$ is an example of an irreducible chain that is wholly transient

► We know the chain is recurrent if $\sum_{x=0}^{\infty} \left(\frac{q}{p}\right)^x = \infty$

- ▶ We know the chain is recurrent if $\sum_{x=0}^{\infty} \left(\frac{q}{p}\right)^x = \infty$
- ▶ When will this chain be positive recurrent?

- ▶ We know the chain is recurrent if $\sum_{x=0}^{\infty} \left(\frac{q}{p}\right)^x = \infty$
- ▶ When will this chain be positive recurrent?
- ▶ We know that an irreducible chain is positive recurrent if and only if it has a stationary distribution.

- ▶ We know the chain is recurrent if $\sum_{x=0}^{\infty} \left(\frac{q}{p}\right)^x = \infty$
- ▶ When will this chain be positive recurrent?
- ▶ We know that an irreducible chain is positive recurrent if and only if it has a stationary distribution.
- ▶ We can check if it has a stationary distribution

- ▶ We know the chain is recurrent if $\sum_{x=0}^{\infty} \left(\frac{q}{p}\right)^x = \infty$
- ▶ When will this chain be positive recurrent?
- ▶ We know that an irreducible chain is positive recurrent if and only if it has a stationary distribution.
- ▶ We can check if it has a stationary distribution
- ▶ The equations that we derived earlier hold for this infinite case also.

- ▶ We derived earlier the equations that a stationary distribution of this chain (if it exists) has to satisfy

- ▶ We derived earlier the equations that a stationary distribution of this chain (if it exists) has to satisfy

$$\pi(n) = \eta_n \pi(0), \quad \text{where} \quad \eta_n = \frac{p_0 p_1 \cdots p_{n-1}}{q_1 q_2 \cdots q_n}, \quad n = 1, 2, \cdots,$$

- ▶ We derived earlier the equations that a stationary distribution of this chain (if it exists) has to satisfy

$$\pi(n) = \eta_n \pi(0), \quad \text{where } \eta_n = \frac{p_0 p_1 \cdots p_{n-1}}{q_1 q_2 \cdots q_n}, \quad n = 1, 2, \cdots,$$

- ▶ Setting $\eta_0 = 1$, we get $\pi(0) \sum_{j=0}^{\infty} \eta_j = 1$

- ▶ We derived earlier the equations that a stationary distribution of this chain (if it exists) has to satisfy

$$\pi(n) = \eta_n \pi(0), \quad \text{where } \eta_n = \frac{p_0 p_1 \cdots p_{n-1}}{q_1 q_2 \cdots q_n}, \quad n = 1, 2, \dots,$$

- ▶ Setting $\eta_0 = 1$, we get $\pi(0) \sum_{j=0}^{\infty} \eta_j = 1$
- ▶ Hence stationary distribution exists iff $\sum_{j=0}^{\infty} \eta_j < \infty$

- ▶ We derived earlier the equations that a stationary distribution of this chain (if it exists) has to satisfy

$$\pi(n) = \eta_n \pi(0), \quad \text{where } \eta_n = \frac{p_0 p_1 \cdots p_{n-1}}{q_1 q_2 \cdots q_n}, \quad n = 1, 2, \dots,$$

- ▶ Setting $\eta_0 = 1$, we get $\pi(0) \sum_{j=0}^{\infty} \eta_j = 1$
- ▶ Hence stationary distribution exists iff $\sum_{j=0}^{\infty} \eta_j < \infty$
- ▶ Let $p_x = p, q_x = q$

- ▶ We derived earlier the equations that a stationary distribution of this chain (if it exists) has to satisfy

$$\pi(n) = \eta_n \pi(0), \quad \text{where } \eta_n = \frac{p_0 p_1 \cdots p_{n-1}}{q_1 q_2 \cdots q_n}, \quad n = 1, 2, \dots,$$

- ▶ Setting $\eta_0 = 1$, we get $\pi(0) \sum_{j=0}^{\infty} \eta_j = 1$
- ▶ Hence stationary distribution exists iff $\sum_{j=0}^{\infty} \eta_j < \infty$
- ▶ Let $p_x = p, q_x = q$

$$\sum_{j=0}^{\infty} \eta_j = \sum_{j=0}^{\infty} \left(\frac{p}{q}\right)^j < \infty \iff p < q$$

- ▶ We derived earlier the equations that a stationary distribution of this chain (if it exists) has to satisfy

$$\pi(n) = \eta_n \pi(0), \quad \text{where } \eta_n = \frac{p_0 p_1 \cdots p_{n-1}}{q_1 q_2 \cdots q_n}, \quad n = 1, 2, \dots,$$

- ▶ Setting $\eta_0 = 1$, we get $\pi(0) \sum_{j=0}^{\infty} \eta_j = 1$
- ▶ Hence stationary distribution exists iff $\sum_{j=0}^{\infty} \eta_j < \infty$
- ▶ Let $p_x = p, q_x = q$

$$\sum_{j=0}^{\infty} \eta_j = \sum_{j=0}^{\infty} \left(\frac{p}{q}\right)^j < \infty \iff p < q$$

- ▶ Thus in this special case, the chain is

- ▶ We derived earlier the equations that a stationary distribution of this chain (if it exists) has to satisfy

$$\pi(n) = \eta_n \pi(0), \quad \text{where } \eta_n = \frac{p_0 p_1 \cdots p_{n-1}}{q_1 q_2 \cdots q_n}, \quad n = 1, 2, \dots,$$

- ▶ Setting $\eta_0 = 1$, we get $\pi(0) \sum_{j=0}^{\infty} \eta_j = 1$
- ▶ Hence stationary distribution exists iff $\sum_{j=0}^{\infty} \eta_j < \infty$
- ▶ Let $p_x = p, q_x = q$

$$\sum_{j=0}^{\infty} \eta_j = \sum_{j=0}^{\infty} \left(\frac{p}{q}\right)^j < \infty \iff p < q$$

- ▶ Thus in this special case, the chain is
 - ▶ transient if $p > q$; recurrent if $p \leq q$

- ▶ We derived earlier the equations that a stationary distribution of this chain (if it exists) has to satisfy

$$\pi(n) = \eta_n \pi(0), \quad \text{where } \eta_n = \frac{p_0 p_1 \cdots p_{n-1}}{q_1 q_2 \cdots q_n}, \quad n = 1, 2, \dots,$$

- ▶ Setting $\eta_0 = 1$, we get $\pi(0) \sum_{j=0}^{\infty} \eta_j = 1$
- ▶ Hence stationary distribution exists iff $\sum_{j=0}^{\infty} \eta_j < \infty$
- ▶ Let $p_x = p, q_x = q$

$$\sum_{j=0}^{\infty} \eta_j = \sum_{j=0}^{\infty} \left(\frac{p}{q}\right)^j < \infty \Leftrightarrow p < q$$

- ▶ Thus in this special case, the chain is
 - ▶ transient if $p > q$; recurrent if $p \leq q$
 - ▶ positive recurrent if $p < q$

- ▶ We derived earlier the equations that a stationary distribution of this chain (if it exists) has to satisfy

$$\pi(n) = \eta_n \pi(0), \quad \text{where } \eta_n = \frac{p_0 p_1 \cdots p_{n-1}}{q_1 q_2 \cdots q_n}, \quad n = 1, 2, \dots,$$


- ▶ Setting $\eta_0 = 1$, we get $\pi(0) \sum_{j=0}^{\infty} \eta_j = 1$
- ▶ Hence stationary distribution exists iff $\sum_{j=0}^{\infty} \eta_j < \infty$
- ▶ Let $p_x = p, q_x = q$

$$\sum_{j=0}^{\infty} \eta_j = \sum_{j=0}^{\infty} \left(\frac{p}{q}\right)^j < \infty \iff p < q$$


- ▶ Thus in this special case, the chain is
 - ▶ transient if $p > q$; recurrent if $p \leq q$
 - ▶ positive recurrent if $p < q$
 - ▶ null recurrent if $p = q$

- ▶ This analysis can handle chains which are infinite in one direction

- ▶ This analysis can handle chains which are infinite in one direction
- ▶ Consider the following random walk chain




- ▶ This analysis can handle chains which are infinite in one direction
- ▶ Consider the following random walk chain




- ▶ The state space here is $\{\dots, -1, 0, +1, \dots\}$

- ▶ This analysis can handle chains which are infinite in one direction
- ▶ Consider the following random walk chain




- ▶ The state space here is $\{\dots, -1, 0, +1, \dots\}$
- ▶ The chain is irreducible and periodic with period 2

- ▶ This analysis can handle chains which are infinite in one direction
- ▶ Consider the following random walk chain




- ▶ The state space here is $\{\dots, -1, 0, +1, \dots\}$
- ▶ The chain is irreducible and periodic with period 2
- ▶ $P^{2n+1}(0, 0) = 0$ and $P^{2n}(0, 0) = {}^{2n}C_n p^n (1-p)^n$.

- ▶ This analysis can handle chains which are infinite in one direction
- ▶ Consider the following random walk chain




- ▶ The state space here is $\{\dots, -1, 0, +1, \dots\}$
- ▶ The chain is irreducible and periodic with period 2
- ▶ $P^{2n+1}(0, 0) = 0$ and $P^{2n}(0, 0) = {}^{2n}C_n p^n (1-p)^n$.
- ▶ State 0 is recurrent if $\sum_n P^{2n}(0, 0) = \infty$


- ▶ This analysis can handle chains which are infinite in one direction
- ▶ Consider the following random walk chain



- ▶ The state space here is $\{\dots, -1, 0, +1, \dots\}$
- ▶ The chain is irreducible and periodic with period 2
- ▶ $P^{2n+1}(0, 0) = 0$ and $P^{2n}(0, 0) = {}^{2n}C_n p^n (1-p)^n$.
- ▶ State 0 is recurrent if $\sum_n P^{2n}(0, 0) = \infty$
- ▶ We can show that the chain is transient if $p \neq 0.5$ and is recurrent if $p = 0.5$.




► We use Stirling approximation: $n! \sim n^{n+0.5} e^{-n} \sqrt{2\pi}$.




► We use Stirling approximation: $n! \sim n^{n+0.5}e^{-n}\sqrt{2\pi}$.

$$P^{2n}(0,0) \sim {}^{2n}C_n p^n (1-p)^n = \frac{(2n)!}{n!n!} p^n (1-p)^n$$




► We use Stirling approximation: $n! \sim n^{n+0.5}e^{-n}\sqrt{2\pi}$.

$$\begin{aligned}
 P^{2n}(0,0) &\sim {}^{2n}C_n p^n (1-p)^n = \frac{(2n)!}{n!n!} p^n (1-p)^n \\
 &= \frac{(2n)^{2n+0.5} e^{-2n} \sqrt{2\pi}}{n^{n+0.5} n^{n+0.5} e^{-n} e^{-n} (2\pi)} (p(1-p))^n
 \end{aligned}$$



► We use Stirling approximation: $n! \sim n^{n+0.5}e^{-n}\sqrt{2\pi}$.

$$\begin{aligned}
 P^{2n}(0,0) &\sim {}^{2n}C_n p^n (1-p)^n = \frac{(2n)!}{n!n!} p^n (1-p)^n \\
 &= \frac{(2n)^{2n+0.5} e^{-2n} \sqrt{2\pi}}{n^{n+0.5} n^{n+0.5} e^{-n} e^{-n} (2\pi)} (p(1-p))^n \\
 &= \frac{2^{2n} \sqrt{2n}}{n \sqrt{2\pi}} (p(1-p))^n
 \end{aligned}$$



► We use Stirling approximation: $n! \sim n^{n+0.5}e^{-n}\sqrt{2\pi}$.

$$\begin{aligned}
 P^{2n}(0,0) &\sim {}^{2n}C_n p^n (1-p)^n = \frac{(2n)!}{n!n!} p^n (1-p)^n \\
 &= \frac{(2n)^{2n+0.5} e^{-2n} \sqrt{2\pi}}{n^{n+0.5} n^{n+0.5} e^{-n} e^{-n} (2\pi)} (p(1-p))^n \\
 &= \frac{2^{2n} \sqrt{2n}}{n \sqrt{2\pi}} (p(1-p))^n \\
 &= \frac{(4p(1-p))^n}{\sqrt{\pi n}}
 \end{aligned}$$

► We got

$$P^{2n}(0,0) \sim \frac{(4p(1-P))^n}{\sqrt{\pi n}}$$

- ▶ We got

$$P^{2n}(0,0) \sim \frac{(4p(1-P))^n}{\sqrt{\pi n}}$$

- ▶ When $a_n \sim b_n$, we have $\sum_n a_n = \infty$ iff $\sum_n b_n = \infty$

- ▶ We got

$$P^{2n}(0,0) \sim \frac{(4p(1-P))^n}{\sqrt{\pi n}}$$

- ▶ When $a_n \sim b_n$, we have $\sum_n a_n = \infty$ iff $\sum_n b_n = \infty$
- ▶ So, for state 0 to be recurrent, we need $\sum_n \frac{(4p(1-P))^n}{\sqrt{\pi n}} = \infty$.

- ▶ We got

$$P^{2n}(0,0) \sim \frac{(4p(1-P))^n}{\sqrt{\pi n}}$$

- ▶ When $a_n \sim b_n$, we have $\sum_n a_n = \infty$ iff $\sum_n b_n = \infty$
- ▶ So, for state 0 to be recurrent, we need $\sum_n \frac{(4p(1-P))^n}{\sqrt{\pi n}} = \infty$.
- ▶ If $p \neq 0.5$, then $4p(1-p) < 1$

- ▶ We got

$$P^{2n}(0,0) \sim \frac{(4p(1-P))^n}{\sqrt{\pi n}}$$

- ▶ When $a_n \sim b_n$, we have $\sum_n a_n = \infty$ iff $\sum_n b_n = \infty$
- ▶ So, for state 0 to be recurrent, we need $\sum_n \frac{(4p(1-P))^n}{\sqrt{\pi n}} = \infty$.
- ▶ If $p \neq 0.5$, then $4p(1-p) < 1$ and $\sum_n \frac{\alpha^n}{\sqrt{n}} < \infty$ if $\alpha < 1$.

- ▶ We got

$$P^{2n}(0,0) \sim \frac{(4p(1-P))^n}{\sqrt{\pi n}}$$

- ▶ When $a_n \sim b_n$, we have $\sum_n a_n = \infty$ iff $\sum_n b_n = \infty$
- ▶ So, for state 0 to be recurrent, we need $\sum_n \frac{(4p(1-P))^n}{\sqrt{\pi n}} = \infty$.
- ▶ If $p \neq 0.5$, then $4p(1-p) < 1$ and $\sum_n \frac{\alpha^n}{\sqrt{n}} < \infty$ if $\alpha < 1$.
- ▶ Hence, if $p \neq 0.5$, then, $\sum_n P^{2n}(0,0) < \infty$

- ▶ We got

$$P^{2n}(0,0) \sim \frac{(4p(1-P))^n}{\sqrt{\pi n}}$$


- ▶ When $a_n \sim b_n$, we have $\sum_n a_n = \infty$ iff $\sum_n b_n = \infty$
- ▶ So, for state 0 to be recurrent, we need $\sum_n \frac{(4p(1-P))^n}{\sqrt{\pi n}} = \infty$.
- ▶ If $p \neq 0.5$, then $4p(1-p) < 1$ and $\sum_n \frac{\alpha^n}{\sqrt{n}} < \infty$ if $\alpha < 1$.
- ▶ Hence, if $p \neq 0.5$, then, $\sum_n P^{2n}(0,0) < \infty$
- ▶ If $p = 0.5$ then $4p(1-p) = 1$ and hence $\sum_n P^{2n}(0,0) = \infty$.

- ▶ We got


$$P^{2n}(0,0) \sim \frac{(4p(1-P))^n}{\sqrt{\pi n}}$$

- ▶ When $a_n \sim b_n$, we have $\sum_n a_n = \infty$ iff $\sum_n b_n = \infty$
- ▶ So, for state 0 to be recurrent, we need $\sum_n \frac{(4p(1-P))^n}{\sqrt{\pi n}} = \infty$.
- ▶ If $p \neq 0.5$, then $4p(1-p) < 1$ and $\sum_n \frac{\alpha^n}{\sqrt{n}} < \infty$ if $\alpha < 1$.
- ▶ Hence, if $p \neq 0.5$, then, $\sum_n P^{2n}(0,0) < \infty$
- ▶ If $p = 0.5$ then $4p(1-p) = 1$ and hence $\sum_n P^{2n}(0,0) = \infty$.
- ▶ The chain is recurrent only when $p = 0.5$

- The one dimensional random walk chain is



- ▶ The one dimensional random walk chain is



- ▶ This chain is transient if $p \neq 0.5$ and is recurrent if $p = 0.5$.

- ▶ Let $\{X_n, n \geq 0\}$ be an irreducible markov chain on a finite state space S with stationary distribution π .

- ▶ Let $\{X_n, n \geq 0\}$ be an irreducible markov chain on a finite state space S with stationary distribution π .
- ▶ Let $r : S \rightarrow \mathbb{R}$ be a bounded function.

- ▶ Let $\{X_n, n \geq 0\}$ be an irreducible markov chain on a finite state space S with stationary distribution π .
- ▶ Let $r : S \rightarrow \mathbb{R}$ be a bounded function.
- ▶ Suppose we want $E[r(X)]$ with respect to the stationary distribution π

- ▶ Let $\{X_n, n \geq 0\}$ be an irreducible markov chain on a finite state space S with stationary distribution π .
- ▶ Let $r : S \rightarrow \Re$ be a bounded function.
- ▶ Suppose we want $E[r(X)]$ with respect to the stationary distribution π ($E[r(X)] = \sum_{j \in S} r(j)\pi(j)$)

- ▶ Let $\{X_n, n \geq 0\}$ be an irreducible markov chain on a finite state space S with stationary distribution π .
- ▶ Let $r : S \rightarrow \Re$ be a bounded function.
- ▶ Suppose we want $E[r(X)]$ with respect to the stationary distribution π ($E[r(X)] = \sum_{j \in S} r(j)\pi(j)$)
- ▶ Let $N_n(j)$ be as earlier. Then

- ▶ Let $\{X_n, n \geq 0\}$ be an irreducible markov chain on a finite state space S with stationary distribution π .
- ▶ Let $r : S \rightarrow \mathbb{R}$ be a bounded function.
- ▶ Suppose we want $E[r(X)]$ with respect to the stationary distribution π ($E[r(X)] = \sum_{j \in S} r(j)\pi(j)$)
- ▶ Let $N_n(j)$ be as earlier. Then

$$\frac{1}{n} \sum_{m=1}^n r(X_m) = \frac{1}{n} \sum_{j \in S} N_n(j) r(j)$$

- ▶ Let $\{X_n, n \geq 0\}$ be an irreducible markov chain on a finite state space S with stationary distribution π .
- ▶ Let $r : S \rightarrow \mathbb{R}$ be a bounded function.
- ▶ Suppose we want $E[r(X)]$ with respect to the stationary distribution π ($E[r(X)] = \sum_{j \in S} r(j)\pi(j)$)
- ▶ Let $N_n(j)$ be as earlier. Then

$$\frac{1}{n} \sum_{m=1}^n r(X_m) = \frac{1}{n} \sum_{j \in S} N_n(j) r(j)$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n r(X_m) = \sum_{j \in S} r(j) \lim_{n \rightarrow \infty} \frac{N_n(j)}{n}$$

- ▶ Let $\{X_n, n \geq 0\}$ be an irreducible markov chain on a finite state space S with stationary distribution π .
- ▶ Let $r : S \rightarrow \mathbb{R}$ be a bounded function.
- ▶ Suppose we want $E[r(X)]$ with respect to the stationary distribution π ($E[r(X)] = \sum_{j \in S} r(j)\pi(j)$)
- ▶ Let $N_n(j)$ be as earlier. Then

$$\frac{1}{n} \sum_{m=1}^n r(X_m) = \frac{1}{n} \sum_{j \in S} N_n(j) r(j)$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n r(X_m) = \sum_{j \in S} r(j) \lim_{n \rightarrow \infty} \frac{N_n(j)}{n} = \sum_{j \in S} r(j) \pi(j)$$

- ▶ Let $\{X_n, n \geq 0\}$ be an irreducible markov chain on a finite state space S with stationary distribution π .
- ▶ Let $r : S \rightarrow \mathbb{R}$ be a bounded function.
- ▶ Suppose we want $E[r(X)]$ with respect to the stationary distribution π ($E[r(X)] = \sum_{j \in S} r(j)\pi(j)$)
- ▶ Let $N_n(j)$ be as earlier. Then

$$\frac{1}{n} \sum_{m=1}^n r(X_m) = \frac{1}{n} \sum_{j \in S} N_n(j) r(j)$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n r(X_m) = \sum_{j \in S} r(j) \lim_{n \rightarrow \infty} \frac{N_n(j)}{n} = \sum_{j \in S} r(j)\pi(j)$$

- ▶ This is known as the ergodic theorem for Markov Chains

MCMC Sampling

MCMC Sampling

- ▶ Consider a distribution over (finite) S : $\pi(x) = \frac{b(x)}{Z}$

MCMC Sampling

- ▶ Consider a distribution over (finite) S : $\pi(x) = \frac{b(x)}{Z}$
- ▶ Since this is a distribution, $Z = \sum_{x \in S} b(x)$

MCMC Sampling

- ▶ Consider a distribution over (finite) S : $\pi(x) = \frac{b(x)}{Z}$
- ▶ Since this is a distribution, $Z = \sum_{x \in S} b(x)$
- ▶ We assume, we can efficiently calculate $b(x)$ for any x but computation of Z is intractable or computationally expensive

MCMC Sampling

- ▶ Consider a distribution over (finite) S : $\pi(x) = \frac{b(x)}{Z}$
 - ▶ Since this is a distribution, $Z = \sum_{x \in S} b(x)$
 - ▶ We assume, we can efficiently calculate $b(x)$ for any x but computation of Z is intractable or computationally expensive
- E.g., the Boltzmann distribution: $b(x) = e^{-E(x)/KT}$

MCMC Sampling

- ▶ Consider a distribution over (finite) S : $\pi(x) = \frac{b(x)}{Z}$
- ▶ Since this is a distribution, $Z = \sum_{x \in S} b(x)$
- ▶ We assume, we can efficiently calculate $b(x)$ for any x but computation of Z is intractable or computationally expensive
E.g., the Boltzmann distribution: $b(x) = e^{-E(x)/KT}$
- ▶ We want $E[g(X)]$ w.r.t. distribution π (for any g)

MCMC Sampling

- ▶ Consider a distribution over (finite) S : $\pi(x) = \frac{b(x)}{Z}$
- ▶ Since this is a distribution, $Z = \sum_{x \in S} b(x)$
- ▶ We assume, we can efficiently calculate $b(x)$ for any x but computation of Z is intractable or computationally expensive
E.g., the Boltzmann distribution: $b(x) = e^{-E(x)/KT}$
- ▶ We want $E[g(X)]$ w.r.t. distribution π (for any g)

$$E[g(X)] = \sum_x g(x) \pi(x)$$

MCMC Sampling

- ▶ Consider a distribution over (finite) S : $\pi(x) = \frac{b(x)}{Z}$
- ▶ Since this is a distribution, $Z = \sum_{x \in S} b(x)$
- ▶ We assume, we can efficiently calculate $b(x)$ for any x but computation of Z is intractable or computationally expensive
E.g., the Boltzmann distribution: $b(x) = e^{-E(x)/KT}$
- ▶ We want $E[g(X)]$ w.r.t. distribution π (for any g)

$$E[g(X)] = \sum_x g(x) \pi(x) \approx \frac{1}{n} \sum_{i=1}^n g(X_i), \quad X_1, \dots, X_n \sim \pi$$

MCMC Sampling

- ▶ Consider a distribution over (finite) S : $\pi(x) = \frac{b(x)}{Z}$
- ▶ Since this is a distribution, $Z = \sum_{x \in S} b(x)$
- ▶ We assume, we can efficiently calculate $b(x)$ for any x but computation of Z is intractable or computationally expensive
E.g., the Boltzmann distribution: $b(x) = e^{-E(x)/KT}$
- ▶ We want $E[g(X)]$ w.r.t. distribution π (for any g)

$$E[g(X)] = \sum_x g(x) \pi(x) \approx \frac{1}{n} \sum_{i=1}^n g(X_i), \quad X_1, \dots, X_n \sim \pi$$

This is the Monte Carlo method for expectations.

MCMC Sampling

- ▶ Consider a distribution over (finite) S : $\pi(x) = \frac{b(x)}{Z}$
- ▶ Since this is a distribution, $Z = \sum_{x \in S} b(x)$
- ▶ We assume, we can efficiently calculate $b(x)$ for any x but computation of Z is intractable or computationally expensive
E.g., the Boltzmann distribution: $b(x) = e^{-E(x)/KT}$
- ▶ We want $E[g(X)]$ w.r.t. distribution π (for any g)

$$E[g(X)] = \sum_x g(x) \pi(x) \approx \frac{1}{n} \sum_{i=1}^n g(X_i), \quad X_1, \dots, X_n \sim \pi$$

This is the Monte Carlo method for expectations.

- ▶ One way to generate samples is to design an ergodic markov chain with stationary distribution π
 - Markov Chain Monte Carlo sampling

- Suppose $\{X_n\}$ is a an irreducible, aperiodic positive recurrent Markov chain with stationary dist $\pi(x) = \frac{b(x)}{Z}$

- ▶ Suppose $\{X_n\}$ is a an irreducible, aperiodic positive recurrent Markov chain with stationary dist $\pi(x) = \frac{b(x)}{Z}$
- ▶ Then we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n g(X_m) = \sum_x g(x) \pi(x)$$

- ▶ Suppose $\{X_n\}$ is a an irreducible, aperiodic positive recurrent Markov chain with stationary dist $\pi(x) = \frac{b(x)}{Z}$
- ▶ Then we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n g(X_m) = \sum_x g(x) \pi(x)$$

- ▶ hence, if we can design a Markov chain with a given stationary distribution, we can use that to calculate the expectation.

- ▶ We can use the chain to generate samples from distribution π

- ▶ We can use the chain to generate samples from distribution π
We can approximate the expectation as

$$\sum_x g(x)\pi(x) \approx \frac{1}{n} \sum_{i=1}^n g(X_{M+i})$$

- ▶ We can use the chain to generate samples from distribution π
We can approximate the expectation as

$$\sum_x g(x)\pi(x) \approx \frac{1}{n} \sum_{i=1}^n g(X_{M+i})$$

with M is large (so that chain is in steady state)

- ▶ We can use the chain to generate samples from distribution π
We can approximate the expectation as

$$\sum_x g(x)\pi(x) \approx \frac{1}{n} \sum_{i=1}^n g(X_{M+i})$$

with M is large (so that chain is in steady state)

- ▶ When we take sample mean, $\frac{1}{n} \sum_{i=1}^n Z_i$, we want Z_i to be uncorrelated

- ▶ We can use the chain to generate samples from distribution π
We can approximate the expectation as

$$\sum_x g(x)\pi(x) \approx \frac{1}{n} \sum_{i=1}^n g(X_{M+i})$$

with M is large (so that chain is in steady state)

- ▶ When we take sample mean, $\frac{1}{n} \sum_{i=1}^n Z_i$, we want Z_i to be uncorrelated
- ▶ We can, for example, use

$$\sum_x g(x)\pi(x) \approx \frac{1}{n} \sum_{i=1}^n g(X_{M+Ki})$$

- ▶ We can use the chain to generate samples from distribution π
We can approximate the expectation as

$$\sum_x g(x)\pi(x) \approx \frac{1}{n} \sum_{i=1}^n g(X_{M+i})$$

with M is large (so that chain is in steady state)

- ▶ When we take sample mean, $\frac{1}{n} \sum_{i=1}^n Z_i$, we want Z_i to be uncorrelated
- ▶ We can, for example, use

$$\sum_x g(x)\pi(x) \approx \frac{1}{n} \sum_{i=1}^n g(X_{M+Ki})$$

- ▶ For all these, we need to design a Markov chain with π as stationary distribution

- ▶ Let $Q = [q(i, j)]$ be the transition probability matrix of an irreducible Markov chain over S .

- ▶ Let $Q = [q(i, j)]$ be the transition probability matrix of an irreducible Markov chain over S .
- ▶ Q is called the proposal distribution

- ▶ Let $Q = [q(i, j)]$ be the transition probability matrix of an irreducible Markov chain over S .
- ▶ Q is called the proposal distribution
- ▶ We start with arbitray X_0 and generate X_{n+1} , $n = 0, 1, 2, \dots$, iteratively as follows

- ▶ Let $Q = [q(i, j)]$ be the transition probability matrix of an irreducible Markov chain over S .
- ▶ Q is called the proposal distribution
- ▶ We start with arbitray X_0 and generate X_{n+1} , $n = 0, 1, 2, \dots$, iteratively as follows
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$

- ▶ Let $Q = [q(i, j)]$ be the transition probability matrix of an irreducible Markov chain over S .
- ▶ Q is called the proposal distribution
- ▶ We start with arbitrary X_0 and generate X_{n+1} , $n = 0, 1, 2, \dots$, iteratively as follows
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j .

- ▶ Let $Q = [q(i, j)]$ be the transition probability matrix of an irreducible Markov chain over S .
- ▶ Q is called the proposal distribution
- ▶ We start with arbitray X_0 and generate X_{n+1} , $n = 0, 1, 2, \dots$, iteratively as follows
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

- ▶ Let $Q = [q(i, j)]$ be the transition probability matrix of an irreducible Markov chain over S .
- ▶ Q is called the proposal distribution
- ▶ We start with arbitray X_0 and generate X_{n+1} , $n = 0, 1, 2, \dots$, iteratively as follows
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

- ▶ $\alpha(i, j)$ is called the acceptance probability

- ▶ Let $Q = [q(i, j)]$ be the transition probability matrix of an irreducible Markov chain over S .
- ▶ Q is called the proposal distribution
- ▶ We start with arbitrary X_0 and generate X_{n+1} , $n = 0, 1, 2, \dots$, iteratively as follows
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

- ▶ $\alpha(i, j)$ is called the acceptance probability
- ▶ We want to choose $\alpha(i, j)$ to make X_n an ergodic Markov chain with stationary probabilities π

- ▶ The stationary distribution π satisfies (with transition probabilities P)

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad \forall y \in S$$

- ▶ The stationary distribution π satisfies (with transition probabilities P)

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad \forall y \in S$$

- ▶ Suppose there is a distribution $g(\cdot)$ that satisfies

$$g(y) P(y, x) = g(x) P(x, y), \quad \forall x, y \in S$$

- ▶ The stationary distribution π satisfies (with transition probabilities P)

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad \forall y \in S$$

- ▶ Suppose there is a distribution $g(\cdot)$ that satisfies

$$g(y) P(y, x) = g(x) P(x, y), \quad \forall x, y \in S$$

This is called detailed balance

- ▶ The stationary distribution π satisfies (with transition probabilities P)

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad \forall y \in S$$

- ▶ Suppose there is a distribution $g(\cdot)$ that satisfies

$$g(y) P(y, x) = g(x) P(x, y), \quad \forall x, y \in S$$

This is called detailed balance

- ▶ Summing both sides above over x give

$$g(y) = \sum_x g(y) P(y, x) = \sum_x g(x) P(x, y), \quad \forall y$$

- ▶ The stationary distribution π satisfies (with transition probabilities P)

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad \forall y \in S$$

- ▶ Suppose there is a distribution $g(\cdot)$ that satisfies

$$g(y) P(y, x) = g(x) P(x, y), \quad \forall x, y \in S$$

This is called detailed balance

- ▶ Summing both sides above over x give

$$g(y) = \sum_x g(y) P(y, x) = \sum_x g(x) P(x, y), \quad \forall y$$

- ▶ Thus if $g(\cdot)$ satisfies detailed balance, then it must be the stationary distribution

- ▶ The stationary distribution π satisfies (with transition probabilities P)

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad \forall y \in S$$

- ▶ Suppose there is a distribution $g(\cdot)$ that satisfies

$$g(y) P(y, x) = g(x) P(x, y), \quad \forall x, y \in S$$

This is called detailed balance

- ▶ Summing both sides above over x give

$$g(y) = \sum_x g(y) P(y, x) = \sum_x g(x) P(x, y), \quad \forall y$$

- ▶ Thus if $g(\cdot)$ satisfies detailed balance, then it must be the stationary distribution
- ▶ Note that it is not necessary for a stationary distribution to satisfy detailed balance

- Any stationary distribution has to satisfy

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad \forall y \in S$$

- ▶ Any stationary distribution has to satisfy

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad \forall y \in S$$

- ▶ If I can find a π that satisfies

$$\pi(x)P(x, y) = \pi(y)P(y, x), \quad \forall x, y \in S,$$

that would be the stationary distribution

- ▶ Any stationary distribution has to satisfy

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad \forall y \in S$$

- ▶ If I can find a π that satisfies

$$\pi(x)P(x, y) = \pi(y)P(y, x), \quad \forall x, y \in S, x \neq y$$

that would be the stationary distribution

- ▶ Any stationary distribution has to satisfy

$$\pi(y) = \sum_x \pi(x) P(x, y), \quad \forall y \in S$$

- ▶ If I can find a π that satisfies

$$\pi(x)P(x, y) = \pi(y)P(y, x), \quad \forall x, y \in S, x \neq y$$

that would be the stationary distribution

- ▶ This is called detailed balance

- ▶ Recall our algorithm for generating X_n , $n = 0, 1, \dots$

- ▶ Recall our algorithm for generating X_n , $n = 0, 1, \dots$
- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n

- ▶ Recall our algorithm for generating X_n , $n = 0, 1, \dots$
- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$

- ▶ Recall our algorithm for generating X_n , $n = 0, 1, \dots$
- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j .

- ▶ Recall our algorithm for generating X_n , $n = 0, 1, \dots$
- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

- ▶ Recall our algorithm for generating X_n , $n = 0, 1, \dots$
- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

- ▶ Hence the transition probabilities for X_n are

- ▶ Recall our algorithm for generating X_n , $n = 0, 1, \dots$
- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

- ▶ Hence the transition probabilities for X_n are

$$P(i, j) = q(i, j) \alpha(i, j), \quad i \neq j$$

- ▶ Recall our algorithm for generating X_n , $n = 0, 1, \dots$
- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

- ▶ Hence the transition probabilities for X_n are

$$\begin{aligned} P(i, j) &= q(i, j) \alpha(i, j), \quad i \neq j \\ P(i, i) &= q(i, i) + \sum_{j \neq i} q(i, j) (1 - \alpha(i, j)) \end{aligned}$$

- ▶ Recall our algorithm for generating X_n , $n = 0, 1, \dots$
- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

- ▶ Hence the transition probabilities for X_n are

$$\begin{aligned} P(i, j) &= q(i, j) \alpha(i, j), \quad i \neq j \\ P(i, i) &= q(i, i) + \sum_{j \neq i} q(i, j) (1 - \alpha(i, j)) \end{aligned}$$

- ▶ $\pi(i) = b(i)/Z$ is the desired stationary distribution

- ▶ Recall our algorithm for generating X_n , $n = 0, 1, \dots$
- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

- ▶ Hence the transition probabilities for X_n are

$$\begin{aligned} P(i, j) &= q(i, j) \alpha(i, j), \quad i \neq j \\ P(i, i) &= q(i, i) + \sum_{j \neq i} q(i, j) (1 - \alpha(i, j)) \end{aligned}$$

- ▶ $\pi(i) = b(i)/Z$ is the desired stationary distribution
- ▶ So, we can try to satisfy

- ▶ Recall our algorithm for generating X_n , $n = 0, 1, \dots$
- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

- ▶ Hence the transition probabilities for X_n are

$$\begin{aligned} P(i, j) &= q(i, j) \alpha(i, j), \quad i \neq j \\ P(i, i) &= q(i, i) + \sum_{j \neq i} q(i, j) (1 - \alpha(i, j)) \end{aligned}$$

- ▶ $\pi(i) = b(i)/Z$ is the desired stationary distribution
- ▶ So, we can try to satisfy

$$\pi(i) P(i, j) = \pi(j) P(j, i), \quad \forall i, j, i \neq j$$

- ▶ Recall our algorithm for generating X_n , $n = 0, 1, \dots$
- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

- ▶ Hence the transition probabilities for X_n are

$$\begin{aligned} P(i, j) &= q(i, j) \alpha(i, j), \quad i \neq j \\ P(i, i) &= q(i, i) + \sum_{j \neq i} q(i, j) (1 - \alpha(i, j)) \end{aligned}$$

- ▶ $\pi(i) = b(i)/Z$ is the desired stationary distribution
- ▶ So, we can try to satisfy

$$\pi(i) P(i, j) = \pi(j) P(j, i), \quad \forall i, j, i \neq j$$

$$\text{that is, } b(i)q(i, j) \alpha(i, j) = b(j)q(j, i) \alpha(j, i)$$

- We want to satisfy

$$b(i)q(i, j) \alpha(i, j) = b(j)q(j, i) \alpha(j, i)$$

- ▶ We want to satisfy

$$b(i)q(i, j) \alpha(i, j) = b(j)q(j, i) \alpha(j, i)$$

- ▶ Choose

$$\alpha(i, j) = \min \left(\frac{\pi(j)q(j, i)}{\pi(i)q(i, j)}, 1 \right) = \min \left(\frac{b(j)q(j, i)}{b(i)q(i, j)}, 1 \right)$$

- ▶ We want to satisfy

$$b(i)q(i, j) \alpha(i, j) = b(j)q(j, i) \alpha(j, i)$$

- ▶ Choose

$$\alpha(i, j) = \min \left(\frac{\pi(j)q(j, i)}{\pi(i)q(i, j)}, 1 \right) = \min \left(\frac{b(j)q(j, i)}{b(i)q(i, j)}, 1 \right)$$

- ▶ Note that one of $\alpha(i, j)$, $\alpha(j, i)$ is 1

- We want to satisfy

$$b(i)q(i, j) \alpha(i, j) = b(j)q(j, i) \alpha(j, i)$$

- Choose

$$\alpha(i, j) = \min \left(\frac{\pi(j)q(j, i)}{\pi(i)q(i, j)}, 1 \right) = \min \left(\frac{b(j)q(j, i)}{b(i)q(i, j)}, 1 \right)$$

- Note that one of $\alpha(i, j)$, $\alpha(j, i)$ is 1

$$\text{suppose } \alpha(i, j) = \frac{\pi(j)q(j, i)}{\pi(i)q(i, j)} < 1$$

- We want to satisfy

$$b(i)q(i, j) \alpha(i, j) = b(j)q(j, i) \alpha(j, i)$$

- Choose

$$\alpha(i, j) = \min \left(\frac{\pi(j)q(j, i)}{\pi(i)q(i, j)}, 1 \right) = \min \left(\frac{b(j)q(j, i)}{b(i)q(i, j)}, 1 \right)$$

- Note that one of $\alpha(i, j)$, $\alpha(j, i)$ is 1

$$\begin{aligned} \text{suppose } \alpha(i, j) &= \frac{\pi(j)q(j, i)}{\pi(i)q(i, j)} < 1 \\ \Rightarrow \pi(i) q(i, j) \alpha(i, j) &= \pi(j) q(j, i) \end{aligned}$$

- ▶ We want to satisfy

$$b(i)q(i, j) \alpha(i, j) = b(j)q(j, i) \alpha(j, i)$$

- ▶ Choose

$$\alpha(i, j) = \min \left(\frac{\pi(j)q(j, i)}{\pi(i)q(i, j)}, 1 \right) = \min \left(\frac{b(j)q(j, i)}{b(i)q(i, j)}, 1 \right)$$

- ▶ Note that one of $\alpha(i, j)$, $\alpha(j, i)$ is 1

$$\begin{aligned} \text{suppose } \alpha(i, j) &= \frac{\pi(j)q(j, i)}{\pi(i)q(i, j)} < 1 \\ \Rightarrow \pi(i) q(i, j) \alpha(i, j) &= \pi(j) q(j, i) \\ &= \pi(j) q(j, i) \alpha(j, i) \end{aligned}$$

- ▶ We want to satisfy

$$b(i)q(i, j) \alpha(i, j) = b(j)q(j, i) \alpha(j, i)$$

- ▶ Choose

$$\alpha(i, j) = \min \left(\frac{\pi(j)q(j, i)}{\pi(i)q(i, j)}, 1 \right) = \min \left(\frac{b(j)q(j, i)}{b(i)q(i, j)}, 1 \right)$$

- ▶ Note that one of $\alpha(i, j)$, $\alpha(j, i)$ is 1

$$\begin{aligned} \text{suppose } \alpha(i, j) &= \frac{\pi(j)q(j, i)}{\pi(i)q(i, j)} < 1 \\ \Rightarrow \pi(i) q(i, j) \alpha(i, j) &= \pi(j) q(j, i) \\ &= \pi(j) q(j, i) \alpha(j, i) \end{aligned}$$

- ▶ Note that $\pi(i)$ above can be replaced by $b(i)$

Metropolis-Hastings Algorithm

- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n

Metropolis-Hastings Algorithm

- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$

Metropolis-Hastings Algorithm

- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j .

Metropolis-Hastings Algorithm

- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

Metropolis-Hastings Algorithm

- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

Where $Q = [q(i, j)]$ is the transition probabilities of an irreducible chain and

Metropolis-Hastings Algorithm

- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

Where $Q = [q(i, j)]$ is the transition probabilities of an irreducible chain and

$$\alpha(i, j) = \min \left(\frac{\pi(j)q(j, i)}{\pi(i)q(i, j)}, 1 \right)$$

Metropolis-Hastings Algorithm

- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

Where $Q = [q(i, j)]$ is the transition probabilities of an irreducible chain and

$$\alpha(i, j) = \min \left(\frac{\pi(j)q(j, i)}{\pi(i)q(i, j)}, 1 \right)$$

- ▶ Then $\{X_n\}$ would be an irreducible, aperiodic chain with stationary distribution π .

Metropolis-Hastings Algorithm

- ▶ Start with arbitrary X_0 and generate X_{n+1} from X_n
 - ▶ If $X_n = i$, we generate Y with $Pr[Y = k] = q(i, k)$
 - ▶ Let the generated value for Y be j . Set

$$X_{n+1} = \begin{cases} j & \text{with probability } \alpha(i, j) \\ X_n & \text{with probability } 1 - \alpha(i, j) \end{cases}$$

Where $Q = [q(i, j)]$ is the transition probabilities of an irreducible chain and

$$\alpha(i, j) = \min \left(\frac{\pi(j)q(j, i)}{\pi(i)q(i, j)}, 1 \right)$$

- ▶ Then $\{X_n\}$ would be an irreducible, aperiodic chain with stationary distribution π .
- ▶ Q is called the proposal chain and $\alpha(i, j)$ is called acceptance probabilities

- ▶ Consider Boltzmann distribution: $b(x) = e^{-E(x)/KT}$

- ▶ Consider Boltzmann distribution: $b(x) = e^{-E(x)/KT}$
- ▶ Take proposal to be uniform: from any state, we go to all other states with equal probabilities

- ▶ Consider Boltzmann distribution: $b(x) = e^{-E(x)/KT}$
- ▶ Take proposal to be uniform: from any state, we go to all other states with equal probabilities
- ▶ Then,

$$\alpha(x, y) = \min \left(\frac{b(y)}{b(x)}, 1 \right) = \min \left(e^{-(E(y)-E(x))/KT}, 1 \right)$$

- ▶ Consider Boltzmann distribution: $b(x) = e^{-E(x)/KT}$
- ▶ Take proposal to be uniform: from any state, we go to all other states with equal probabilities
- ▶ Then,

$$\alpha(x, y) = \min \left(\frac{b(y)}{b(x)}, 1 \right) = \min \left(e^{-(E(y)-E(x))/KT}, 1 \right)$$

- ▶ In state x you generate a random new state y .
If $E(y) \leq E(x)$ you always go there;

- ▶ Consider Boltzmann distribution: $b(x) = e^{-E(x)/KT}$
- ▶ Take proposal to be uniform: from any state, we go to all other states with equal probabilities
- ▶ Then,

$$\alpha(x, y) = \min \left(\frac{b(y)}{b(x)}, 1 \right) = \min \left(e^{-(E(y)-E(x))/KT}, 1 \right)$$

- ▶ In state x you generate a random new state y .
If $E(y) \leq E(x)$ you always go there;
if $E(y) > E(x)$, accept with probability $e^{-(E(y)-E(x))/KT}$

- ▶ Consider Boltzmann distribution: $b(x) = e^{-E(x)/KT}$
- ▶ Take proposal to be uniform: from any state, we go to all other states with equal probabilities
- ▶ Then,

$$\alpha(x, y) = \min \left(\frac{b(y)}{b(x)}, 1 \right) = \min \left(e^{-(E(y)-E(x))/KT}, 1 \right)$$

- ▶ In state x you generate a random new state y .
If $E(y) \leq E(x)$ you always go there;
if $E(y) > E(x)$, accept with probability $e^{-(E(y)-E(x))/KT}$
- ▶ An interesting way to simulate Boltzmann distribution

- ▶ Consider Boltzmann distribution: $b(x) = e^{-E(x)/KT}$
- ▶ Take proposal to be uniform: from any state, we go to all other states with equal probabilities
- ▶ Then,

$$\alpha(x, y) = \min \left(\frac{b(y)}{b(x)}, 1 \right) = \min \left(e^{-(E(y)-E(x))/KT}, 1 \right)$$

- ▶ In state x you generate a random new state y .
If $E(y) \leq E(x)$ you always go there;
if $E(y) > E(x)$, accept with probability $e^{-(E(y)-E(x))/KT}$
- ▶ An interesting way to simulate Boltzmann distribution
- ▶ We could have chosen Q to be 'uniform over neighbours'

► Suppose $E : S \rightarrow \mathfrak{R}$ is some function.

- ▶ Suppose $E : S \rightarrow \mathbb{R}$ is some function.
- ▶ We want to find $x \in S$ where E is *globally* minimized.

- ▶ Suppose $E : S \rightarrow \mathbb{R}$ is some function.
- ▶ We want to find $x \in S$ where E is *globally* minimized.
- ▶ A gradient descent type method tries to find a locally minimizing direction and hence gives only a ‘local’ minimum.

- ▶ Suppose $E : S \rightarrow \mathbb{R}$ is some function.
- ▶ We want to find $x \in S$ where E is *globally* minimized.
- ▶ A gradient descent type method tries to find a locally minimizing direction and hence gives only a ‘local’ minimum.
- ▶ The Metropolis-Hastings algorithm gives another view point on how such optimization problems can be handled.

- ▶ Suppose $E : S \rightarrow \mathbb{R}$ is some function.
- ▶ We want to find $x \in S$ where E is *globally* minimized.
- ▶ A gradient descent type method tries to find a locally minimizing direction and hence gives only a ‘local’ minimum.
- ▶ The Metropolis-Hastings algorithm gives another view point on how such optimization problems can be handled.
- ▶ We can think of E as the energy function in a Boltzmann distribution

- ▶ Let $b(x) = e^{-E(x)/T}$ where T is a parameter called 'temperature'

- ▶ Let $b(x) = e^{-E(x)/T}$ where T is a parameter called 'temparature'
- ▶ $\{X_n\}$ be Markov chain with stationary dist $\pi(x) = \frac{b(x)}{Z}$

- ▶ Let $b(x) = e^{-E(x)/T}$ where T is a parameter called 'temperature'
- ▶ $\{X_n\}$ be Markov chain with stationary dist $\pi(x) = \frac{b(x)}{Z}$
- ▶ We can find relative occupation of different states by the chain by collecting statistics during steady state

- ▶ Let $b(x) = e^{-E(x)/T}$ where T is a parameter called 'temparature'
- ▶ $\{X_n\}$ be Markov chain with stationary dist $\pi(x) = \frac{b(x)}{Z}$
- ▶ We can find relative occupation of different states by the chain by collecting statistics during steady state
- ▶ We know

$$\frac{\pi(x_1)}{\pi(x_2)} = \frac{b(x_1)}{b(x_2)} = e^{-(E(x_1)-E(x_2))/T}$$

- ▶ Let $b(x) = e^{-E(x)/T}$ where T is a parameter called 'temperature'
- ▶ $\{X_n\}$ be Markov chain with stationary dist $\pi(x) = \frac{b(x)}{Z}$
- ▶ We can find relative occupation of different states by the chain by collecting statistics during steady state
- ▶ We know

$$\frac{\pi(x_1)}{\pi(x_2)} = \frac{b(x_1)}{b(x_2)} = e^{-(E(x_1)-E(x_2))/T}$$

- ▶ We spend more time in global minimum

- ▶ Let $b(x) = e^{-E(x)/T}$ where T is a parameter called 'temperature'
- ▶ $\{X_n\}$ be Markov chain with stationary dist $\pi(x) = \frac{b(x)}{Z}$
- ▶ We can find relative occupation of different states by the chain by collecting statistics during steady state
- ▶ We know

$$\frac{\pi(x_1)}{\pi(x_2)} = \frac{b(x_1)}{b(x_2)} = e^{-(E(x_1)-E(x_2))/T}$$

- ▶ We spend more time in global minimum
We can increase the relative fraction of time spent in global minimum by decreasing T

- ▶ Let $b(x) = e^{-E(x)/T}$ where T is a parameter called 'temperature'
- ▶ $\{X_n\}$ be Markov chain with stationary dist $\pi(x) = \frac{b(x)}{Z}$
- ▶ We can find relative occupation of different states by the chain by collecting statistics during steady state
- ▶ We know

$$\frac{\pi(x_1)}{\pi(x_2)} = \frac{b(x_1)}{b(x_2)} = e^{-(E(x_1)-E(x_2))/T}$$

- ▶ We spend more time in global minimum
We can increase the relative fraction of time spent in global minimum by decreasing T (There is a price to pay!)

- ▶ Let $b(x) = e^{-E(x)/T}$ where T is a parameter called 'temparature'
- ▶ $\{X_n\}$ be Markov chain with stationary dist $\pi(x) = \frac{b(x)}{Z}$
- ▶ We can find relative occupation of different states by the chain by collecting statistics during steady state
- ▶ We know

$$\frac{\pi(x_1)}{\pi(x_2)} = \frac{b(x_1)}{b(x_2)} = e^{-(E(x_1)-E(x_2))/T}$$

- ▶ We spend more time in global minimum
We can increase the relative fraction of time spent in global minimum by decreasing T (There is a price to pay!)
- ▶ Gives rise to interesting optimization technique called simulated annealing

- ▶ In most applications of MCMC, $x \in \mathcal{S}$ is a vector.

- ▶ In most applications of MCMC, $x \in \mathcal{S}$ is a vector.
- ▶ One normally changes one component at a time. That is how neighbours can be defined

- ▶ In most applications of MCMC, $x \in \mathcal{S}$ is a vector.
- ▶ One normally changes one component at a time. That is how neighbours can be defined
- ▶ A special case of proposal distribution is the conditional distribution.

- ▶ In most applications of MCMC, $x \in \mathcal{S}$ is a vector.
- ▶ One normally changes one component at a time. That is how neighbours can be defined
- ▶ A special case of proposal distribution is the conditional distribution.
- ▶ Suppose $X = (X_1, \dots, X_N)$. To propose a value for X_i , we use $f_{X_i|X_{-i}}$

- ▶ In most applications of MCMC, $x \in \mathcal{S}$ is a vector.
- ▶ One normally changes one component at a time. That is how neighbours can be defined
- ▶ A special case of proposal distribution is the conditional distribution.
- ▶ Suppose $X = (X_1, \dots, X_N)$. To propose a value for X_i , we use $f_{X_i|X_{-i}}$ ($= f_{X_i|X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_N}$)

- ▶ In most applications of MCMC, $x \in \mathcal{S}$ is a vector.
- ▶ One normally changes one component at a time. That is how neighbours can be defined
- ▶ A special case of proposal distribution is the conditional distribution.
- ▶ Suppose $X = (X_1, \dots, X_N)$. To propose a value for X_i , we use $f_{X_i|X_{-i}}$ ($= f_{X_i|X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_N}$)
- ▶ Here the conditional distribution is calculated using the target π as the joint distribution.

- ▶ In most applications of MCMC, $x \in S$ is a vector.
- ▶ One normally changes one component at a time. That is how neighbours can be defined
- ▶ A special case of proposal distribution is the conditional distribution.
- ▶ Suppose $X = (X_1, \dots, X_N)$. To propose a value for X_i , we use $f_{X_i|X_{-i}}$ ($= f_{X_i|X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_N}$)
- ▶ Here the conditional distribution is calculated using the target π as the joint distribution.
- ▶ With such a proposal distribution, one can show that $\alpha(i, j)$ is always 1

- ▶ In most applications of MCMC, $x \in \mathcal{S}$ is a vector.
- ▶ One normally changes one component at a time. That is how neighbours can be defined
- ▶ A special case of proposal distribution is the conditional distribution.
- ▶ Suppose $X = (X_1, \dots, X_N)$. To propose a value for X_i , we use $f_{X_i|X_{-i}} (= f_{X_i|X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_N})$
- ▶ Here the conditional distribution is calculated using the target π as the joint distribution.
- ▶ With such a proposal distribution, one can show that $\alpha(i, j)$ is always 1
- ▶ This is known as Gibbs sampling