

# Review of basic probability

We use the following Notation:

# Review of basic probability

We use the following Notation:

- ▶ Sample space –  $\Omega$

# Review of basic probability

We use the following Notation:

- ▶ Sample space –  $\Omega$   
Elements of  $\Omega$  are the outcomes of the random experiment

# Review of basic probability

We use the following Notation:

- ▶ Sample space –  $\Omega$

Elements of  $\Omega$  are the outcomes of the random experiment

We write  $\Omega = \{\omega_1, \omega_2, \dots\}$  when it is countable

# Review of basic probability

We use the following Notation:

- ▶ Sample space –  $\Omega$

Elements of  $\Omega$  are the outcomes of the random experiment

We write  $\Omega = \{\omega_1, \omega_2, \dots\}$  when it is countable

- ▶ An event is, by definition, a subset of  $\Omega$

# Review of basic probability

We use the following Notation:

- ▶ Sample space –  $\Omega$   
Elements of  $\Omega$  are the outcomes of the random experiment  
We write  $\Omega = \{\omega_1, \omega_2, \dots\}$  when it is countable
- ▶ An event is, by definition, a subset of  $\Omega$
- ▶ Set of all possible events –  $\mathcal{F} \subset 2^\Omega$  (power set of  $\Omega$ )  
**Each event is a subset of  $\Omega$**

# Review of basic probability

We use the following Notation:

- ▶ Sample space –  $\Omega$   
Elements of  $\Omega$  are the outcomes of the random experiment  
We write  $\Omega = \{\omega_1, \omega_2, \dots\}$  when it is countable
- ▶ An event is, by definition, a subset of  $\Omega$
- ▶ Set of all possible events –  $\mathcal{F} \subset 2^\Omega$  (power set of  $\Omega$ )  
**Each event is a subset of  $\Omega$**   
For now, we take  $\mathcal{F} = 2^\Omega$  (power set of  $\Omega$ )

# Probability axioms

Probability (or probability measure) is a function that assigns a number to each event and satisfies some properties.

$$P : \mathcal{F} \rightarrow \mathbb{R}, \quad \mathcal{F} \subset 2^\Omega$$



# Probability axioms

Probability (or probability measure) is a function that assigns a number to each event and satisfies some properties.

$$P : \mathcal{F} \rightarrow \mathbb{R}, \quad \mathcal{F} \subset 2^\Omega$$

$$\text{A1 } P(A) \geq 0, \forall A \in \mathcal{F}$$

$$\text{A2 } P(\Omega) = 1$$

$$\text{A3 } \text{If } A_i \cap A_j = \phi, \forall i \neq j \text{ then } P(\cup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} P(A_i)$$

# Some consequences of the axioms

►  $0 \leq P(A) \leq 1$

# Some consequences of the axioms

- ▶  $0 \leq P(A) \leq 1$
- ▶  $P(A^c) = 1 - P(A)$

# Some consequences of the axioms

- ▶  $0 \leq P(A) \leq 1$
- ▶  $P(A^c) = 1 - P(A)$
- ▶ If  $A \subset B$  then,  $P(A) \leq P(B)$

# Some consequences of the axioms

- ▶  $0 \leq P(A) \leq 1$
- ▶  $P(A^c) = 1 - P(A)$
- ▶ If  $A \subset B$  then,  $P(A) \leq P(B)$
- ▶ If  $A \subset B$  then,  $P(B - A) = P(B) - P(A)$

# Some consequences of the axioms

- ▶  $0 \leq P(A) \leq 1$
- ▶  $P(A^c) = 1 - P(A)$
- ▶ If  $A \subset B$  then,  $P(A) \leq P(B)$
- ▶ If  $A \subset B$  then,  $P(B - A) = P(B) - P(A)$
- ▶  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

# Case of finite $\Omega$ – Example

## Case of finite $\Omega$ – Example

- ▶ Let  $\Omega = \{\omega_1, \dots, \omega_n\}$ ,  $\mathcal{F} = 2^\Omega$ , and  $P$  is specified through ‘equally likely’ assumption.



## Case of finite $\Omega$ – Example

- ▶ Let  $\Omega = \{\omega_1, \dots, \omega_n\}$ ,  $\mathcal{F} = 2^\Omega$ , and  $P$  is specified through ‘equally likely’ assumption.
- ▶ That is,  $P(\{\omega_i\}) = \frac{1}{n}$ .

## Case of finite $\Omega$ – Example

- ▶ Let  $\Omega = \{\omega_1, \dots, \omega_n\}$ ,  $\mathcal{F} = 2^\Omega$ , and  $P$  is specified through ‘equally likely’ assumption.
- ▶ That is,  $P(\{\omega_i\}) = \frac{1}{n}$ . (Note the notation)

## Case of finite $\Omega$ – Example

- ▶ Let  $\Omega = \{\omega_1, \dots, \omega_n\}$ ,  $\mathcal{F} = 2^\Omega$ , and  $P$  is specified through ‘equally likely’ assumption.
- ▶ That is,  $P(\{\omega_i\}) = \frac{1}{n}$ . (Note the notation)
- ▶ Suppose  $A = \{\omega_1, \omega_2, \omega_3\}$ . Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\})$$

## Case of finite $\Omega$ – Example

- ▶ Let  $\Omega = \{\omega_1, \dots, \omega_n\}$ ,  $\mathcal{F} = 2^\Omega$ , and  $P$  is specified through ‘equally likely’ assumption.
- ▶ That is,  $P(\{\omega_i\}) = \frac{1}{n}$ . (Note the notation)
- ▶ Suppose  $A = \{\omega_1, \omega_2, \omega_3\}$ . Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\}) = \sum_{i=1}^3 P(\{\omega_i\})$$

## Case of finite $\Omega$ – Example

- ▶ Let  $\Omega = \{\omega_1, \dots, \omega_n\}$ ,  $\mathcal{F} = 2^\Omega$ , and  $P$  is specified through ‘equally likely’ assumption.
- ▶ That is,  $P(\{\omega_i\}) = \frac{1}{n}$ . (Note the notation)
- ▶ Suppose  $A = \{\omega_1, \omega_2, \omega_3\}$ . Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\}) = \sum_{i=1}^3 P(\{\omega_i\}) = \frac{3}{n} = \frac{|A|}{|\Omega|}$$

## Case of finite $\Omega$ – Example

- ▶ Let  $\Omega = \{\omega_1, \dots, \omega_n\}$ ,  $\mathcal{F} = 2^\Omega$ , and  $P$  is specified through ‘equally likely’ assumption.
- ▶ That is,  $P(\{\omega_i\}) = \frac{1}{n}$ . (Note the notation)
- ▶ Suppose  $A = \{\omega_1, \omega_2, \omega_3\}$ . Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\}) = \sum_{i=1}^3 P(\{\omega_i\}) = \frac{3}{n} = \frac{|A|}{|\Omega|}$$

- ▶ We can easily see this to be true for any event,  $A$ .

## Case of finite $\Omega$ – Example

- ▶ Let  $\Omega = \{\omega_1, \dots, \omega_n\}$ ,  $\mathcal{F} = 2^\Omega$ , and  $P$  is specified through ‘equally likely’ assumption.
- ▶ That is,  $P(\{\omega_i\}) = \frac{1}{n}$ . (Note the notation)
- ▶ Suppose  $A = \{\omega_1, \omega_2, \omega_3\}$ . Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\}) = \sum_{i=1}^3 P(\{\omega_i\}) = \frac{3}{n} = \frac{|A|}{|\Omega|}$$

- ▶ We can easily see this to be true for any event,  $A$ .
- ▶ This is the usual familiar formula: number of favourable outcomes by total number of outcomes.

## Case of finite $\Omega$ – Example

- ▶ Let  $\Omega = \{\omega_1, \dots, \omega_n\}$ ,  $\mathcal{F} = 2^\Omega$ , and  $P$  is specified through ‘equally likely’ assumption.
- ▶ That is,  $P(\{\omega_i\}) = \frac{1}{n}$ . (Note the notation)
- ▶ Suppose  $A = \{\omega_1, \omega_2, \omega_3\}$ . Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\}) = \sum_{i=1}^3 P(\{\omega_i\}) = \frac{3}{n} = \frac{|A|}{|\Omega|}$$

- ▶ We can easily see this to be true for any event,  $A$ .
- ▶ This is the usual familiar formula: number of favourable outcomes by total number of outcomes.
- ▶ Thus, ‘equally likely’ is one way of specifying the probability function (in case of finite  $\Omega$ ).



## Case of finite $\Omega$ – Example

- ▶ Let  $\Omega = \{\omega_1, \dots, \omega_n\}$ ,  $\mathcal{F} = 2^\Omega$ , and  $P$  is specified through ‘equally likely’ assumption.
- ▶ That is,  $P(\{\omega_i\}) = \frac{1}{n}$ . (Note the notation)
- ▶ Suppose  $A = \{\omega_1, \omega_2, \omega_3\}$ . Then

$$P(A) = P(\{\omega_1\} \cup \{\omega_2\} \cup \{\omega_3\}) = \sum_{i=1}^3 P(\{\omega_i\}) = \frac{3}{n} = \frac{|A|}{|\Omega|}$$

- ▶ We can easily see this to be true for any event,  $A$ .
- ▶ This is the usual familiar formula: number of favourable outcomes by total number of outcomes.
- ▶ Thus, ‘equally likely’ is one way of specifying the probability function (in case of finite  $\Omega$ ).
- ▶ An obvious point worth remembering: specifying  $P$  for singleton events fixes it for all other events.

# Case of Countably infinite $\Omega$

- ▶ Let  $\Omega = \{\omega_1, \omega_2, \dots\}$ .

# Case of Countably infinite $\Omega$

- ▶ Let  $\Omega = \{\omega_1, \omega_2, \dots\}$ .
- ▶ Once again, any  $A \subset \Omega$  can be written as mutually exclusive union of singleton sets.

# Case of Countably infinite $\Omega$

- ▶ Let  $\Omega = \{\omega_1, \omega_2, \dots\}$ .
- ▶ Once again, any  $A \subset \Omega$  can be written as mutually exclusive union of singleton sets.
- ▶ Let  $q_i, i = 1, 2, \dots$  be numbers such that  $q_i \geq 0$  and  $\sum_i q_i = 1$ .

## Case of Countably infinite $\Omega$

- ▶ Let  $\Omega = \{\omega_1, \omega_2, \dots\}$ .
- ▶ Once again, any  $A \subset \Omega$  can be written as mutually exclusive union of singleton sets.
- ▶ Let  $q_i, i = 1, 2, \dots$  be numbers such that  $q_i \geq 0$  and  $\sum_i q_i = 1$ .
- ▶ We can now set  $P(\{\omega_i\}) = q_i, i = 1, 2, \dots$ .

## Case of Countably infinite $\Omega$

- ▶ Let  $\Omega = \{\omega_1, \omega_2, \dots\}$ .
- ▶ Once again, any  $A \subset \Omega$  can be written as mutually exclusive union of singleton sets.
- ▶ Let  $q_i, i = 1, 2, \dots$  be numbers such that  $q_i \geq 0$  and  $\sum_i q_i = 1$ .
- ▶ We can now set  $P(\{\omega_i\}) = q_i, i = 1, 2, \dots$ .  
(Assumptions on  $q_i$  needed to satisfy  $P(A) \geq 0$  and  $P(\Omega) = 1$ ).

# Case of Countably infinite $\Omega$

- ▶ Let  $\Omega = \{\omega_1, \omega_2, \dots\}$ .
- ▶ Once again, any  $A \subset \Omega$  can be written as mutually exclusive union of singleton sets.
- ▶ Let  $q_i, i = 1, 2, \dots$  be numbers such that  $q_i \geq 0$  and  $\sum_i q_i = 1$ .
- ▶ We can now set  $P(\{\omega_i\}) = q_i, i = 1, 2, \dots$ .  
(Assumptions on  $q_i$  needed to satisfy  $P(A) \geq 0$  and  $P(\Omega) = 1$ ).
- ▶ This fixes  $P$  for all events:  $P(A) = \sum_{\omega \in A} P(\{\omega\})$

## Case of Countably infinite $\Omega$

- ▶ Let  $\Omega = \{\omega_1, \omega_2, \dots\}$ .
- ▶ Once again, any  $A \subset \Omega$  can be written as mutually exclusive union of singleton sets.
- ▶ Let  $q_i, i = 1, 2, \dots$  be numbers such that  $q_i \geq 0$  and  $\sum_i q_i = 1$ .
- ▶ We can now set  $P(\{\omega_i\}) = q_i, i = 1, 2, \dots$ .  
(Assumptions on  $q_i$  needed to satisfy  $P(A) \geq 0$  and  $P(\Omega) = 1$ ).
- ▶ This fixes  $P$  for all events:  $P(A) = \sum_{\omega \in A} P(\{\omega\})$
- ▶ This is how we normally define a probability measure on countably infinite  $\Omega$ .



# Case of Countably infinite $\Omega$

- ▶ Let  $\Omega = \{\omega_1, \omega_2, \dots\}$ .
- ▶ Once again, any  $A \subset \Omega$  can be written as mutually exclusive union of singleton sets.
- ▶ Let  $q_i, i = 1, 2, \dots$  be numbers such that  $q_i \geq 0$  and  $\sum_i q_i = 1$ .
- ▶ We can now set  $P(\{\omega_i\}) = q_i, i = 1, 2, \dots$ .  
(Assumptions on  $q_i$  needed to satisfy  $P(A) \geq 0$  and  $P(\Omega) = 1$ ).
- ▶ This fixes  $P$  for all events:  $P(A) = \sum_{\omega \in A} P(\{\omega\})$
- ▶ This is how we normally define a probability measure on countably infinite  $\Omega$ .
- ▶ This can be done for finite  $\Omega$  too.

## Example: countably infinite $\Omega$

- ▶ Let  $\Omega = \{0, 1, 2, \dots\}$

## Example: countably infinite $\Omega$

- ▶ Let  $\Omega = \{0, 1, 2, \dots\}$
- ▶ Let  $q_i = (1 - p)^i p$  for some  $p$ ,  $0 < p < 1$ .

## Example: countably infinite $\Omega$

- ▶ Let  $\Omega = \{0, 1, 2, \dots\}$
- ▶ Let  $q_i = (1 - p)^i p$  for some  $p$ ,  $0 < p < 1$ .
- ▶ Easy to see:  $q_i \geq 0$  and  $\sum_{i=0}^{\infty} q_i = 1$ .

## Example: countably infinite $\Omega$

- ▶ Let  $\Omega = \{0, 1, 2, \dots\}$
- ▶ Let  $q_i = (1 - p)^i p$  for some  $p$ ,  $0 < p < 1$ .
- ▶ Easy to see:  $q_i \geq 0$  and  $\sum_{i=0}^{\infty} q_i = 1$ .
- ▶ We can assign  $P(\{k\}) = (1 - p)^k p$ ,  $k = 0, 1, \dots$

## Example: countably infinite $\Omega$

- ▶ Let  $\Omega = \{0, 1, 2, \dots\}$
- ▶ Let  $q_i = (1 - p)^i p$  for some  $p$ ,  $0 < p < 1$ .
- ▶ Easy to see:  $q_i \geq 0$  and  $\sum_{i=0}^{\infty} q_i = 1$ .
- ▶ We can assign  $P(\{k\}) = (1 - p)^k p$ ,  $k = 0, 1, \dots$
- ▶ Consider a random experiment of tossing a biased coin repeatedly till we get a head. We take the outcome of the experiment to be the number of tails we had before the first head.

## Example: countably infinite $\Omega$

- ▶ Let  $\Omega = \{0, 1, 2, \dots\}$
- ▶ Let  $q_i = (1 - p)^i p$  for some  $p$ ,  $0 < p < 1$ .
- ▶ Easy to see:  $q_i \geq 0$  and  $\sum_{i=0}^{\infty} q_i = 1$ .
- ▶ We can assign  $P(\{k\}) = (1 - p)^k p$ ,  $k = 0, 1, \dots$
- ▶ Consider a random experiment of tossing a biased coin repeatedly till we get a head. We take the outcome of the experiment to be the number of tails we had before the first head.
- ▶ A (reasonable) probability assignment is:

$$P(\{k\}) = (1 - p)^k p, k = 0, 1, \dots$$

where  $p$  is the probability of head and  $0 < p < 1$ .  
(We assume you understand the idea of ‘independent’ tosses here).

# Case of uncountably infinite $\Omega$

- ▶ We would mostly be considering only the cases where  $\Omega$  is a subset of  $\Re^d$  for some  $d$ .



# Case of uncountably infinite $\Omega$

- ▶ We would mostly be considering only the cases where  $\Omega$  is a subset of  $\Re^d$  for some  $d$ .
- ▶ Note that now an event need not be a countable union of singleton sets.

# Case of uncountably infinite $\Omega$

- ▶ We would mostly be considering only the cases where  $\Omega$  is a subset of  $\Re^d$  for some  $d$ .
- ▶ Note that now an event need not be a countable union of singleton sets.
- ▶ For now we would only consider a simple intuitive extension of the 'equally likely' idea.

# Case of uncountably infinite $\Omega$

- ▶ We would mostly be considering only the cases where  $\Omega$  is a subset of  $\Re^d$  for some  $d$ .
- ▶ Note that now an event need not be a countable union of singleton sets.
- ▶ For now we would only consider a simple intuitive extension of the 'equally likely' idea.
- ▶ Suppose  $\Omega$  is a finite interval of  $\Re$ . Then we will take  $P(A) = \frac{m(A)}{m(\Omega)}$  where  $m(A)$  is length of the set  $A$ .

# Case of uncountably infinite $\Omega$

- ▶ We would mostly be considering only the cases where  $\Omega$  is a subset of  $\Re^d$  for some  $d$ .
- ▶ Note that now an event need not be a countable union of singleton sets.
- ▶ For now we would only consider a simple intuitive extension of the 'equally likely' idea.
- ▶ Suppose  $\Omega$  is a finite interval of  $\Re$ . Then we will take  $P(A) = \frac{m(A)}{m(\Omega)}$  where  $m(A)$  is length of the set  $A$ .
- ▶ We can use this in higher dimensions also by taking  $m(\cdot)$  to be an appropriate 'measure' of a set.

# Case of uncountably infinite $\Omega$

- ▶ We would mostly be considering only the cases where  $\Omega$  is a subset of  $\mathbb{R}^d$  for some  $d$ .
- ▶ Note that now an event need not be a countable union of singleton sets.
- ▶ For now we would only consider a simple intuitive extension of the 'equally likely' idea.
- ▶ Suppose  $\Omega$  is a finite interval of  $\mathbb{R}$ . Then we will take  $P(A) = \frac{m(A)}{m(\Omega)}$  where  $m(A)$  is length of the set  $A$ .
- ▶ We can use this in higher dimensions also by taking  $m(\cdot)$  to be an appropriate 'measure' of a set.
- ▶ For example, in  $\mathbb{R}^2$ ,  $m(A)$  denotes area of  $A$ , in  $\mathbb{R}^3$  it would be volume and so on.

# Case of uncountably infinite $\Omega$

- ▶ We would mostly be considering only the cases where  $\Omega$  is a subset of  $\mathbb{R}^d$  for some  $d$ .
- ▶ Note that now an event need not be a countable union of singleton sets.
- ▶ For now we would only consider a simple intuitive extension of the 'equally likely' idea.
- ▶ Suppose  $\Omega$  is a finite interval of  $\mathbb{R}$ . Then we will take  $P(A) = \frac{m(A)}{m(\Omega)}$  where  $m(A)$  is length of the set  $A$ .
- ▶ We can use this in higher dimensions also by taking  $m(\cdot)$  to be an appropriate 'measure' of a set.
- ▶ For example, in  $\mathbb{R}^2$ ,  $m(A)$  denotes area of  $A$ , in  $\mathbb{R}^3$  it would be volume and so on.

(There are many issues that need more attention here).

## Example: Uncountably infinite $\Omega$

**Problem:** A rod of unit length is broken at two random points. What is the probability that the three pieces so formed would make a triangle.

## Example: Uncountably infinite $\Omega$

**Problem:** A rod of unit length is broken at two random points. What is the probability that the three pieces so formed would make a triangle.

- ▶ Let us take left end of the rod as origin and let  $x, y$  denote the two successive points where the rod is broken.



## Example: Uncountably infinite $\Omega$

**Problem:** A rod of unit length is broken at two random points. What is the probability that the three pieces so formed would make a triangle.

- ▶ Let us take left end of the rod as origin and let  $x, y$  denote the two successive points where the rod is broken.
- ▶ Then the random experiment is picking two numbers  $x, y$  with  $0 < x < y < 1$ .

## Example: Uncountably infinite $\Omega$

**Problem:** A rod of unit length is broken at two random points. What is the probability that the three pieces so formed would make a triangle.

- ▶ Let us take left end of the rod as origin and let  $x, y$  denote the two successive points where the rod is broken.
- ▶ Then the random experiment is picking two numbers  $x, y$  with  $0 < x < y < 1$ .
- ▶ We can take  $\Omega = \{(x, y) : 0 < x < y < 1\} \subset \mathbb{R}^2$ .

## Example: Uncountably infinite $\Omega$

**Problem:** A rod of unit length is broken at two random points. What is the probability that the three pieces so formed would make a triangle.

- ▶ Let us take left end of the rod as origin and let  $x, y$  denote the two successive points where the rod is broken.
- ▶ Then the random experiment is picking two numbers  $x, y$  with  $0 < x < y < 1$ .
- ▶ We can take  $\Omega = \{(x, y) : 0 < x < y < 1\} \subset \mathbb{R}^2$ .
- ▶ For the pieces to make a triangle, sum of lengths of any two should be more than the third.

- ▶ The lengths are:  $x, (y - x), (1 - y)$ .

- ▶ The lengths are:  $x, (y - x), (1 - y)$ . So we need

- The lengths are:  $x, (y - x), (1 - y)$ . So we need

$$x + (y - x) > (1 - y) \Rightarrow y > 0.5$$

- The lengths are:  $x, (y - x), (1 - y)$ . So we need

$$x + (y - x) > (1 - y) \Rightarrow y > 0.5$$

$$x + (1 - y) > (y - x) \Rightarrow y < x + 0.5;$$

- The lengths are:  $x, (y - x), (1 - y)$ . So we need

$$x + (y - x) > (1 - y) \Rightarrow y > 0.5$$

$$x + (1 - y) > (y - x) \Rightarrow y < x + 0.5;$$

$$(y - x) + 1 - y > x \Rightarrow x < 0.5$$



- The lengths are:  $x, (y - x), (1 - y)$ . So we need

$$x + (y - x) > (1 - y) \Rightarrow y > 0.5$$

$$x + (1 - y) > (y - x) \Rightarrow y < x + 0.5;$$

$$(y - x) + 1 - y > x \Rightarrow x < 0.5$$

- So the event of interest is:

$$A = \{(x, y) : y > 0.5; x < 0.5; y < x + 0.5, 0 < x, y < 1\}$$

► We have


$$\Omega = \{(x, y) : 0 < x < y < 1\}$$

$$A = \{(x, y) \in \Omega : y > 0.5; x < 0.5; y < x + 0.5\}$$

► We have

$$\Omega = \{(x, y) : 0 < x < y < 1\}$$

$$A = \{(x, y) \in \Omega : y > 0.5; x < 0.5; y < x + 0.5\}$$




► We can visualize it as follows

- We have

$$\Omega = \{(x, y) : 0 < x < y < 1\}$$

$$A = \{(x, y) \in \Omega : y > 0.5; x < 0.5; y < x + 0.5\}$$



- We can visualize it as follows
- The required probability is area of  $A$  divided by area of  $\Omega$  which gives the answer as 0.25

- Everything we do in probability theory is always in reference to an underlying probability space:  $(\Omega, \mathcal{F}, P)$  where

- ▶ Everything we do in probability theory is always in reference to an underlying probability space:  $(\Omega, \mathcal{F}, P)$  where
  - ▶  $\Omega$  is the sample space

- ▶ Everything we do in probability theory is always in reference to an underlying probability space:  $(\Omega, \mathcal{F}, P)$  where
  - ▶  $\Omega$  is the sample space
  - ▶  $\mathcal{F} \subset 2^\Omega$  set of events; each event is a subset of  $\Omega$

- ▶ Everything we do in probability theory is always in reference to an underlying probability space:  $(\Omega, \mathcal{F}, P)$  where
  - ▶  $\Omega$  is the sample space
  - ▶  $\mathcal{F} \subset 2^\Omega$  set of events; each event is a subset of  $\Omega$
  - ▶  $P : \mathcal{F} \rightarrow [0, 1]$  is a probability (measure) that assigns a number between 0 and 1 to every event (satisfying the three axioms).



- ▶ Everything we do in probability theory is always in reference to an underlying probability space:  $(\Omega, \mathcal{F}, P)$  where
  - ▶  $\Omega$  is the sample space
  - ▶  $\mathcal{F} \subset 2^\Omega$  set of events; each event is a subset of  $\Omega$
  - ▶  $P : \mathcal{F} \rightarrow [0, 1]$  is a probability (measure) that assigns a number between 0 and 1 to every event (satisfying the three axioms).
- ▶ We saw some examples

- ▶ Everything we do in probability theory is always in reference to an underlying probability space:  $(\Omega, \mathcal{F}, P)$  where
  - ▶  $\Omega$  is the sample space
  - ▶  $\mathcal{F} \subset 2^\Omega$  set of events; each event is a subset of  $\Omega$
  - ▶  $P : \mathcal{F} \rightarrow [0, 1]$  is a probability (measure) that assigns a number between 0 and 1 to every event (satisfying the three axioms).
- ▶ We saw some examples
- ▶ Given an  $\Omega$  and  $\mathcal{F}$ , there can be many  $P$  that satisfy the axioms

- ▶ Everything we do in probability theory is always in reference to an underlying probability space:  $(\Omega, \mathcal{F}, P)$  where
  - ▶  $\Omega$  is the sample space
  - ▶  $\mathcal{F} \subset 2^\Omega$  set of events; each event is a subset of  $\Omega$
  - ▶  $P : \mathcal{F} \rightarrow [0, 1]$  is a probability (measure) that assigns a number between 0 and 1 to every event (satisfying the three axioms).
- ▶ We saw some examples
- ▶ Given an  $\Omega$  and  $\mathcal{F}$ , there can be many  $P$  that satisfy the axioms – many "probability models"

# Conditional Probability

# Conditional Probability

- ▶ Let  $B$  be an event with  $P(B) > 0$ . We define conditional probability, conditioned on  $B$ , of any event,  $A$ , as

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)} = \frac{P(AB)}{P(B)}$$

# Conditional Probability

- ▶ Let  $B$  be an event with  $P(B) > 0$ . We define conditional probability, conditioned on  $B$ , of any event,  $A$ , as

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)} = \frac{P(AB)}{P(B)}$$

- ▶ The above is a notation.

# Conditional Probability

- ▶ Let  $B$  be an event with  $P(B) > 0$ . We define conditional probability, conditioned on  $B$ , of any event,  $A$ , as

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)} = \frac{P(AB)}{P(B)}$$

- ▶ The above is a notation. “ $A \mid B$ ” does not represent any set operation!

# Conditional Probability

- ▶ Let  $B$  be an event with  $P(B) > 0$ . We define conditional probability, conditioned on  $B$ , of any event,  $A$ , as

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)} = \frac{P(AB)}{P(B)}$$

- ▶ The above is a notation. “ $A \mid B$ ” does not represent any set operation! (This is an abuse of notation!)



# Conditional Probability

- ▶ Let  $B$  be an event with  $P(B) > 0$ . We define conditional probability, conditioned on  $B$ , of any event,  $A$ , as

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)} = \frac{P(AB)}{P(B)}$$

- ▶ The above is a notation. “ $A \mid B$ ” does not represent any set operation! (This is an abuse of notation!)
- ▶ Given a  $B$ , conditional probability is a new probability assignment to any event.

# Conditional Probability

- ▶ Let  $B$  be an event with  $P(B) > 0$ . We define conditional probability, conditioned on  $B$ , of any event,  $A$ , as

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)} = \frac{P(AB)}{P(B)}$$

- ▶ The above is a notation. “ $A \mid B$ ” does not represent any set operation! (This is an abuse of notation!)
- ▶ Given a  $B$ , conditional probability is a new probability assignment to any event.
- ▶ That is, given  $B$  with  $P(B) > 0$ ,

# Conditional Probability

- ▶ Let  $B$  be an event with  $P(B) > 0$ . We define conditional probability, conditioned on  $B$ , of any event,  $A$ , as

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)} = \frac{P(AB)}{P(B)}$$

- ▶ The above is a notation. “ $A \mid B$ ” does not represent any set operation! (This is an abuse of notation!)
- ▶ Given a  $B$ , conditional probability is a new probability assignment to any event.
- ▶ That is, given  $B$  with  $P(B) > 0$ , we define a new probability  $P_B : \mathcal{F} \rightarrow [0, 1]$  by

# Conditional Probability

- ▶ Let  $B$  be an event with  $P(B) > 0$ . We define conditional probability, conditioned on  $B$ , of any event,  $A$ , as

$$P(A \mid B) = \frac{P(A \cap B)}{P(B)} = \frac{P(AB)}{P(B)}$$

- ▶ The above is a notation. “ $A \mid B$ ” does not represent any set operation! (This is an abuse of notation!)
- ▶ Given a  $B$ , conditional probability is a new probability assignment to any event.
- ▶ That is, given  $B$  with  $P(B) > 0$ , we define a new probability  $P_B : \mathcal{F} \rightarrow [0, 1]$  by

$$P_B(A) = \frac{P(AB)}{P(B)}$$

- ▶ Conditional probability is a probability.

- ▶ Conditional probability is a probability.  
What does this mean?

- ▶ Conditional probability is a probability.  
What does this mean?
- ▶ The new function we defined,  $P_B : \mathcal{F} \rightarrow [0, 1]$ ,  
$$P_B(A) = \frac{P(AB)}{P(B)},$$
satisfies the three axioms of probability.

- ▶ Conditional probability is a probability.  
What does this mean?
- ▶ The new function we defined,  $P_B : \mathcal{F} \rightarrow [0, 1]$ ,  
$$P_B(A) = \frac{P(AB)}{P(B)},$$
satisfies the three axioms of probability.
- ▶ Easy to see:  $P_B(A) \geq 0$



- ▶ Conditional probability is a probability.  
What does this mean?
- ▶ The new function we defined,  $P_B : \mathcal{F} \rightarrow [0, 1]$ ,  
$$P_B(A) = \frac{P(AB)}{P(B)},$$
satisfies the three axioms of probability.
- ▶ Easy to see:  $P_B(A) \geq 0$  and  $P_B(\Omega) = 1$ .

- ▶ Conditional probability is a probability.  
What does this mean?
- ▶ The new function we defined,  $P_B : \mathcal{F} \rightarrow [0, 1]$ ,  
$$P_B(A) = \frac{P(AB)}{P(B)},$$
satisfies the three axioms of probability.
- ▶ Easy to see:  $P_B(A) \geq 0$  and  $P_B(\Omega) = 1$ .
- ▶ If  $A_1, A_2$  are mutually exclusive then  $A_1B$  and  $A_2B$  are also mutually exclusive

- ▶ Conditional probability is a probability.  
What does this mean?
- ▶ The new function we defined,  $P_B : \mathcal{F} \rightarrow [0, 1]$ ,  
$$P_B(A) = \frac{P(AB)}{P(B)},$$
satisfies the three axioms of probability.
- ▶ Easy to see:  $P_B(A) \geq 0$  and  $P_B(\Omega) = 1$ .
- ▶ If  $A_1, A_2$  are mutually exclusive then  $A_1B$  and  $A_2B$  are also mutually exclusive and hence

$$P_B(A_1 + A_2) = \frac{P((A_1 + A_2)B)}{P(B)}$$

- ▶ Conditional probability is a probability.  
What does this mean?
- ▶ The new function we defined,  $P_B : \mathcal{F} \rightarrow [0, 1]$ ,  
 $P_B(A) = \frac{P(AB)}{P(B)}$ ,  
satisfies the three axioms of probability.
- ▶ Easy to see:  $P_B(A) \geq 0$  and  $P_B(\Omega) = 1$ .
- ▶ If  $A_1, A_2$  are mutually exclusive then  $A_1B$  and  $A_2B$  are also mutually exclusive and hence

$$P_B(A_1 + A_2) = \frac{P((A_1 + A_2)B)}{P(B)} = \frac{P(A_1B + A_2B)}{P(B)}$$

- ▶ Conditional probability is a probability.  
What does this mean?
- ▶ The new function we defined,  $P_B : \mathcal{F} \rightarrow [0, 1]$ ,  
 $P_B(A) = \frac{P(AB)}{P(B)}$ ,  
satisfies the three axioms of probability.
- ▶ Easy to see:  $P_B(A) \geq 0$  and  $P_B(\Omega) = 1$ .
- ▶ If  $A_1, A_2$  are mutually exclusive then  $A_1B$  and  $A_2B$  are also mutually exclusive and hence

$$\begin{aligned} P_B(A_1 + A_2) &= \frac{P((A_1 + A_2)B)}{P(B)} = \frac{P(A_1B + A_2B)}{P(B)} \\ &= \frac{P(A_1B) + P(A_2B)}{P(B)} = P_B(A_1) + P_B(A_2) \end{aligned}$$

- ▶ Conditional probability is a probability.  
What does this mean?
- ▶ The new function we defined,  $P_B : \mathcal{F} \rightarrow [0, 1]$ ,  
 $P_B(A) = \frac{P(AB)}{P(B)}$ ,  
satisfies the three axioms of probability.
- ▶ Easy to see:  $P_B(A) \geq 0$  and  $P_B(\Omega) = 1$ .
- ▶ If  $A_1, A_2$  are mutually exclusive then  $A_1B$  and  $A_2B$  are also mutually exclusive and hence

$$\begin{aligned} P_B(A_1 + A_2) &= \frac{P((A_1 + A_2)B)}{P(B)} = \frac{P(A_1B + A_2B)}{P(B)} \\ &= \frac{P(A_1B) + P(A_2B)}{P(B)} = P_B(A_1) + P_B(A_2) \end{aligned}$$

- ▶ Once we understand conditional probability is a new probability assignment, we go back to the 'standard notation'

$$P(A \mid B) = \frac{P(AB)}{P(B)}$$

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- Note  $P(B|B) = 1$  and  $P(A|B) > 0$  only if  $P(AB) > 0$ .



$$P(A | B) = \frac{P(AB)}{P(B)}$$

- ▶ Note  $P(B|B) = 1$  and  $P(A|B) > 0$  only if  $P(AB) > 0$ .
- ▶ Now the 'new' probability of each event is determined by what it has in common with  $B$ .

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- ▶ Note  $P(B|B) = 1$  and  $P(A|B) > 0$  only if  $P(AB) > 0$ .
- ▶ Now the 'new' probability of each event is determined by what it has in common with  $B$ .
- ▶ If we know the event  $B$  has occurred, then based on this knowledge we can readjust probabilities of all events and that is given by the conditional probability.

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- ▶ Note  $P(B|B) = 1$  and  $P(A|B) > 0$  only if  $P(AB) > 0$ .
- ▶ Now the 'new' probability of each event is determined by what it has in common with  $B$ .
- ▶ If we know the event  $B$  has occurred, then based on this knowledge we can readjust probabilities of all events and that is given by the conditional probability.
- ▶ Intuitively it is as if the sample space is now reduced to  $B$  because we are given the information that  $B$  has occurred.

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- ▶ Note  $P(B|B) = 1$  and  $P(A|B) > 0$  only if  $P(AB) > 0$ .
- ▶ Now the 'new' probability of each event is determined by what it has in common with  $B$ .
- ▶ If we know the event  $B$  has occurred, then based on this knowledge we can readjust probabilities of all events and that is given by the conditional probability.
- ▶ Intuitively it is as if the sample space is now reduced to  $B$  because we are given the information that  $B$  has occurred.
- ▶ This is a useful intuition as long as we understand it properly.

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- ▶ Note  $P(B|B) = 1$  and  $P(A|B) > 0$  only if  $P(AB) > 0$ .
- ▶ Now the 'new' probability of each event is determined by what it has in common with  $B$ .
- ▶ If we know the event  $B$  has occurred, then based on this knowledge we can readjust probabilities of all events and that is given by the conditional probability.
- ▶ Intuitively it is as if the sample space is now reduced to  $B$  because we are given the information that  $B$  has occurred.
- ▶ This is a useful intuition as long as we understand it properly.
- ▶ It is not as if we talk about conditional probability only for subsets of  $B$ . Conditional probability is also with respect to the original probability space. Every element of  $\mathcal{F}$  has conditional probability defined.

$$P(A \mid B) = \frac{P(AB)}{P(B)}$$

$$P(A \mid B) = \frac{P(AB)}{P(B)}$$

► Suppose  $P(A \mid B) > P(A)$

$$P(A \mid B) = \frac{P(AB)}{P(B)}$$

- ▶ Suppose  $P(A \mid B) > P(A)$   
Does it mean “B **causes** A”?



$$P(A | B) = \frac{P(AB)}{P(B)}$$

- Suppose  $P(A | B) > P(A)$   
Does it mean “B **causes** A”?

$$P(A | B) > P(A) \Rightarrow P(AB) > P(A)P(B)$$

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- Suppose  $P(A | B) > P(A)$   
Does it mean “B **causes** A”?

$$\begin{aligned} P(A | B) > P(A) &\Rightarrow P(AB) > P(A)P(B) \\ &\Rightarrow \frac{P(AB)}{P(A)} > P(B) \end{aligned}$$

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- Suppose  $P(A | B) > P(A)$   
Does it mean “B **causes** A”?

$$\begin{aligned} P(A | B) > P(A) &\Rightarrow P(AB) > P(A)P(B) \\ &\Rightarrow \frac{P(AB)}{P(A)} > P(B) \\ &\Rightarrow P(B | A) > P(B) \end{aligned}$$

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- Suppose  $P(A | B) > P(A)$   
Does it mean “B **causes** A”?

$$\begin{aligned} P(A | B) > P(A) &\Rightarrow P(AB) > P(A)P(B) \\ &\Rightarrow \frac{P(AB)}{P(A)} > P(B) \\ &\Rightarrow P(B | A) > P(B) \end{aligned}$$

- Hence, conditional probabilities cannot actually capture causal influences.

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- ▶ Suppose  $P(A | B) > P(A)$   
Does it mean “B **causes** A”?

$$\begin{aligned}P(A | B) > P(A) &\Rightarrow P(AB) > P(A)P(B) \\&\Rightarrow \frac{P(AB)}{P(A)} > P(B) \\&\Rightarrow P(B | A) > P(B)\end{aligned}$$

- ▶ Hence, conditional probabilities cannot actually capture causal influences.
- ▶ There are probabilistic methods to capture causation (but far beyond the scope of this course!)

- ▶ In a conditional probability, the conditioning event can be any event (with positive probability)

- ▶ In a conditional probability, the conditioning event can be any event (with positive probability)
- ▶ In particular, it could be intersection of events.

- ▶ In a conditional probability, the conditioning event can be any event (with positive probability)
- ▶ In particular, it could be intersection of events.
- ▶ We think of that as conditioning on multiple events.



- ▶ In a conditional probability, the conditioning event can be any event (with positive probability)
- ▶ In particular, it could be intersection of events.
- ▶ We think of that as conditioning on multiple events.

$$P(A \mid B, C) = P(A \mid BC)$$

- ▶ In a conditional probability, the conditioning event can be any event (with positive probability)
- ▶ In particular, it could be intersection of events.
- ▶ We think of that as conditioning on multiple events.

$$P(A \mid B, C) = P(A \mid BC) = \frac{P(ABC)}{P(BC)}$$

- The conditional probability is defined by

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- ▶ The conditional probability is defined by

$$P(A \mid B) = \frac{P(AB)}{P(B)}$$

- ▶ This gives us a useful identity

$$P(AB) = P(A \mid B)P(B)$$

- ▶ The conditional probability is defined by

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- ▶ This gives us a useful identity

$$P(AB) = P(A | B)P(B)$$

- ▶ We can iterate this for multiple events

- ▶ The conditional probability is defined by

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- ▶ This gives us a useful identity

$$P(AB) = P(A | B)P(B)$$

- ▶ We can iterate this for multiple events

$$P(ABC) = P(A | BC)P(BC)$$

- ▶ The conditional probability is defined by

$$P(A | B) = \frac{P(AB)}{P(B)}$$

- ▶ This gives us a useful identity

$$P(AB) = P(A | B)P(B)$$

- ▶ We can iterate this for multiple events

$$P(ABC) = P(A | BC)P(BC) = P(A | BC)P(B | C)P(C)$$

- Let  $B_1, \dots, B_m$  be events such that  $\cup_{i=1}^m B_i = \Omega$  and  $B_i B_j = \phi, \forall i \neq j$ .



- ▶ Let  $B_1, \dots, B_m$  be events such that  $\cup_{i=1}^m B_i = \Omega$  and  $B_i B_j = \phi, \forall i \neq j$ .
- ▶ Such a collection of events is said to be a partition of  $\Omega$ .

- ▶ Let  $B_1, \dots, B_m$  be events such that  $\cup_{i=1}^m B_i = \Omega$  and  $B_i B_j = \phi, \forall i \neq j$ .
- ▶ Such a collection of events is said to be a partition of  $\Omega$ . (They are also sometimes said to be mutually exclusive and collectively exhaustive).

- ▶ Let  $B_1, \dots, B_m$  be events such that  $\cup_{i=1}^m B_i = \Omega$  and  $B_i B_j = \phi, \forall i \neq j$ .
- ▶ Such a collection of events is said to be a partition of  $\Omega$ . (They are also sometimes said to be mutually exclusive and collectively exhaustive).
- ▶ Given this partition, any other event can be represented as a mutually exclusive union as

$$A = AB_1 + \dots + AB_m$$

- ▶ Let  $B_1, \dots, B_m$  be events such that  $\cup_{i=1}^m B_i = \Omega$  and  $B_i B_j = \phi, \forall i \neq j$ .
- ▶ Such a collection of events is said to be a partition of  $\Omega$ . (They are also sometimes said to be mutually exclusive and collectively exhaustive).
- ▶ Given this partition, any other event can be represented as a mutually exclusive union as

$$A = AB_1 + \dots + AB_m$$

To explain the notation again

- ▶ Let  $B_1, \dots, B_m$  be events such that  $\cup_{i=1}^m B_i = \Omega$  and  $B_i B_j = \phi, \forall i \neq j$ .
- ▶ Such a collection of events is said to be a partition of  $\Omega$ . (They are also sometimes said to be mutually exclusive and collectively exhaustive).
- ▶ Given this partition, any other event can be represented as a mutually exclusive union as

$$A = AB_1 + \dots + AB_m$$

To explain the notation again

$$A = A \cap \Omega$$

- ▶ Let  $B_1, \dots, B_m$  be events such that  $\cup_{i=1}^m B_i = \Omega$  and  $B_i B_j = \phi, \forall i \neq j$ .
- ▶ Such a collection of events is said to be a partition of  $\Omega$ . (They are also sometimes said to be mutually exclusive and collectively exhaustive).
- ▶ Given this partition, any other event can be represented as a mutually exclusive union as

$$A = AB_1 + \dots + AB_m$$

To explain the notation again

$$A = A \cap \Omega = A \cap (B_1 \cup \dots \cup B_m)$$

- ▶ Let  $B_1, \dots, B_m$  be events such that  $\cup_{i=1}^m B_i = \Omega$  and  $B_i B_j = \phi, \forall i \neq j$ .
- ▶ Such a collection of events is said to be a partition of  $\Omega$ . (They are also sometimes said to be mutually exclusive and collectively exhaustive).
- ▶ Given this partition, any other event can be represented as a mutually exclusive union as

$$A = AB_1 + \dots + AB_m$$

To explain the notation again

$$A = A \cap \Omega = A \cap (B_1 \cup \dots \cup B_m) = (A \cap B_1) \cup \dots \cup (A \cap B_m)$$

- ▶ Let  $B_1, \dots, B_m$  be events such that  $\cup_{i=1}^m B_i = \Omega$  and  $B_i B_j = \phi, \forall i \neq j$ .
- ▶ Such a collection of events is said to be a partition of  $\Omega$ . (They are also sometimes said to be mutually exclusive and collectively exhaustive).
- ▶ Given this partition, any other event can be represented as a mutually exclusive union as

$$A = AB_1 + \dots + AB_m$$

To explain the notation again

$$A = A \cap \Omega = A \cap (B_1 \cup \dots \cup B_m) = (A \cap B_1) \cup \dots \cup (A \cap B_m)$$

$$\text{Hence, } A = AB_1 + \dots + AB_m$$



- ▶ Let  $B_1, \dots, B_m$  be events such that  $\cup_{i=1}^m B_i = \Omega$  and  $B_i B_j = \phi, \forall i \neq j$ .
- ▶ Such a collection of events is said to be a partition of  $\Omega$ . (They are also sometimes said to be mutually exclusive and collectively exhaustive).
- ▶ Given this partition, any other event can be represented as a mutually exclusive union as

$$A = AB_1 + \dots + AB_m$$

To explain the notation again

$$A = A \cap \Omega = A \cap (B_1 \cup \dots \cup B_m) = (A \cap B_1) \cup \dots \cup (A \cap B_m)$$

Hence,  $A = AB_1 + \dots + AB_m$  when  $B_1, \dots, B_m$  be a partition of  $\Omega$ .

# Total Probability rule

- ▶ Let  $B_1, \dots, B_m$  be a partition of  $\Omega$ .

# Total Probability rule

- ▶ Let  $B_1, \dots, B_m$  be a partition of  $\Omega$ .
- ▶ Then, for any event  $A$ , we have

$$P(A) = P(AB_1 + \dots + AB_m)$$

# Total Probability rule

- ▶ Let  $B_1, \dots, B_m$  be a partition of  $\Omega$ .
- ▶ Then, for any event  $A$ , we have

$$\begin{aligned} P(A) &= P(AB_1 + \dots + AB_m) \\ &= P(AB_1) + \dots + P(AB_m) \end{aligned}$$

# Total Probability rule

- ▶ Let  $B_1, \dots, B_m$  be a partition of  $\Omega$ .
- ▶ Then, for any event  $A$ , we have

$$\begin{aligned}P(A) &= P(AB_1 + \dots + AB_m) \\&= P(AB_1) + \dots + P(AB_m) \\&= P(A | B_1)P(B_1) + \dots + P(A | B_m)P(B_m)\end{aligned}$$

# Total Probability rule

- ▶ Let  $B_1, \dots, B_m$  be a partition of  $\Omega$ .
- ▶ Then, for any event  $A$ , we have

$$\begin{aligned}P(A) &= P(AB_1 + \dots + AB_m) \\&= P(AB_1) + \dots + P(AB_m) \\&= P(A | B_1)P(B_1) + \dots + P(A | B_m)P(B_m)\end{aligned}$$

- ▶ The formula (where  $B_i$  form a partition)

$$P(A) = \sum_i P(A | B_i)P(B_i)$$

is known as **total probability rule** or total probability law or total probability formula.

# Total Probability rule

- ▶ Let  $B_1, \dots, B_m$  be a partition of  $\Omega$ .
- ▶ Then, for any event  $A$ , we have

$$\begin{aligned}P(A) &= P(AB_1 + \dots + AB_m) \\&= P(AB_1) + \dots + P(AB_m) \\&= P(A | B_1)P(B_1) + \dots + P(A | B_m)P(B_m)\end{aligned}$$

- ▶ The formula (where  $B_i$  form a partition)

$$P(A) = \sum_i P(A | B_i)P(B_i)$$

is known as **total probability rule** or total probability law or total probability formula.

- ▶ This is a very useful in many situations. (“arguing by cases”)

## Example: Polya's Urn

An urn contains  $r$  red balls and  $b$  black balls. We draw a ball at random, note its color, and put back that ball along with  $c$  balls of the same color. We keep repeating this process. Let  $R_n$  ( $B_n$ ) denote the event of drawing a red (black) ball at the  $n^{th}$  draw. We want to calculate the probabilities of all these events.



## Example: Polya's Urn

An urn contains  $r$  red balls and  $b$  black balls. We draw a ball at random, note its color, and put back that ball along with  $c$  balls of the same color. We keep repeating this process. Let  $R_n$  ( $B_n$ ) denote the event of drawing a red (black) ball at the  $n^{th}$  draw. We want to calculate the probabilities of all these events.

► It is easy to see that  $P(R_1) = \frac{r}{r+b}$

## Example: Polya's Urn

An urn contains  $r$  red balls and  $b$  black balls. We draw a ball at random, note its color, and put back that ball along with  $c$  balls of the same color. We keep repeating this process. Let  $R_n$  ( $B_n$ ) denote the event of drawing a red (black) ball at the  $n^{th}$  draw. We want to calculate the probabilities of all these events.

► It is easy to see that  $P(R_1) = \frac{r}{r+b}$  and  $P(B_1) = \frac{b}{r+b}$ .

## Example: Polya's Urn

An urn contains  $r$  red balls and  $b$  black balls. We draw a ball at random, note its color, and put back that ball along with  $c$  balls of the same color. We keep repeating this process. Let  $R_n$  ( $B_n$ ) denote the event of drawing a red (black) ball at the  $n^{th}$  draw. We want to calculate the probabilities of all these events.

- ▶ It is easy to see that  $P(R_1) = \frac{r}{r+b}$  and  $P(B_1) = \frac{b}{r+b}$ .
- ▶ For  $R_2$  we have, using total probability rule,

## Example: Polya's Urn

An urn contains  $r$  red balls and  $b$  black balls. We draw a ball at random, note its color, and put back that ball along with  $c$  balls of the same color. We keep repeating this process. Let  $R_n$  ( $B_n$ ) denote the event of drawing a red (black) ball at the  $n^{th}$  draw. We want to calculate the probabilities of all these events.

- ▶ It is easy to see that  $P(R_1) = \frac{r}{r+b}$  and  $P(B_1) = \frac{b}{r+b}$ .
- ▶ For  $R_2$  we have, using total probability rule,

$$P(R_2) = P(R_2 \mid R_1)P(R_1) + P(R_2 \mid B_1)P(B_1)$$

## Example: Polya's Urn

An urn contains  $r$  red balls and  $b$  black balls. We draw a ball at random, note its color, and put back that ball along with  $c$  balls of the same color. We keep repeating this process. Let  $R_n$  ( $B_n$ ) denote the event of drawing a red (black) ball at the  $n^{th}$  draw. We want to calculate the probabilities of all these events.

- ▶ It is easy to see that  $P(R_1) = \frac{r}{r+b}$  and  $P(B_1) = \frac{b}{r+b}$ .
- ▶ For  $R_2$  we have, using total probability rule,

$$\begin{aligned} P(R_2) &= P(R_2 \mid R_1)P(R_1) + P(R_2 \mid B_1)P(B_1) \\ &= \frac{r+c}{r+c+b} \end{aligned}$$

## Example: Polya's Urn

An urn contains  $r$  red balls and  $b$  black balls. We draw a ball at random, note its color, and put back that ball along with  $c$  balls of the same color. We keep repeating this process. Let  $R_n$  ( $B_n$ ) denote the event of drawing a red (black) ball at the  $n^{th}$  draw. We want to calculate the probabilities of all these events.

- ▶ It is easy to see that  $P(R_1) = \frac{r}{r+b}$  and  $P(B_1) = \frac{b}{r+b}$ .
- ▶ For  $R_2$  we have, using total probability rule,

$$\begin{aligned} P(R_2) &= P(R_2 \mid R_1)P(R_1) + P(R_2 \mid B_1)P(B_1) \\ &= \frac{r+c}{r+c+b} \frac{r}{r+b} + \end{aligned}$$

## Example: Polya's Urn

An urn contains  $r$  red balls and  $b$  black balls. We draw a ball at random, note its color, and put back that ball along with  $c$  balls of the same color. We keep repeating this process. Let  $R_n$  ( $B_n$ ) denote the event of drawing a red (black) ball at the  $n^{th}$  draw. We want to calculate the probabilities of all these events.

- ▶ It is easy to see that  $P(R_1) = \frac{r}{r+b}$  and  $P(B_1) = \frac{b}{r+b}$ .
- ▶ For  $R_2$  we have, using total probability rule,

$$\begin{aligned} P(R_2) &= P(R_2 \mid R_1)P(R_1) + P(R_2 \mid B_1)P(B_1) \\ &= \frac{r+c}{r+c+b} \frac{r}{r+b} + \frac{r}{r+b+c} \frac{b}{r+b} \end{aligned}$$

## Example: Polya's Urn

An urn contains  $r$  red balls and  $b$  black balls. We draw a ball at random, note its color, and put back that ball along with  $c$  balls of the same color. We keep repeating this process. Let  $R_n$  ( $B_n$ ) denote the event of drawing a red (black) ball at the  $n^{th}$  draw. We want to calculate the probabilities of all these events.

- ▶ It is easy to see that  $P(R_1) = \frac{r}{r+b}$  and  $P(B_1) = \frac{b}{r+b}$ .
- ▶ For  $R_2$  we have, using total probability rule,

$$\begin{aligned}P(R_2) &= P(R_2 \mid R_1)P(R_1) + P(R_2 \mid B_1)P(B_1) \\&= \frac{r+c}{r+c+b} \frac{r}{r+b} + \frac{r}{r+b+c} \frac{b}{r+b} \\&= \frac{r(r+c+b)}{(r+c+b)(r+b)}\end{aligned}$$



## Example: Polya's Urn

An urn contains  $r$  red balls and  $b$  black balls. We draw a ball at random, note its color, and put back that ball along with  $c$  balls of the same color. We keep repeating this process. Let  $R_n$  ( $B_n$ ) denote the event of drawing a red (black) ball at the  $n^{th}$  draw. We want to calculate the probabilities of all these events.

- ▶ It is easy to see that  $P(R_1) = \frac{r}{r+b}$  and  $P(B_1) = \frac{b}{r+b}$ .
- ▶ For  $R_2$  we have, using total probability rule,

$$\begin{aligned}P(R_2) &= P(R_2 \mid R_1)P(R_1) + P(R_2 \mid B_1)P(B_1) \\&= \frac{r+c}{r+c+b} \frac{r}{r+b} + \frac{r}{r+b+c} \frac{b}{r+b} \\&= \frac{r(r+c+b)}{(r+c+b)(r+b)} = \frac{r}{r+b}\end{aligned}$$

## Example: Polya's Urn

An urn contains  $r$  red balls and  $b$  black balls. We draw a ball at random, note its color, and put back that ball along with  $c$  balls of the same color. We keep repeating this process. Let  $R_n$  ( $B_n$ ) denote the event of drawing a red (black) ball at the  $n^{\text{th}}$  draw. We want to calculate the probabilities of all these events.

- ▶ It is easy to see that  $P(R_1) = \frac{r}{r+b}$  and  $P(B_1) = \frac{b}{r+b}$ .
- ▶ For  $R_2$  we have, using total probability rule,

$$\begin{aligned}P(R_2) &= P(R_2 \mid R_1)P(R_1) + P(R_2 \mid B_1)P(B_1) \\&= \frac{r+c}{r+c+b} \frac{r}{r+b} + \frac{r}{r+b+c} \frac{b}{r+b} \\&= \frac{r(r+c+b)}{(r+c+b)(r+b)} = \frac{r}{r+b} = P(R_1)\end{aligned}$$

- ▶ Similarly we can show that  $P(B_2) = P(B_1)$ .

- ▶ Similarly we can show that  $P(B_2) = P(B_1)$ .
- ▶ One can show by mathematical induction that  $P(R_n) = P(R_1)$  and  $P(B_n) = P(B_1)$  for all  $n$ .  
(Left as an exercise for you!)

- ▶ Similarly we can show that  $P(B_2) = P(B_1)$ .
- ▶ One can show by mathematical induction that  $P(R_n) = P(R_1)$  and  $P(B_n) = P(B_1)$  for all  $n$ .  
(Left as an exercise for you!)
- ▶ This does not depend on the value of  $c$ !

# Bayes Rule

- ▶ Another important formula based on conditional probability is Bayes Rule:

# Bayes Rule

- ▶ Another important formula based on conditional probability is Bayes Rule:

$$P(A | B) = \frac{P(AB)}{P(B)} = \frac{P(B | A)P(A)}{P(B)}$$

# Bayes Rule

- ▶ Another important formula based on conditional probability is Bayes Rule:

$$P(A | B) = \frac{P(AB)}{P(B)} = \frac{P(B | A)P(A)}{P(B)}$$

- ▶ This allows one to calculate  $P(A | B)$  if we know  $P(B | A)$ .



# Bayes Rule

- ▶ Another important formula based on conditional probability is Bayes Rule:

$$P(A | B) = \frac{P(AB)}{P(B)} = \frac{P(B | A)P(A)}{P(B)}$$

- ▶ This allows one to calculate  $P(A | B)$  if we know  $P(B | A)$ .
- ▶ Useful in many applications because one conditional probability may be more easier to obtain (or estimate) than the other.

# Bayes Rule

- ▶ Another important formula based on conditional probability is Bayes Rule:

$$P(A | B) = \frac{P(AB)}{P(B)} = \frac{P(B | A)P(A)}{P(B)}$$

- ▶ This allows one to calculate  $P(A | B)$  if we know  $P(B | A)$ .
- ▶ Useful in many applications because one conditional probability may be more easier to obtain (or estimate) than the other.
- ▶ Often one uses total probability rule to calculate the denominator in the RHS above:

# Bayes Rule

- ▶ Another important formula based on conditional probability is Bayes Rule:

$$P(A | B) = \frac{P(AB)}{P(B)} = \frac{P(B | A)P(A)}{P(B)}$$

- ▶ This allows one to calculate  $P(A | B)$  if we know  $P(B | A)$ .
- ▶ Useful in many applications because one conditional probability may be more easier to obtain (or estimate) than the other.
- ▶ Often one uses total probability rule to calculate the denominator in the RHS above:

$$P(A | B) = \frac{P(B | A)P(A)}{P(B | A)P(A) + P(B | A^c)P(A^c)}$$

## Example: Bayes Rule

Let  $D$  and  $D^c$  denote someone being diagnosed as having a disease or not having it. Let  $T_+$  and  $T_-$  denote the events of a test for it being positive or negative. (Note that  $T_+^c = T_-$ ). We want to calculate  $P(D|T_+)$ .

## Example: Bayes Rule

Let  $D$  and  $D^c$  denote someone being diagnosed as having a disease or not having it. Let  $T_+$  and  $T_-$  denote the events of a test for it being positive or negative. (Note that  $T_+^c = T_-$ ). We want to calculate  $P(D|T_+)$ .

► We have, by Bayes rule,

$$P(D|T_+) = \frac{P(T_+|D)P(D)}{P(T_+|D)P(D) + P(T_+|D^c)P(D^c)}$$

## Example: Bayes Rule

Let  $D$  and  $D^c$  denote someone being diagnosed as having a disease or not having it. Let  $T_+$  and  $T_-$  denote the events of a test for it being positive or negative. (Note that  $T_+^c = T_-$ ). We want to calculate  $P(D|T_+)$ .

- ▶ We have, by Bayes rule,

$$P(D|T_+) = \frac{P(T_+|D)P(D)}{P(T_+|D)P(D) + P(T_+|D^c)P(D^c)}$$

- ▶ The probabilities  $P(T_+|D)$  and  $P(T_+|D^c)$  can be obtained through, for example, laboratory experiments.

## Example: Bayes Rule

Let  $D$  and  $D^c$  denote someone being diagnosed as having a disease or not having it. Let  $T_+$  and  $T_-$  denote the events of a test for it being positive or negative. (Note that  $T_+^c = T_-$ ). We want to calculate  $P(D|T_+)$ .

- ▶ We have, by Bayes rule,

$$P(D|T_+) = \frac{P(T_+|D)P(D)}{P(T_+|D)P(D) + P(T_+|D^c)P(D^c)}$$

- ▶ The probabilities  $P(T_+|D)$  and  $P(T_+|D^c)$  can be obtained through, for example, laboratory experiments.
- ▶  $P(T_+|D)$  is called the true positive rate and  $P(T_+|D^c)$  is called false positive rate.

## Example: Bayes Rule

Let  $D$  and  $D^c$  denote someone being diagnosed as having a disease or not having it. Let  $T_+$  and  $T_-$  denote the events of a test for it being positive or negative. (Note that  $T_+^c = T_-$ ). We want to calculate  $P(D|T_+)$ .

- ▶ We have, by Bayes rule,

$$P(D|T_+) = \frac{P(T_+|D)P(D)}{P(T_+|D)P(D) + P(T_+|D^c)P(D^c)}$$

- ▶ The probabilities  $P(T_+|D)$  and  $P(T_+|D^c)$  can be obtained through, for example, laboratory experiments.
- ▶  $P(T_+|D)$  is called the true positive rate and  $P(T_+|D^c)$  is called false positive rate.
- ▶ We also need  $P(D)$ , the probability of a random person having the disease.



- ▶ Let us take some specific numbers

- ▶ Let us take some specific numbers
- ▶ Let:  $P(D) = 0.5$ ,  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .

- ▶ Let us take some specific numbers
- ▶ Let:  $P(D) = 0.5$ ,  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .

$$P(D|T_+) = \frac{0.99 * 0.5}{0.99 * 0.5 + 0.05 * 0.5}$$

- ▶ Let us take some specific numbers
- ▶ Let:  $P(D) = 0.5$ ,  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .

$$P(D|T_+) = \frac{0.99 * 0.5}{0.99 * 0.5 + 0.05 * 0.5} = 0.95$$

- ▶ Let us take some specific numbers
- ▶ Let:  $P(D) = 0.5$ ,  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .

$$P(D|T_+) = \frac{0.99 * 0.5}{0.99 * 0.5 + 0.05 * 0.5} = 0.95$$

That is pretty good.

- ▶ Let us take some specific numbers
- ▶ Let:  $P(D) = 0.5$ ,  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .

$$P(D|T_+) = \frac{0.99 * 0.5}{0.99 * 0.5 + 0.05 * 0.5} = 0.95$$

That is pretty good.

- ▶ But taking  $P(D) = 0.5$  is not realistic. Let us take  $P(D) = 0.1$ .

- ▶ Let us take some specific numbers
- ▶ Let:  $P(D) = 0.5$ ,  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .

$$P(D|T_+) = \frac{0.99 * 0.5}{0.99 * 0.5 + 0.05 * 0.5} = 0.95$$

That is pretty good.

- ▶ But taking  $P(D) = 0.5$  is not realistic. Let us take  $P(D) = 0.1$ .

$$P(D|T_+) = \frac{0.99 * 0.1}{0.99 * 0.1 + 0.05 * 0.9}$$

- ▶ Let us take some specific numbers
- ▶ Let:  $P(D) = 0.5$ ,  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .

$$P(D|T_+) = \frac{0.99 * 0.5}{0.99 * 0.5 + 0.05 * 0.5} = 0.95$$

That is pretty good.

- ▶ But taking  $P(D) = 0.5$  is not realistic. Let us take  $P(D) = 0.1$ .

$$P(D|T_+) = \frac{0.99 * 0.1}{0.99 * 0.1 + 0.05 * 0.9} = 0.69$$



- ▶ Let us take some specific numbers
- ▶ Let:  $P(D) = 0.5$ ,  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .

$$P(D|T_+) = \frac{0.99 * 0.5}{0.99 * 0.5 + 0.05 * 0.5} = 0.95$$

That is pretty good.

- ▶ But taking  $P(D) = 0.5$  is not realistic. Let us take  $P(D) = 0.1$ .

$$P(D|T_+) = \frac{0.99 * 0.1}{0.99 * 0.1 + 0.05 * 0.9} = 0.69$$

- ▶ Now suppose we can improve the test so that  $P(T_+|D^c) = 0.01$

- ▶ Let us take some specific numbers
- ▶ Let:  $P(D) = 0.5$ ,  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .

$$P(D|T_+) = \frac{0.99 * 0.5}{0.99 * 0.5 + 0.05 * 0.5} = 0.95$$

That is pretty good.

- ▶ But taking  $P(D) = 0.5$  is not realistic. Let us take  $P(D) = 0.1$ .

$$P(D|T_+) = \frac{0.99 * 0.1}{0.99 * 0.1 + 0.05 * 0.9} = 0.69$$

- ▶ Now suppose we can improve the test so that  $P(T_+|D^c) = 0.01$

$$P(D|T_+) = \frac{0.99 * 0.1}{0.99 * 0.1 + 0.01 * 0.9}$$

- ▶ Let us take some specific numbers
- ▶ Let:  $P(D) = 0.5$ ,  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .

$$P(D|T_+) = \frac{0.99 * 0.5}{0.99 * 0.5 + 0.05 * 0.5} = 0.95$$

That is pretty good.

- ▶ But taking  $P(D) = 0.5$  is not realistic. Let us take  $P(D) = 0.1$ .

$$P(D|T_+) = \frac{0.99 * 0.1}{0.99 * 0.1 + 0.05 * 0.9} = 0.69$$

- ▶ Now suppose we can improve the test so that  $P(T_+|D^c) = 0.01$

$$P(D|T_+) = \frac{0.99 * 0.1}{0.99 * 0.1 + 0.01 * 0.9} = 0.92$$

- ▶ Let us take some specific numbers
- ▶ Let:  $P(D) = 0.5$ ,  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .

$$P(D|T_+) = \frac{0.99 * 0.5}{0.99 * 0.5 + 0.05 * 0.5} = 0.95$$

That is pretty good.

- ▶ But taking  $P(D) = 0.5$  is not realistic. Let us take  $P(D) = 0.1$ .

$$P(D|T_+) = \frac{0.99 * 0.1}{0.99 * 0.1 + 0.05 * 0.9} = 0.69$$

- ▶ Now suppose we can improve the test so that  $P(T_+|D^c) = 0.01$

$$P(D|T_+) = \frac{0.99 * 0.1}{0.99 * 0.1 + 0.01 * 0.9} = 0.92$$

- ▶ These different cases are important in understanding the role of false positives rate.

- ▶  $P(D)$  is the probability that a random person has the disease. We call it the prior probability.

- ▶  $P(D)$  is the probability that a random person has the disease. We call it the prior probability.
- ▶  $P(D|T_+)$  is the probability of the random person having disease once we do a test and it came positive. We call it the posterior probability.

- ▶  $P(D)$  is the probability that a random person has the disease. We call it the prior probability.
- ▶  $P(D|T_+)$  is the probability of the random person having disease once we do a test and it came positive. We call it the posterior probability.
- ▶ Bayes rule essentially transforms the prior probability to posterior probability.

- ▶ In many applications of Bayes rule the same generic situation exists



- ▶ In many applications of Bayes rule the same generic situation exists
- ▶ Based on a measurement we want to predict (what may be called) the state of nature.

- ▶ In many applications of Bayes rule the same generic situation exists
- ▶ Based on a measurement we want to predict (what may be called) the state of nature.
- ▶ For another example, take a simple communication system.

- ▶ In many applications of Bayes rule the same generic situation exists
- ▶ Based on a measurement we want to predict (what may be called) the state of nature.
- ▶ For another example, take a simple communication system.
  - ▶  $D$  can represent the event that the transmitter sent bit 1.

- ▶ In many applications of Bayes rule the same generic situation exists
- ▶ Based on a measurement we want to predict (what may be called) the state of nature.
- ▶ For another example, take a simple communication system.
  - ▶  $D$  can represent the event that the transmitter sent bit 1.
  - ▶  $T_+$  can represent an event about the measurement we made at the receiver.

- ▶ In many applications of Bayes rule the same generic situation exists
- ▶ Based on a measurement we want to predict (what may be called) the state of nature.
- ▶ For another example, take a simple communication system.
  - ▶  $D$  can represent the event that the transmitter sent bit 1.
  - ▶  $T_+$  can represent an event about the measurement we made at the receiver.
  - ▶ We want the probability that bit 1 is sent based on the measurement.

- ▶ In many applications of Bayes rule the same generic situation exists
- ▶ Based on a measurement we want to predict (what may be called) the state of nature.
- ▶ For another example, take a simple communication system.
  - ▶  $D$  can represent the event that the transmitter sent bit 1.
  - ▶  $T_+$  can represent an event about the measurement we made at the receiver.
  - ▶ We want the probability that bit 1 is sent based on the measurement.
  - ▶ The knowledge we need is  $P(T_+|D)$ ,  $P(T_+|D^c)$  which can be determined through experiment or modelling of channel.



- ▶ Not all applications of Bayes rule involve a 'binary' situation



- ▶ Not all applications of Bayes rule involve a 'binary' situation
- ▶ Suppose  $D_1, D_2, D_3$  are the (exclusive) possibilities and  $T$  is an event about a measurement.

- ▶ Not all applications of Bayes rule involve a 'binary' situation
- ▶ Suppose  $D_1, D_2, D_3$  are the (exclusive) possibilities and  $T$  is an event about a measurement.

$$P(D_1|T) = \frac{P(T|D_1)P(D_1)}{P(T)}$$

- ▶ Not all applications of Bayes rule involve a ‘binary’ situation
- ▶ Suppose  $D_1, D_2, D_3$  are the (exclusive) possibilities and  $T$  is an event about a measurement.

$$\begin{aligned} P(D_1|T) &= \frac{P(T|D_1)P(D_1)}{P(T)} \\ &= \frac{P(T|D_1)P(D_1)}{P(T|D_1)P(D_1) + P(T|D_2)P(D_2) + P(T|D_3)P(D_3)} \end{aligned}$$

- ▶ Not all applications of Bayes rule involve a 'binary' situation
- ▶ Suppose  $D_1, D_2, D_3$  are the (exclusive) possibilities and  $T$  is an event about a measurement.

$$\begin{aligned} P(D_1|T) &= \frac{P(T|D_1)P(D_1)}{P(T)} \\ &= \frac{P(T|D_1)P(D_1)}{P(T|D_1)P(D_1) + P(T|D_2)P(D_2) + P(T|D_3)P(D_3)} \\ &= \frac{P(T|D_1)P(D_1)}{\sum_i P(T|D_i)P(D_i)} \end{aligned}$$

- ▶ Not all applications of Bayes rule involve a 'binary' situation
- ▶ Suppose  $D_1, D_2, D_3$  are the (exclusive) possibilities and  $T$  is an event about a measurement.

$$\begin{aligned} P(D_1|T) &= \frac{P(T|D_1)P(D_1)}{P(T)} \\ &= \frac{P(T|D_1)P(D_1)}{P(T|D_1)P(D_1) + P(T|D_2)P(D_2) + P(T|D_3)P(D_3)} \\ &= \frac{P(T|D_1)P(D_1)}{\sum_i P(T|D_i)P(D_i)} \end{aligned}$$

- ▶ Example: I have three coins with probability of heads being 0.1, 0.5, 0.8. I choose one at random and toss it twice and see heads both times. What is the probability it is the fair coin?

$$P(D|T_+) = \frac{P(T_+|D)P(D)}{P(T_+|D)P(D) + P(T_+|D^c)P(D^c)}$$

$$P(D|T_+) = \frac{P(T_+|D)P(D)}{P(T_+|D)P(D) + P(T_+|D^c)P(D^c)}$$

- In the binary situation we can think of Bayes rule in a slightly modified form too.

$$P(D|T_+) = \frac{P(T_+|D)P(D)}{P(T_+|D)P(D) + P(T_+|D^c)P(D^c)}$$

- In the binary situation we can think of Bayes rule in a slightly modified form too.

$$\frac{P(D|T_+)}{P(D^c|T_+)} = \frac{P(T_+|D)}{P(T_+|D^c)} \frac{P(D)}{P(D^c)}$$



$$P(D|T_+) = \frac{P(T_+|D)P(D)}{P(T_+|D)P(D) + P(T_+|D^c)P(D^c)}$$

- In the binary situation we can think of Bayes rule in a slightly modified form too.

$$\frac{P(D|T_+)}{P(D^c|T_+)} = \frac{P(T_+|D)}{P(T_+|D^c)} \frac{P(D)}{P(D^c)}$$

- This is called the odds-likelihood form of Bayes rule

$$P(D|T_+) = \frac{P(T_+|D)P(D)}{P(T_+|D)P(D) + P(T_+|D^c)P(D^c)}$$

- In the binary situation we can think of Bayes rule in a slightly modified form too.

$$\frac{P(D|T_+)}{P(D^c|T_+)} = \frac{P(T_+|D)}{P(T_+|D^c)} \frac{P(D)}{P(D^c)}$$

- This is called the odds-likelihood form of Bayes rule  
(The ratio of  $P(A)$  to  $P(A^c)$  is called odds for  $A$ )

# Independent Events

# Independent Events

- ▶ Two events  $A, B$  are said to be independent if

$$P(AB) = P(A)P(B)$$

# Independent Events

- ▶ Two events  $A, B$  are said to be independent if

$$P(AB) = P(A)P(B)$$

- ▶ Note that this is a definition. Two events are independent if and only if they satisfy the above.

# Independent Events

- ▶ Two events  $A, B$  are said to be independent if

$$P(AB) = P(A)P(B)$$

- ▶ Note that this is a definition. Two events are independent if and only if they satisfy the above.
- ▶ Suppose  $P(A), P(B) > 0$ . Then, if they are independent

# Independent Events

- ▶ Two events  $A, B$  are said to be independent if

$$P(AB) = P(A)P(B)$$

- ▶ Note that this is a definition. Two events are independent if and only if they satisfy the above.
- ▶ Suppose  $P(A), P(B) > 0$ . Then, if they are independent

$$P(A|B) = \frac{P(AB)}{P(B)} = P(A);$$

# Independent Events

- ▶ Two events  $A, B$  are said to be independent if

$$P(AB) = P(A)P(B)$$

- ▶ Note that this is a definition. Two events are independent if and only if they satisfy the above.
- ▶ Suppose  $P(A), P(B) > 0$ . Then, if they are independent

$$P(A|B) = \frac{P(AB)}{P(B)} = P(A); \text{ similarly } P(B|A) = P(B)$$



# Independent Events

- ▶ Two events  $A, B$  are said to be independent if

$$P(AB) = P(A)P(B)$$

- ▶ Note that this is a definition. Two events are independent if and only if they satisfy the above.
- ▶ Suppose  $P(A), P(B) > 0$ . Then, if they are independent

$$P(A|B) = \frac{P(AB)}{P(B)} = P(A); \text{ similarly } P(B|A) = P(B)$$

- ▶ This gives an intuitive feel for independence.

# Independent Events

- ▶ Two events  $A, B$  are said to be independent if

$$P(AB) = P(A)P(B)$$

- ▶ Note that this is a definition. Two events are independent if and only if they satisfy the above.
- ▶ Suppose  $P(A), P(B) > 0$ . Then, if they are independent

$$P(A|B) = \frac{P(AB)}{P(B)} = P(A); \text{ similarly } P(B|A) = P(B)$$

- ▶ This gives an intuitive feel for independence.
- ▶ Independence is an important (often confusing!) concept.

# Example: Independence

## Example: Independence

A class has 20 female and 30 male course (MTech) students and 6 female and 9 male research (PhD) students. Are gender and degree independent?

## Example: Independence

A class has 20 female and 30 male course (MTech) students and 6 female and 9 male research (PhD) students. Are gender and degree independent?

- ▶ Let  $F, M, C, R$  denote events of female, male, course, research students

## Example: Independence

A class has 20 female and 30 male course (MTech) students and 6 female and 9 male research (PhD) students. Are gender and degree independent?

- ▶ Let  $F, M, C, R$  denote events of female, male, course, research students
- ▶ From the given numbers, we can easily calculate the following:

## Example: Independence

A class has 20 female and 30 male course (MTech) students and 6 female and 9 male research (PhD) students. Are gender and degree independent?

- ▶ Let  $F, M, C, R$  denote events of female, male, course, research students
- ▶ From the given numbers, we can easily calculate the following:

$$P(F) = \frac{26}{65} = \frac{2}{5};$$

## Example: Independence

A class has 20 female and 30 male course (MTech) students and 6 female and 9 male research (PhD) students. Are gender and degree independent?

- ▶ Let  $F, M, C, R$  denote events of female, male, course, research students
- ▶ From the given numbers, we can easily calculate the following:

$$P(F) = \frac{26}{65} = \frac{2}{5}; \quad P(C) = \frac{50}{65} = \frac{10}{13};$$



## Example: Independence

A class has 20 female and 30 male course (MTech) students and 6 female and 9 male research (PhD) students. Are gender and degree independent?

- ▶ Let  $F, M, C, R$  denote events of female, male, course, research students
- ▶ From the given numbers, we can easily calculate the following:

$$P(F) = \frac{26}{65} = \frac{2}{5}; \quad P(C) = \frac{50}{65} = \frac{10}{13}; \quad P(FC) = \frac{20}{65} = \frac{4}{13}$$

## Example: Independence

A class has 20 female and 30 male course (MTech) students and 6 female and 9 male research (PhD) students. Are gender and degree independent?

- ▶ Let  $F, M, C, R$  denote events of female, male, course, research students
- ▶ From the given numbers, we can easily calculate the following:

$$P(F) = \frac{26}{65} = \frac{2}{5}; \quad P(C) = \frac{50}{65} = \frac{10}{13}; \quad P(FC) = \frac{20}{65} = \frac{4}{13}$$

- ▶ Hence we can verify

$$P(F)P(C) = \frac{2}{5} \frac{10}{13}$$

## Example: Independence

A class has 20 female and 30 male course (MTech) students and 6 female and 9 male research (PhD) students. Are gender and degree independent?

- ▶ Let  $F, M, C, R$  denote events of female, male, course, research students
- ▶ From the given numbers, we can easily calculate the following:

$$P(F) = \frac{26}{65} = \frac{2}{5}; \quad P(C) = \frac{50}{65} = \frac{10}{13}; \quad P(FC) = \frac{20}{65} = \frac{4}{13}$$

- ▶ Hence we can verify

$$P(F)P(C) = \frac{2}{5} \frac{10}{13} = \frac{4}{13}$$

## Example: Independence

A class has 20 female and 30 male course (MTech) students and 6 female and 9 male research (PhD) students. Are gender and degree independent?

- ▶ Let  $F, M, C, R$  denote events of female, male, course, research students
- ▶ From the given numbers, we can easily calculate the following:

$$P(F) = \frac{26}{65} = \frac{2}{5}; \quad P(C) = \frac{50}{65} = \frac{10}{13}; \quad P(FC) = \frac{20}{65} = \frac{4}{13}$$

- ▶ Hence we can verify

$$P(F)P(C) = \frac{2}{5} \frac{10}{13} = \frac{4}{13} = P(FC)$$

## Example: Independence

A class has 20 female and 30 male course (MTech) students and 6 female and 9 male research (PhD) students. Are gender and degree independent?

- ▶ Let  $F, M, C, R$  denote events of female, male, course, research students
- ▶ From the given numbers, we can easily calculate the following:

$$P(F) = \frac{26}{65} = \frac{2}{5}; \quad P(C) = \frac{50}{65} = \frac{10}{13}; \quad P(FC) = \frac{20}{65} = \frac{4}{13}$$

- ▶ Hence we can verify

$$P(F)P(C) = \frac{2}{5} \frac{10}{13} = \frac{4}{13} = P(FC)$$

and conclude that  $F$  and  $C$  are independent.

## Example: Independence

A class has 20 female and 30 male course (MTech) students and 6 female and 9 male research (PhD) students. Are gender and degree independent?

- ▶ Let  $F, M, C, R$  denote events of female, male, course, research students
- ▶ From the given numbers, we can easily calculate the following:

$$P(F) = \frac{26}{65} = \frac{2}{5}; \quad P(C) = \frac{50}{65} = \frac{10}{13}; \quad P(FC) = \frac{20}{65} = \frac{4}{13}$$

- ▶ Hence we can verify

$$P(F)P(C) = \frac{2}{5} \frac{10}{13} = \frac{4}{13} = P(FC)$$

and conclude that  $F$  and  $C$  are independent.  
Similarly we can show for others.

- ▶ In this example, if we keep all other numbers same but change the number of male research students to, say, 12 then the independence no longer holds.

- In this example, if we keep all other numbers same but change the number of male research students to, say, 12 then the independence no longer holds.

$$\left(\frac{26}{68} \frac{50}{68} \neq \frac{20}{68}\right)$$



- ▶ In this example, if we keep all other numbers same but change the number of male research students to, say, 12 then the independence no longer holds.

$$\left(\frac{26}{68} \frac{50}{68} \neq \frac{20}{68}\right)$$

- ▶ One needs to be careful about independence!

- ▶ In this example, if we keep all other numbers same but change the number of male research students to, say, 12 then the independence no longer holds.  
$$\left(\frac{26}{68} \frac{50}{68} \neq \frac{20}{68}\right)$$
- ▶ One needs to be careful about independence!
- ▶ We always have an underlying probability space  $(\Omega, \mathcal{F}, P)$

- ▶ In this example, if we keep all other numbers same but change the number of male research students to, say, 12 then the independence no longer holds.  
 $(\frac{26}{68} \frac{50}{68} \neq \frac{20}{68})$
- ▶ One needs to be careful about independence!
- ▶ We always have an underlying probability space  $(\Omega, \mathcal{F}, P)$
- ▶ Once that is given, the probabilities of all events are fixed.

- ▶ In this example, if we keep all other numbers same but change the number of male research students to, say, 12 then the independence no longer holds.

$$\left(\frac{26}{68} \frac{50}{68} \neq \frac{20}{68}\right)$$

- ▶ One needs to be careful about independence!
- ▶ We always have an underlying probability space  $(\Omega, \mathcal{F}, P)$
- ▶ Once that is given, the probabilities of all events are fixed.
- ▶ Hence whether or not two events are independent is a matter of 'calculation'

- ▶ If  $A$  and  $B$  are independent then so are  $A$  and  $B^c$ .

- ▶ If  $A$  and  $B$  are independent then so are  $A$  and  $B^c$ .
- ▶ Using  $A = AB + AB^c$ , and  $AB \subset A$ , we have

- ▶ If  $A$  and  $B$  are independent then so are  $A$  and  $B^c$ .
- ▶ Using  $A = AB + AB^c$ , and  $AB \subset A$ , we have

$$P(AB^c) = P(A - AB)$$

- ▶ If  $A$  and  $B$  are independent then so are  $A$  and  $B^c$ .
- ▶ Using  $A = AB + AB^c$ , and  $AB \subset A$ , we have

$$P(AB^c) = P(A - AB) = P(A) - P(AB)$$



- ▶ If  $A$  and  $B$  are independent then so are  $A$  and  $B^c$ .
- ▶ Using  $A = AB + AB^c$ , and  $AB \subset A$ , we have

$$P(AB^c) = P(A - AB) = P(A) - P(AB) = P(A)(1 - P(B))$$

- ▶ If  $A$  and  $B$  are independent then so are  $A$  and  $B^c$ .
- ▶ Using  $A = AB + AB^c$ , and  $AB \subset A$ , we have

$$P(AB^c) = P(A - AB) = P(A) - P(AB) = P(A)(1 - P(B)) = P(A)P(B^c)$$

- ▶ If  $A$  and  $B$  are independent then so are  $A$  and  $B^c$ .
- ▶ Using  $A = AB + AB^c$ , and  $AB \subset A$ , we have

$$P(AB^c) = P(A - AB) = P(A) - P(AB) = P(A)(1 - P(B)) = P(A)P(B^c)$$

- ▶ This also shows that  $A^c$  and  $B$  are independent and so are  $A^c$  and  $B^c$ .

- ▶ If  $A$  and  $B$  are independent then so are  $A$  and  $B^c$ .
- ▶ Using  $A = AB + AB^c$ , and  $AB \subset A$ , we have

$$P(AB^c) = P(A - AB) = P(A) - P(AB) = P(A)(1 - P(B)) = P(A)P(B^c)$$

- ▶ This also shows that  $A^c$  and  $B$  are independent and so are  $A^c$  and  $B^c$ .
- ▶ For example, in the previous problem, once we saw that  $F$  and  $C$  are independent, we can conclude  $M$  and  $C$  are also independent (because in this example we are taking  $F^c = M$ ).

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.
- ▶ That is,  $P(\{HH\}) = \frac{1}{4}$ ,  $P(\{HT\}) = \frac{1}{4}$  and so on



- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.
- ▶ That is,  $P(\{HH\}) = \frac{1}{4}$ ,  $P(\{HT\}) = \frac{1}{4}$  and so on
- ▶ Let  $A = \text{'H on 1st toss'}$

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.
- ▶ That is,  $P(\{HH\}) = \frac{1}{4}$ ,  $P(\{HT\}) = \frac{1}{4}$  and so on
- ▶ Let  $A = \text{'H on 1st toss'} = \{HH, HT\}$

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.
- ▶ That is,  $P(\{HH\}) = \frac{1}{4}$ ,  $P(\{HT\}) = \frac{1}{4}$  and so on
- ▶ Let  $A = \text{'H on 1st toss'} = \{HH, HT\}$  ( $P(A) = \frac{1}{2}$ )

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.
- ▶ That is,  $P(\{HH\}) = \frac{1}{4}$ ,  $P(\{HT\}) = \frac{1}{4}$  and so on
- ▶ Let  $A = \text{'H on 1st toss'} = \{HH, HT\}$  ( $P(A) = \frac{1}{2}$ )  
Let  $B = \text{'T on second toss'}$

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.
- ▶ That is,  $P(\{HH\}) = \frac{1}{4}$ ,  $P(\{HT\}) = \frac{1}{4}$  and so on
- ▶ Let  $A = \text{'H on 1st toss'} = \{HH, HT\}$  ( $P(A) = \frac{1}{2}$ )  
Let  $B = \text{'T on second toss'} = \{HT, TT\}$

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.
- ▶ That is,  $P(\{HH\}) = \frac{1}{4}$ ,  $P(\{HT\}) = \frac{1}{4}$  and so on
- ▶ Let  $A = \text{'H on 1st toss'} = \{HH, HT\}$  ( $P(A) = \frac{1}{2}$ )  
Let  $B = \text{'T on second toss'} = \{HT, TT\}$  ( $P(B) = \frac{1}{2}$ )

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.
- ▶ That is,  $P(\{HH\}) = \frac{1}{4}$ ,  $P(\{HT\}) = \frac{1}{4}$  and so on
- ▶ Let  $A = \text{'H on 1st toss'} = \{HH, HT\}$  ( $P(A) = \frac{1}{2}$ )  
Let  $B = \text{'T on second toss'} = \{HT, TT\}$  ( $P(B) = \frac{1}{2}$ )
- ▶ We have  $P(AB) = P(\{HT\}) = 0.25$

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.
- ▶ That is,  $P(\{HH\}) = \frac{1}{4}$ ,  $P(\{HT\}) = \frac{1}{4}$  and so on
- ▶ Let  $A = \text{'H on 1st toss'} = \{HH, HT\}$  ( $P(A) = \frac{1}{2}$ )  
Let  $B = \text{'T on second toss'} = \{HT, TT\}$  ( $P(B) = \frac{1}{2}$ )
- ▶ We have  $P(AB) = P(\{HT\}) = 0.25$
- ▶ Since  $P(A)P(B) = \frac{1}{2} \frac{1}{2}$



- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.
- ▶ That is,  $P(\{HH\}) = \frac{1}{4}$ ,  $P(\{HT\}) = \frac{1}{4}$  and so on
- ▶ Let  $A = \text{'H on 1st toss'} = \{HH, HT\}$  ( $P(A) = \frac{1}{2}$ )  
Let  $B = \text{'T on second toss'} = \{HT, TT\}$  ( $P(B) = \frac{1}{2}$ )
- ▶ We have  $P(AB) = P(\{HT\}) = 0.25$
- ▶ Since  $P(A)P(B) = \frac{1}{2} \frac{1}{2} = \frac{1}{4} = P(AB)$ ,

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.
- ▶ That is,  $P(\{HH\}) = \frac{1}{4}$ ,  $P(\{HT\}) = \frac{1}{4}$  and so on
- ▶ Let  $A = \text{'H on 1st toss'} = \{HH, HT\}$  ( $P(A) = \frac{1}{2}$ )  
Let  $B = \text{'T on second toss'} = \{HT, TT\}$  ( $P(B) = \frac{1}{2}$ )
- ▶ We have  $P(AB) = P(\{HT\}) = 0.25$
- ▶ Since  $P(A)P(B) = \frac{1}{2} \frac{1}{2} = \frac{1}{4} = P(AB)$ ,  
 $A, B$  are independent.

- ▶ Consider the random experiment of tossing two fair coins (or tossing a coin twice).
- ▶  $\Omega = \{HH, HT, TH, TT\}$ .  
Suppose we employ 'equally likely idea'.
- ▶ That is,  $P(\{HH\}) = \frac{1}{4}$ ,  $P(\{HT\}) = \frac{1}{4}$  and so on
- ▶ Let  $A = \text{'H on 1st toss'} = \{HH, HT\}$  ( $P(A) = \frac{1}{2}$ )  
Let  $B = \text{'T on second toss'} = \{HT, TT\}$  ( $P(B) = \frac{1}{2}$ )
- ▶ We have  $P(AB) = P(\{HT\}) = 0.25$
- ▶ Since  $P(A)P(B) = \frac{1}{2} \frac{1}{2} = \frac{1}{4} = P(AB)$ ,  
 $A, B$  are independent.
- ▶ Hence, in multiple tosses, assuming all outcomes are equally likely implies outcome of one toss is independent of another.

- ▶ In multiple tosses, assuming all outcomes are equally likely is alright if the coin is fair.

- ▶ In multiple tosses, assuming all outcomes are equally likely is alright if the coin is fair.
- ▶ Suppose we toss a biased coin two times.

- ▶ In multiple tosses, assuming all outcomes are equally likely is alright if the coin is fair.
- ▶ Suppose we toss a biased coin two times.
- ▶ Then the four outcomes are, obviously, not 'equally likely'

- ▶ In multiple tosses, assuming all outcomes are equally likely is alright if the coin is fair.
- ▶ Suppose we toss a biased coin two times.
- ▶ Then the four outcomes are, obviously, not 'equally likely'
- ▶ How should we then assign these probabilities?

- ▶ In multiple tosses, assuming all outcomes are equally likely is alright if the coin is fair.
- ▶ Suppose we toss a biased coin two times.
- ▶ Then the four outcomes are, obviously, not 'equally likely'
- ▶ How should we then assign these probabilities?
- ▶ If we assume tosses are independent then we can assign probabilities easily.



- Consider toss of a biased coin:

$$\Omega^1 = \{H, T\}, P(\{H\}) = p \text{ and } P(\{T\}) = 1 - p.$$

- ▶ Consider toss of a biased coin:  
 $\Omega^1 = \{H, T\}$ ,  $P(\{H\}) = p$  and  $P(\{T\}) = 1 - p$ .
- ▶ If we toss this twice then  $\Omega^2 = \{HH, HT, TH, TT\}$  and we assign

- Consider toss of a biased coin:

$$\Omega^1 = \{H, T\}, P(\{H\}) = p \text{ and } P(\{T\}) = 1 - p.$$

- If we toss this twice then  $\Omega^2 = \{HH, HT, TH, TT\}$  and we assign

$$P(\{HH\}) = p^2, P(\{HT\}) = p(1 - p),$$

- Consider toss of a biased coin:

$$\Omega^1 = \{H, T\}, P(\{H\}) = p \text{ and } P(\{T\}) = 1 - p.$$

- If we toss this twice then  $\Omega^2 = \{HH, HT, TH, TT\}$  and we assign

$$P(\{HH\}) = p^2, P(\{HT\}) = p(1 - p),$$

$$P(\{TH\}) = (1 - p)p, P(\{TT\}) = (1 - p)^2.$$

- ▶ Consider toss of a biased coin:  
 $\Omega^1 = \{H, T\}$ ,  $P(\{H\}) = p$  and  $P(\{T\}) = 1 - p$ .
- ▶ If we toss this twice then  $\Omega^2 = \{HH, HT, TH, TT\}$  and we assign  
 $P(\{HH\}) = p^2$ ,  $P(\{HT\}) = p(1 - p)$ ,  
 $P(\{TH\}) = (1 - p)p$ ,  $P(\{TT\}) = (1 - p)^2$ .
- ▶  $P(\{HH, HT\}) =$

- ▶ Consider toss of a biased coin:  
 $\Omega^1 = \{H, T\}$ ,  $P(\{H\}) = p$  and  $P(\{T\}) = 1 - p$ .
- ▶ If we toss this twice then  $\Omega^2 = \{HH, HT, TH, TT\}$  and we assign  
 $P(\{HH\}) = p^2$ ,  $P(\{HT\}) = p(1 - p)$ ,  
 $P(\{TH\}) = (1 - p)p$ ,  $P(\{TT\}) = (1 - p)^2$ .
- ▶  $P(\{HH, HT\}) = p^2 + p(1 - p)$

- ▶ Consider toss of a biased coin:  
 $\Omega^1 = \{H, T\}$ ,  $P(\{H\}) = p$  and  $P(\{T\}) = 1 - p$ .
- ▶ If we toss this twice then  $\Omega^2 = \{HH, HT, TH, TT\}$  and we assign  
 $P(\{HH\}) = p^2$ ,  $P(\{HT\}) = p(1 - p)$ ,  
 $P(\{TH\}) = (1 - p)p$ ,  $P(\{TT\}) = (1 - p)^2$ .
- ▶  $P(\{HH, HT\}) = p^2 + p(1 - p) = p$

- ▶ Consider toss of a biased coin:  
 $\Omega^1 = \{H, T\}$ ,  $P(\{H\}) = p$  and  $P(\{T\}) = 1 - p$ .
- ▶ If we toss this twice then  $\Omega^2 = \{HH, HT, TH, TT\}$  and we assign  
 $P(\{HH\}) = p^2$ ,  $P(\{HT\}) = p(1 - p)$ ,  
 $P(\{TH\}) = (1 - p)p$ ,  $P(\{TT\}) = (1 - p)^2$ .
- ▶  $P(\{HH, HT\}) = p^2 + p(1 - p) = p = P(\{HH, TH\})$



- ▶ Consider toss of a biased coin:  
 $\Omega^1 = \{H, T\}$ ,  $P(\{H\}) = p$  and  $P(\{T\}) = 1 - p$ .
- ▶ If we toss this twice then  $\Omega^2 = \{HH, HT, TH, TT\}$  and we assign  
 $P(\{HH\}) = p^2$ ,  $P(\{HT\}) = p(1 - p)$ ,  
 $P(\{TH\}) = (1 - p)p$ ,  $P(\{TT\}) = (1 - p)^2$ .
- ▶  $P(\{HH, HT\}) = p^2 + p(1 - p) = p = P(\{HH, TH\})$
- ▶ This assignment ensures that  $P(\{HH\})$  equals product of probability of  $H$  on 1st toss and  $H$  on second toss.

- ▶ Consider toss of a biased coin:  
 $\Omega^1 = \{H, T\}$ ,  $P(\{H\}) = p$  and  $P(\{T\}) = 1 - p$ .
- ▶ If we toss this twice then  $\Omega^2 = \{HH, HT, TH, TT\}$  and we assign  
 $P(\{HH\}) = p^2$ ,  $P(\{HT\}) = p(1 - p)$ ,  
 $P(\{TH\}) = (1 - p)p$ ,  $P(\{TT\}) = (1 - p)^2$ .
- ▶  $P(\{HH, HT\}) = p^2 + p(1 - p) = p = P(\{HH, TH\})$
- ▶ This assignment ensures that  $P(\{HH\})$  equals product of probability of  $H$  on 1st toss and  $H$  on second toss.
- ▶  $\Omega^2$  is a cartesian product of  $\Omega^1$  with itself and we essentially used products of the corresponding probabilities.

- ▶ Consider toss of a biased coin:  
 $\Omega^1 = \{H, T\}$ ,  $P(\{H\}) = p$  and  $P(\{T\}) = 1 - p$ .
- ▶ If we toss this twice then  $\Omega^2 = \{HH, HT, TH, TT\}$  and we assign  
 $P(\{HH\}) = p^2$ ,  $P(\{HT\}) = p(1 - p)$ ,  
 $P(\{TH\}) = (1 - p)p$ ,  $P(\{TT\}) = (1 - p)^2$ .
- ▶  $P(\{HH, HT\}) = p^2 + p(1 - p) = p = P(\{HH, TH\})$
- ▶ This assignment ensures that  $P(\{HH\})$  equals product of probability of  $H$  on 1st toss and  $H$  on second toss.
- ▶  $\Omega^2$  is a cartesian product of  $\Omega^1$  with itself and we essentially used products of the corresponding probabilities.
- ▶ For any independent repetitions of a random experiment we follow this.

- ▶ Consider toss of a biased coin:  
 $\Omega^1 = \{H, T\}$ ,  $P(\{H\}) = p$  and  $P(\{T\}) = 1 - p$ .
- ▶ If we toss this twice then  $\Omega^2 = \{HH, HT, TH, TT\}$  and we assign  
 $P(\{HH\}) = p^2$ ,  $P(\{HT\}) = p(1 - p)$ ,  
 $P(\{TH\}) = (1 - p)p$ ,  $P(\{TT\}) = (1 - p)^2$ .
- ▶  $P(\{HH, HT\}) = p^2 + p(1 - p) = p = P(\{HH, TH\})$
- ▶ This assignment ensures that  $P(\{HH\})$  equals product of probability of  $H$  on 1st toss and  $H$  on second toss.
- ▶  $\Omega^2$  is a cartesian product of  $\Omega^1$  with itself and we essentially used products of the corresponding probabilities.
- ▶ For any independent repetitions of a random experiment we follow this.  
 (We will look at it more formally when we consider multiple random variables).

- ▶ In many situations calculating probabilities of intersection of events is difficult.

- ▶ In many situations calculating probabilities of intersection of events is difficult.
- ▶ One often **assumes**  $A$  and  $B$  are independent to calculate  $P(AB)$ .

- ▶ In many situations calculating probabilities of intersection of events is difficult.
- ▶ One often **assumes**  $A$  and  $B$  are independent to calculate  $P(AB)$ .
- ▶ As we saw, if  $A$  and  $B$  are independent, then  $P(A|B) = P(A)$

- ▶ In many situations calculating probabilities of intersection of events is difficult.
- ▶ One often **assumes**  $A$  and  $B$  are independent to calculate  $P(AB)$ .
- ▶ As we saw, if  $A$  and  $B$  are independent, then  $P(A|B) = P(A)$
- ▶ This is often used, at an intuitive level, to justify assumption of independence.



# Independence of multiple events

# Independence of multiple events

- ▶ Events  $A_1, A_2, \dots, A_n$  are said to be (totally) independent if for any  $k$ ,  $1 \leq k \leq n$ , and any indices  $i_1, \dots, i_k$ , we have

$$P(A_{i_1} \cdots A_{i_k}) = P(A_{i_1}) \cdots P(A_{i_k})$$

# Independence of multiple events

- ▶ Events  $A_1, A_2, \dots, A_n$  are said to be (totally) independent if for any  $k$ ,  $1 \leq k \leq n$ , and any indices  $i_1, \dots, i_k$ , we have

$$P(A_{i_1} \cdots A_{i_k}) = P(A_{i_1}) \cdots P(A_{i_k})$$

- ▶ For example,  $A, B, C$  are independent if

$$P(AB) = P(A)P(B);$$

# Independence of multiple events

- ▶ Events  $A_1, A_2, \dots, A_n$  are said to be (totally) independent if for any  $k$ ,  $1 \leq k \leq n$ , and any indices  $i_1, \dots, i_k$ , we have

$$P(A_{i_1} \cdots A_{i_k}) = P(A_{i_1}) \cdots P(A_{i_k})$$

- ▶ For example,  $A, B, C$  are independent if

$$P(AB) = P(A)P(B); \quad P(AC) = P(A)P(C);$$

# Independence of multiple events

- ▶ Events  $A_1, A_2, \dots, A_n$  are said to be (totally) independent if for any  $k$ ,  $1 \leq k \leq n$ , and any indices  $i_1, \dots, i_k$ , we have

$$P(A_{i_1} \cdots A_{i_k}) = P(A_{i_1}) \cdots P(A_{i_k})$$

- ▶ For example,  $A, B, C$  are independent if

$$P(AB) = P(A)P(B); \quad P(AC) = P(A)P(C);$$

$$P(BC) = P(B)P(C);$$

# Independence of multiple events

- ▶ Events  $A_1, A_2, \dots, A_n$  are said to be (totally) independent if for any  $k$ ,  $1 \leq k \leq n$ , and any indices  $i_1, \dots, i_k$ , we have

$$P(A_{i_1} \cdots A_{i_k}) = P(A_{i_1}) \cdots P(A_{i_k})$$

- ▶ For example,  $A, B, C$  are independent if

$$P(AB) = P(A)P(B); \quad P(AC) = P(A)P(C);$$

$$P(BC) = P(B)P(C); \quad P(ABC) = P(A)P(B)P(C)$$

# Pair-wise independence

# Pair-wise independence

- Events  $A_1, A_2, \dots, A_n$  are said to be pair-wise independent if

$$P(A_i A_j) = P(A_i)P(A_j), \forall i \neq j$$



# Pair-wise independence

- ▶ Events  $A_1, A_2, \dots, A_n$  are said to be pair-wise independent if

$$P(A_i A_j) = P(A_i)P(A_j), \forall i \neq j$$

- ▶ Events may be pair-wise independent but not (totally) independent.

# Pair-wise independence

- ▶ Events  $A_1, A_2, \dots, A_n$  are said to be pair-wise independent if

$$P(A_i A_j) = P(A_i)P(A_j), \forall i \neq j$$

- ▶ Events may be pair-wise independent but not (totally) independent.
- ▶ Example: Four balls in a box inscribed with '1', '2', '3' and '123'. Let  $E_i$  be the event that number 'i' appears on a randomly drawn ball,  $i = 1, 2, 3$ .

# Pair-wise independence

- ▶ Events  $A_1, A_2, \dots, A_n$  are said to be pair-wise independent if

$$P(A_i A_j) = P(A_i)P(A_j), \forall i \neq j$$

- ▶ Events may be pair-wise independent but not (totally) independent.
- ▶ Example: Four balls in a box inscribed with '1', '2', '3' and '123'. Let  $E_i$  be the event that number 'i' appears on a randomly drawn ball,  $i = 1, 2, 3$ .
- ▶ Easy to see:  $P(E_i) = 0.5$ ,  $i = 1, 2, 3$ .

# Pair-wise independence

- ▶ Events  $A_1, A_2, \dots, A_n$  are said to be pair-wise independent if

$$P(A_i A_j) = P(A_i)P(A_j), \forall i \neq j$$

- ▶ Events may be pair-wise independent but not (totally) independent.
- ▶ Example: Four balls in a box inscribed with '1', '2', '3' and '123'. Let  $E_i$  be the event that number 'i' appears on a randomly drawn ball,  $i = 1, 2, 3$ .
- ▶ Easy to see:  $P(E_i) = 0.5$ ,  $i = 1, 2, 3$ .
- ▶  $P(E_i E_j) = 0.25$  ( $i \neq j$ )

# Pair-wise independence

- ▶ Events  $A_1, A_2, \dots, A_n$  are said to be pair-wise independent if

$$P(A_i A_j) = P(A_i)P(A_j), \forall i \neq j$$

- ▶ Events may be pair-wise independent but not (totally) independent.
- ▶ Example: Four balls in a box inscribed with '1', '2', '3' and '123'. Let  $E_i$  be the event that number 'i' appears on a randomly drawn ball,  $i = 1, 2, 3$ .
- ▶ Easy to see:  $P(E_i) = 0.5$ ,  $i = 1, 2, 3$ .
- ▶  $P(E_i E_j) = 0.25$  ( $i \neq j$ )  $\Rightarrow$  pairwise independent

# Pair-wise independence

- ▶ Events  $A_1, A_2, \dots, A_n$  are said to be pair-wise independent if

$$P(A_i A_j) = P(A_i)P(A_j), \forall i \neq j$$

- ▶ Events may be pair-wise independent but not (totally) independent.
- ▶ Example: Four balls in a box inscribed with '1', '2', '3' and '123'. Let  $E_i$  be the event that number 'i' appears on a randomly drawn ball,  $i = 1, 2, 3$ .
- ▶ Easy to see:  $P(E_i) = 0.5$ ,  $i = 1, 2, 3$ .
- ▶  $P(E_i E_j) = 0.25$  ( $i \neq j$ )  $\Rightarrow$  pairwise independent
- ▶ But,  $P(E_1 E_2 E_3) = 0.25 \neq (0.5)^3$

# Conditional Independence

- ▶ Events  $A, B$  are said to be conditionally independent given  $C$  if

$$P(AB|C) = P(A|C)P(B|C)$$

# Conditional Independence

- ▶ Events  $A, B$  are said to be conditionally independent given  $C$  if

$$P(AB|C) = P(A|C)P(B|C)$$

- ▶ If the above holds

$$P(A|BC) = \frac{P(ABC)}{P(BC)} =$$



# Conditional Independence

- ▶ Events  $A, B$  are said to be conditionally independent given  $C$  if

$$P(AB|C) = P(A|C)P(B|C)$$

- ▶ If the above holds

$$P(A|BC) = \frac{P(ABC)}{P(BC)} = \frac{P(AB|C)P(C)}{P(BC)}$$

# Conditional Independence

- ▶ Events  $A, B$  are said to be conditionally independent given  $C$  if

$$P(AB|C) = P(A|C)P(B|C)$$

- ▶ If the above holds

$$\begin{aligned} P(A|BC) &= \frac{P(ABC)}{P(BC)} = \frac{P(AB|C)P(C)}{P(BC)} \\ &= \frac{P(A|C) P(B|C)P(C)}{P(BC)} \end{aligned}$$

# Conditional Independence

- ▶ Events  $A, B$  are said to be conditionally independent given  $C$  if

$$P(AB|C) = P(A|C)P(B|C)$$

- ▶ If the above holds

$$\begin{aligned} P(A|BC) &= \frac{P(ABC)}{P(BC)} = \frac{P(AB|C)P(C)}{P(BC)} \\ &= \frac{P(A|C) P(B|C)P(C)}{P(BC)} = P(A|C) \end{aligned}$$

# Conditional Independence

- ▶ Events  $A, B$  are said to be conditionally independent given  $C$  if

$$P(AB|C) = P(A|C)P(B|C)$$

- ▶ If the above holds

$$\begin{aligned} P(A|BC) &= \frac{P(ABC)}{P(BC)} = \frac{P(AB|C)P(C)}{P(BC)} \\ &= \frac{P(A|C) P(B|C)P(C)}{P(BC)} = P(A|C) \end{aligned}$$

- ▶ Events may be conditionally independent but not independent. (e.g., 'independent' multiple tests for confirming a disease)

# Conditional Independence

- ▶ Events  $A, B$  are said to be conditionally independent given  $C$  if

$$P(AB|C) = P(A|C)P(B|C)$$

- ▶ If the above holds

$$\begin{aligned} P(A|BC) &= \frac{P(ABC)}{P(BC)} = \frac{P(AB|C)P(C)}{P(BC)} \\ &= \frac{P(A|C)P(B|C)P(C)}{P(BC)} = P(A|C) \end{aligned}$$

- ▶ Events may be conditionally independent but not independent. (e.g., 'independent' multiple tests for confirming a disease)
- ▶ It is also possible that  $A, B$  are independent but are not conditionally independent given some other event  $C$ .

# Use of conditional independence in Bayes rule

- ▶ We can write Bayes rule with multiple conditioning events.

# Use of conditional independence in Bayes rule

- We can write Bayes rule with multiple conditioning events.

$$P(A|BC) = \frac{P(BC|A)P(A)}{P(BC|A)P(A) + P(BC|A^c)P(A^c)}$$

# Use of conditional independence in Bayes rule

- ▶ We can write Bayes rule with multiple conditioning events.

$$P(A|BC) = \frac{P(BC|A)P(A)}{P(BC|A)P(A) + P(BC|A^c)P(A^c)}$$

- ▶ The above gets simplified if we assume  
 $P(BC|A) = P(B|A)P(C|A)$ ,  
 $P(BC|A^c) = P(B|A^c)P(C|A^c)$



# Use of conditional independence in Bayes rule

- ▶ We can write Bayes rule with multiple conditioning events.

$$P(A|BC) = \frac{P(BC|A)P(A)}{P(BC|A)P(A) + P(BC|A^c)P(A^c)}$$

- ▶ The above gets simplified if we assume  
 $P(BC|A) = P(B|A)P(C|A)$ ,  
 $P(BC|A^c) = P(B|A^c)P(C|A^c)$
- ▶ Consider the old example, where now we repeat the test for the disease.

# Use of conditional independence in Bayes rule

- ▶ We can write Bayes rule with multiple conditioning events.

$$P(A|BC) = \frac{P(BC|A)P(A)}{P(BC|A)P(A) + P(BC|A^c)P(A^c)}$$

- ▶ The above gets simplified if we assume  
 $P(BC|A) = P(B|A)P(C|A),$   
 $P(BC|A^c) = P(B|A^c)P(C|A^c)$
- ▶ Consider the old example, where now we repeat the test for the disease.
- ▶ Take:  $A = D, B = T_+^1, C = T_+^2.$

# Use of conditional independence in Bayes rule

- ▶ We can write Bayes rule with multiple conditioning events.

$$P(A|BC) = \frac{P(BC|A)P(A)}{P(BC|A)P(A) + P(BC|A^c)P(A^c)}$$

- ▶ The above gets simplified if we assume
$$P(BC|A) = P(B|A)P(C|A),$$
$$P(BC|A^c) = P(B|A^c)P(C|A^c)$$
- ▶ Consider the old example, where now we repeat the test for the disease.
- ▶ Take:  $A = D$ ,  $B = T_+^1$ ,  $C = T_+^2$ .
- ▶ Assuming conditional independence we can calculate the new posterior probability using the same information we had about true positive and false positive rate.

- ▶ Let us consider the example with  $P(T_+|D) = 0.99$ ,  
 $P(T_+|D^c) = 0.05$ .  $P(D) = 0.1$ .

- ▶ Let us consider the example with  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .  $P(D) = 0.1$ .
- ▶ Recall that we got  $P(D|T_+) = 0.69$ .

- ▶ Let us consider the example with  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .  $P(D) = 0.1$ .
- ▶ Recall that we got  $P(D|T_+) = 0.69$ .
- ▶ Let us suppose the same test is repeated.

- ▶ Let us consider the example with  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .  $P(D) = 0.1$ .
- ▶ Recall that we got  $P(D|T_+) = 0.69$ .
- ▶ Let us suppose the same test is repeated.

$$P(D \mid T_+^1 T_+^2) = \frac{P(T_+^1 T_+^2 \mid D)P(D)}{P(T_+^1 T_+^2 \mid D)P(D) + P(T_+^1 T_+^2 \mid D^c)P(D^c)}$$

- ▶ Let us consider the example with  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .  $P(D) = 0.1$ .
- ▶ Recall that we got  $P(D|T_+) = 0.69$ .
- ▶ Let us suppose the same test is repeated.

$$\begin{aligned}
 P(D | T_+^1 T_+^2) &= \frac{P(T_+^1 T_+^2 | D)P(D)}{P(T_+^1 T_+^2 | D)P(D) + P(T_+^1 T_+^2 | D^c)P(D^c)} \\
 &= \frac{P(T_+^1 | D)P(T_+^2 | D)P(D)}{P(T_+^1 | D)P(T_+^2 | D)P(D) + P(T_+^1 | D^c)P(T_+^2 | D^c)P(D^c)}
 \end{aligned}$$



- ▶ Let us consider the example with  $P(T_+|D) = 0.99$ ,  $P(T_+|D^c) = 0.05$ .  $P(D) = 0.1$ .
- ▶ Recall that we got  $P(D|T_+) = 0.69$ .
- ▶ Let us suppose the same test is repeated.

$$\begin{aligned}
 P(D | T_+^1 T_+^2) &= \frac{P(T_+^1 T_+^2 | D)P(D)}{P(T_+^1 T_+^2 | D)P(D) + P(T_+^1 T_+^2 | D^c)P(D^c)} \\
 &= \frac{P(T_+^1 | D)P(T_+^2 | D)P(D)}{P(T_+^1 | D)P(T_+^2 | D)P(D) + P(T_+^1 | D^c)P(T_+^2 | D^c)P(D^c)} \\
 &= \frac{0.99 * 0.99 * 0.1}{0.99 * 0.99 * 0.1 + 0.05 * 0.05 * 0.9} = 0.97
 \end{aligned}$$

