

# Recap: Continuous-Time Markov Chains

- ▶  $\{X(t), t \geq 0\}$  is a continuous-time Markov chain if

$$\begin{aligned} Pr[X(t+s) = j \mid X(s) = i, X(u) \in A_u, 0 \leq u < s] \\ = Pr[X(t+s) = j \mid X(s) = i] \end{aligned}$$

# Recap: Continuous-Time Markov Chains

- ▶  $\{X(t), t \geq 0\}$  is a continuous-time Markov chain if

$$\begin{aligned} Pr[X(t+s) = j \mid X(s) = i, X(u) \in A_u, 0 \leq u < s] \\ = Pr[X(t+s) = j \mid X(s) = i] \end{aligned}$$

implies

$$\begin{aligned} Pr[X(s) \in B_s, s \in (t, t+\tau] \mid X(s) = i, X(s'), 0 \leq s' < s < t] \\ = Pr[X(s) \in B_s, s \in (t, t+\tau] \mid X(s) = i] \end{aligned}$$

# Recap: Continuous-Time Markov Chains

- ▶  $\{X(t), t \geq 0\}$  is a continuous-time Markov chain if

$$\begin{aligned} Pr[X(t+s) = j \mid X(s) = i, X(u) \in A_u, 0 \leq u < s] \\ = Pr[X(t+s) = j \mid X(s) = i] \end{aligned}$$

implies

$$\begin{aligned} Pr[X(s) \in B_s, s \in (t, t+\tau] \mid X(s) = i, X(s'), 0 \leq s' < s < t] \\ = Pr[X(s) \in B_s, s \in (t, t+\tau] \mid X(s) = i] \end{aligned}$$

- ▶ It is called homogeneous chain if

$$Pr[X(t+s) = j \mid X(s) = i] = Pr[X(t) = j \mid X(0) = i], \forall s$$

# Recap: Transition Structure

## ► Define

$$P_{ij}(t) = Pr[X(t) = j \mid X(0) = i] = Pr[X(t+s) = j \mid X(s) = i]$$

# Recap: Transition Structure

- ▶ Define

$$P_{ij}(t) = Pr[X(t) = j \mid X(0) = i] = Pr[X(t+s) = j \mid X(s) = i]$$

- ▶ The Chain can be described as follows.

# Recap: Transition Structure

- ▶ Define

$$P_{ij}(t) = Pr[X(t) = j \mid X(0) = i] = Pr[X(t+s) = j \mid X(s) = i]$$

- ▶ The Chain can be described as follows.
- ▶ By Markov property and homogeneity, time spent in a state  $i$  is  $T_i \sim \text{exponential}(\nu_i)$

# Recap: Transition Structure

- ▶ Define

$$P_{ij}(t) = Pr[X(t) = j \mid X(0) = i] = Pr[X(t+s) = j \mid X(s) = i]$$

- ▶ The Chain can be described as follows.
- ▶ By Markov property and homogeneity, time spent in a state  $i$  is  $T_i \sim \text{exponential}(\nu_i)$
- ▶ Then, when it leaves  $i$ , it transits to state  $j$  with probability, say,  $z_{ij}$  ( $z_{ij} \geq 0$ ,  $\sum_j z_{ij} = 1$ ,  $z_{ii} = 0$ )

# Recap: Transition Structure

- ▶ Define

$$P_{ij}(t) = Pr[X(t) = j \mid X(0) = i] = Pr[X(t+s) = j \mid X(s) = i]$$

- ▶ The Chain can be described as follows.
- ▶ By Markov property and homogeneity, time spent in a state  $i$  is  $T_i \sim \text{exponential}(\nu_i)$
- ▶ Then, when it leaves  $i$ , it transits to state  $j$  with probability, say,  $z_{ij}$  ( $z_{ij} \geq 0$ ,  $\sum_j z_{ij} = 1$ ,  $z_{ii} = 0$ )
- ▶ Note that  $P_{ij}(t)$  is different from these  $z_{ij}$



## Recap: Birth-Death process

- ▶ From  $i$  the process can only go to  $i + 1$  or  $i - 1$

# Recap: Birth-Death process

- ▶ From  $i$  the process can only go to  $i + 1$  or  $i - 1$
- ▶ When in  $i$ , a 'birth event' takes it to  $i + 1$  and a 'death event' takes it to  $i - 1$

# Recap: Birth-Death process

- ▶ From  $i$  the process can only go to  $i + 1$  or  $i - 1$
- ▶ When in  $i$ , a 'birth event' takes it to  $i + 1$  and a 'death event' takes it to  $i - 1$
- ▶ In state  $i$ , birth rate is  $\lambda_i$   
(i.e., time till next birth event is  $\exp(\lambda_i)$ ).

# Recap: Birth-Death process

- ▶ From  $i$  the process can only go to  $i + 1$  or  $i - 1$
- ▶ When in  $i$ , a 'birth event' takes it to  $i + 1$  and a 'death event' takes it to  $i - 1$
- ▶ In state  $i$ , birth rate is  $\lambda_i$   
(i.e., time till next birth event is  $\exp(\lambda_i)$ ).
- ▶ In state  $i$ , death rate is  $\mu_i$   
(i.e., time till next death event is  $\exp(\mu_i)$ ).

## Recap: Birth-Death process

- ▶ From  $i$  the process can only go to  $i + 1$  or  $i - 1$
- ▶ When in  $i$ , a 'birth event' takes it to  $i + 1$  and a 'death event' takes it to  $i - 1$
- ▶ In state  $i$ , birth rate is  $\lambda_i$   
(i.e., time till next birth event is  $\exp(\lambda_i)$ ).
- ▶ In state  $i$ , death rate is  $\mu_i$   
(i.e., time till next death event is  $\exp(\mu_i)$ ).
- ▶ For a birth-death process

$$z_{i,i+1} = \frac{\lambda_i}{\lambda_i + \mu_i}; \quad z_{i,i-1} = \frac{\mu_i}{\lambda_i + \mu_i}; \quad \nu_i = \lambda_i + \mu_i$$

## Recap: Birth-Death process

- ▶ From  $i$  the process can only go to  $i + 1$  or  $i - 1$
- ▶ When in  $i$ , a 'birth event' takes it to  $i + 1$  and a 'death event' takes it to  $i - 1$
- ▶ In state  $i$ , birth rate is  $\lambda_i$   
(i.e., time till next birth event is  $\exp(\lambda_i)$ ).
- ▶ In state  $i$ , death rate is  $\mu_i$   
(i.e., time till next death event is  $\exp(\mu_i)$ ).
- ▶ For a birth-death process

$$z_{i,i+1} = \frac{\lambda_i}{\lambda_i + \mu_i}; \quad z_{i,i-1} = \frac{\mu_i}{\lambda_i + \mu_i}; \quad \nu_i = \lambda_i + \mu_i$$

- ▶ Poisson process is a special case with  $\lambda_i = \lambda$  and  $\mu_i = 0$

# Recap: Queuing system

- ▶ The state is number of people in the system

## Recap: Queuing system

- ▶ The state is number of people in the system
- ▶ People joining the queue is a Poisson process with rate  $\lambda$



## Recap: Queuing system

- ▶ The state is number of people in the system
- ▶ People joining the queue is a Poisson process with rate  $\lambda$
- ▶ The time to service each customer is independent and exponential with parameter  $\mu$ .

# Recap: Queuing system

- ▶ The state is number of people in the system
- ▶ People joining the queue is a Poisson process with rate  $\lambda$
- ▶ The time to service each customer is independent and exponential with parameter  $\mu$ .
- ▶ We assume that the arrival and service processes are independent.

## Recap: Queuing system

- ▶ The state is number of people in the system
- ▶ People joining the queue is a Poisson process with rate  $\lambda$
- ▶ The time to service each customer is independent and exponential with parameter  $\mu$ .
- ▶ We assume that the arrival and service processes are independent.
- ▶ Then this is a birth death process with

$$\lambda_n = \lambda, \quad n \geq 0 \quad \text{and} \quad \mu_n = \mu, \quad n \geq 1$$

## Recap: Queuing system

- ▶ The state is number of people in the system
- ▶ People joining the queue is a Poisson process with rate  $\lambda$
- ▶ The time to service each customer is independent and exponential with parameter  $\mu$ .
- ▶ We assume that the arrival and service processes are independent.
- ▶ Then this is a birth death process with

$$\lambda_n = \lambda, \quad n \geq 0 \quad \text{and} \quad \mu_n = \mu, \quad n \geq 1$$

- ▶ This is known as an  $M/M/1$  queue

## Recap: Queuing system

- ▶ The state is number of people in the system
- ▶ People joining the queue is a Poisson process with rate  $\lambda$
- ▶ The time to service each customer is independent and exponential with parameter  $\mu$ .
- ▶ We assume that the arrival and service processes are independent.
- ▶ Then this is a birth death process with

$$\lambda_n = \lambda, \quad n \geq 0 \quad \text{and} \quad \mu_n = \mu, \quad n \geq 1$$

- ▶ This is known as an  $M/M/1$  queue
- ▶ A variation:  $M/M/K$  queue

## Recap: Queuing system

- ▶ The state is number of people in the system
- ▶ People joining the queue is a Poisson process with rate  $\lambda$
- ▶ The time to service each customer is independent and exponential with parameter  $\mu$ .
- ▶ We assume that the arrival and service processes are independent.
- ▶ Then this is a birth death process with

$$\lambda_n = \lambda, \quad n \geq 0 \quad \text{and} \quad \mu_n = \mu, \quad n \geq 1$$

- ▶ This is known as an  $M/M/1$  queue
- ▶ A variation:  $M/M/K$  queue

$$\lambda_n = \lambda, \quad n \geq 0 \quad \text{and} \quad \mu_n = \begin{cases} n\mu & 1 \leq n \leq K \\ K\mu & n > K \end{cases}$$

## Recap: Kolmogorov backward equation

- ▶ The  $P_{ij}(t)$  satisfy the Chapman-Kolmogorov equations:

$$P_{ij}(t+s) = \sum_k P_{ik}(s)P_{kj}(t)$$

# Recap: Kolmogorov backward equation

- ▶ The  $P_{ij}(t)$  satisfy the Chapman-Kolmogorov equations:

$$P_{ij}(t+s) = \sum_k P_{ik}(s)P_{kj}(t)$$

- ▶ For a finite chain, in Matrix notation:

$$P(t+s) = P(s)P(t)$$



## Recap: Kolmogorov backward equation

- ▶ The  $P_{ij}(t)$  satisfy the Chapman-Kolmogorov equations:

$$P_{ij}(t+s) = \sum_k P_{ik}(s)P_{kj}(t)$$

- ▶ For a finite chain, in Matrix notation:

$$P(t+s) = P(s)P(t)$$

- ▶ Using this, we get Kolmogorov backward equation

$$P'_{ij}(t) = \sum_{k \neq i} q_{ik} P_{kj}(t) - \left( \sum_{k \neq i} q_{ik} \right) P_{ij}(t), \quad \forall i, j$$

where

$$q_{ij} = \lim_{h \rightarrow 0} \frac{P_{ij}(h) - \delta_{ij}}{h}, \quad i \neq j$$

## Recap: Kolmogorov backward equation

- ▶ The  $P_{ij}(t)$  satisfy the Chapman-Kolmogorov equations:

$$P_{ij}(t+s) = \sum_k P_{ik}(s)P_{kj}(t)$$

- ▶ For a finite chain, in Matrix notation:

$$P(t+s) = P(s)P(t)$$

- ▶ Using this, we get Kolmogorov backward equation

$$P'_{ij}(t) = \sum_{k \neq i} q_{ik}P_{kj}(t) - \left( \sum_{k \neq i} q_{ik} \right) P_{ij}(t), \quad \forall i, j$$

where

$$q_{ij} = \lim_{h \rightarrow 0} \frac{P_{ij}(h) - \delta_{ij}}{h}, \quad i \neq j$$

- ▶ The  $q_{ij}$  – the infinitesimal generator of the chain.

## Recap: Kolmogorov backward equation

- ▶ The  $P_{ij}(t)$  satisfy the Chapman-Kolmogorov equations:

$$P_{ij}(t+s) = \sum_k P_{ik}(s)P_{kj}(t)$$

- ▶ For a finite chain, in Matrix notation:

$$P(t+s) = P(s)P(t)$$

- ▶ Using this, we get Kolmogorov backward equation

$$P'_{ij}(t) = \sum_{k \neq i} q_{ik} P_{kj}(t) - \left( \sum_{k \neq i} q_{ik} \right) P_{ij}(t), \quad \forall i, j$$

where

$$q_{ij} = \lim_{h \rightarrow 0} \frac{P_{ij}(h) - \delta_{ij}}{h}, \quad i \neq j$$

- ▶ The  $q_{ij}$  – the infinitesimal generator of the chain.
- ▶  $q_{ij}$  is the rate of transitions out of  $i$  into  $j$

# Recap: Kolmogorov backward equation

- ▶ The  $P_{ij}(t)$  satisfy the Chapman-Kolmogorov equations:

$$P_{ij}(t+s) = \sum_k P_{ik}(s)P_{kj}(t)$$

- ▶ For a finite chain, in Matrix notation:

$$P(t+s) = P(s)P(t)$$

- ▶ Using this, we get Kolmogorov backward equation

$$P'_{ij}(t) = \sum_{k \neq i} q_{ik}P_{kj}(t) - \left( \sum_{k \neq i} q_{ik} \right) P_{ij}(t), \quad \forall i, j$$

where

$$q_{ij} = \lim_{h \rightarrow 0} \frac{P_{ij}(h) - \delta_{ij}}{h}, \quad i \neq j$$

- ▶ The  $q_{ij}$  – the infinitesimal generator of the chain.
- ▶  $q_{ij}$  is the rate of transitions out of  $i$  into  $j$
- ▶ For a birth-death chain,  $q_{i,i+1} = \lambda_i$  and  $q_{i,i-1} = \mu_i$

## Recap: Obtaining $P_{ij}(t)$

- The Kolmogorov backward equation is

$$P'_{ij}(t) = \sum_{k \neq i} q_{ik} P_{kj}(t) - \left( \sum_{k \neq i} q_{ik} \right) P_{ij}(t)$$

## Recap: Obtaining $P_{ij}(t)$

- ▶ The Kolmogorov backward equation is

$$P'_{ij}(t) = \sum_{k \neq i} q_{ik} P_{kj}(t) - \left( \sum_{k \neq i} q_{ik} \right) P_{ij}(t)$$

- ▶ For a birth-death chain the equation becomes

$$P'_{ij}(t) = \lambda_i P_{i+1,j}(t) + \mu_i P_{i-1,j}(t) - (\lambda_i + \mu_i) P_{ij}(t)$$

## Recap: Obtaining $P_{ij}(t)$

- ▶ The Kolmogorov backward equation is

$$P'_{ij}(t) = \sum_{k \neq i} q_{ik} P_{kj}(t) - \left( \sum_{k \neq i} q_{ik} \right) P_{ij}(t)$$

- ▶ For a birth-death chain the equation becomes

$$P'_{ij}(t) = \lambda_i P_{i+1,j}(t) + \mu_i P_{i-1,j}(t) - (\lambda_i + \mu_i) P_{ij}(t)$$

- ▶ We can solve these to get  $P_{ij}(t)$

## Recap: 2-state birth-death chain

$$P'_{ij}(t) = \lambda_i P_{i+1,j}(t) + \mu_i P_{i-1,j}(t) - (\lambda_i + \mu_i) P_{ij}(t)$$

- For the two state chain, these equations are

$$P'_{00}(t) = \lambda_0 P_{10}(t) - \lambda_0 P_{00}(t)$$

$$P'_{01}(t) = \lambda_0 P_{11}(t) - \lambda_0 P_{01}(t)$$

$$P'_{10}(t) = \mu_1 P_{00}(t) - \mu_1 P_{10}(t)$$

$$P'_{11}(t) = \mu_1 P_{01}(t) - \mu_1 P_{11}(t)$$



## Recap: 2-state birth-death chain

$$P'_{ij}(t) = \lambda_i P_{i+1,j}(t) + \mu_i P_{i-1,j}(t) - (\lambda_i + \mu_i) P_{ij}(t)$$

- For the two state chain, these equations are

$$P'_{00}(t) = \lambda_0 P_{10}(t) - \lambda_0 P_{00}(t)$$

$$P'_{01}(t) = \lambda_0 P_{11}(t) - \lambda_0 P_{01}(t)$$

$$P'_{10}(t) = \mu_1 P_{00}(t) - \mu_1 P_{10}(t)$$

$$P'_{11}(t) = \mu_1 P_{01}(t) - \mu_1 P_{11}(t)$$

- Solving these we can show

$$P_{00}(t) = \frac{\lambda}{\lambda + \mu} e^{-(\lambda + \mu)t} + \frac{\mu}{\lambda + \mu}$$

## Recap: 2-state birth-death chain

$$P'_{ij}(t) = \lambda_i P_{i+1,j}(t) + \mu_i P_{i-1,j}(t) - (\lambda_i + \mu_i) P_{ij}(t)$$

- For the two state chain, these equations are

$$P'_{00}(t) = \lambda_0 P_{10}(t) - \lambda_0 P_{00}(t)$$

$$P'_{01}(t) = \lambda_0 P_{11}(t) - \lambda_0 P_{01}(t)$$

$$P'_{10}(t) = \mu_1 P_{00}(t) - \mu_1 P_{10}(t)$$

$$P'_{11}(t) = \mu_1 P_{01}(t) - \mu_1 P_{11}(t)$$

- Solving these we can show

$$P_{00}(t) = \frac{\lambda}{\lambda + \mu} e^{-(\lambda + \mu)t} + \frac{\mu}{\lambda + \mu}$$

- We get a system of equations like this for any finite chain.

## Recap: Finite chains

- ▶ The Chapman-Kolmogorov equation gives

$$P(t + s) = P(t) P(s)$$

## Recap: Finite chains

- ▶ The Chapman-Kolmogorov equation gives

$$P(t + s) = P(t) P(s)$$

- ▶ Differentiating the above with respect to  $t$  and at  $t = 0$ ,

$$P'(s) = Q P(s), \text{ where } Q = P'(0)$$

## Recap: Finite chains

- ▶ The Chapman-Kolmogorov equation gives

$$P(t + s) = P(t) P(s)$$

- ▶ Differentiating the above with respect to  $t$  and at  $t = 0$ ,

$$P'(s) = Q P(s), \quad \text{where } Q = P'(0)$$

- ▶ The solution for the above ODE is

$$P(t) = e^{tQ}, \quad \text{because } P(0) = I$$

# Recap: Finite chains

- ▶ The Chapman-Kolmogorov equation gives

$$P(t + s) = P(t) P(s)$$

- ▶ Differentiating the above with respect to  $t$  and at  $t = 0$ ,

$$P'(s) = Q P(s), \quad \text{where } Q = P'(0)$$

- ▶ The solution for the above ODE is

$$P(t) = e^{tQ}, \quad \text{because } P(0) = I$$

- ▶ The  $Q$  matrix has elements  $q_{ij}$  defined earlier with  $q_{ii} = -\sum_{k \neq i} q_{ik}$ .

# Recap: Kolmogorov Forward equation

- ▶ The Chapman-Kolmogorov equation also gives us

$$P_{ij}(t+h) = \sum_k P_{ik}(t) P_{kj}(h)$$

# Recap: Kolmogorov Forward equation

- ▶ The Chapman-Kolmogorov equation also gives us

$$P_{ij}(t+h) = \sum_k P_{ik}(t) P_{kj}(h)$$

- ▶ Similar algebra as earlier gives us

$$P'_{ij}(t) = \sum_{k \neq j} P_{ik}(t) q_{kj} - \left( \sum_{k \neq j} q_{jk} \right) P_{ij}(t)$$



# Recap: Kolmogorov Forward equation

- ▶ The Chapman-Kolmogorov equation also gives us

$$P_{ij}(t+h) = \sum_k P_{ik}(t) P_{kj}(h)$$

- ▶ Similar algebra as earlier gives us

$$P'_{ij}(t) = \sum_{k \neq j} P_{ik}(t) q_{kj} - \left( \sum_{k \neq j} q_{jk} \right) P_{ij}(t)$$

- ▶ This is known as Kolmogorov forward equation

# Recap: Transient and recurrent states

► We define

$$T_i = \min\{t > 0 : X(t) \neq i\} \quad f_i = \min\{t : t > T_i, X(t) = i\}$$

# Recap: Transient and recurrent states

- We define

$$T_i = \min\{t > 0 : X(t) \neq i\} \quad f_i = \min\{t : t > T_i, X(t) = i\}$$

- A state  $i$  is said to be

# Recap: Transient and recurrent states

- ▶ We define

$$T_i = \min\{t > 0 : X(t) \neq i\} \quad f_i = \min\{t : t > T_i, X(t) = i\}$$

- ▶ A state  $i$  is said to be
  - ▶ Transient if  $Pr[f_i < \infty \mid X(0) = i] < 1$

# Recap: Transient and recurrent states

- ▶ We define

$$T_i = \min\{t > 0 : X(t) \neq i\} \quad f_i = \min\{t : t > T_i, X(t) = i\}$$

- ▶ A state  $i$  is said to be
  - ▶ Transient if  $Pr[f_i < \infty \mid X(0) = i] < 1$
  - ▶ Recurrent if  $Pr[f_i < \infty \mid X(0) = i] = 1$

# Recap: Irreducibility, positive recurrence

- ▶ Most of the other definitions are also similar to the case of discrete chains

# Recap: Irreducibility, positive recurrence

- ▶ Most of the other definitions are also similar to the case of discrete chains
- ▶ The chain is irreducible if for all  $i, j$  there is a positive probability of going from  $i$  to  $j$  in some finite time:  
 $P_{ij}(t) > 0$  for some  $t$

# Recap: Irreducibility, positive recurrence

- ▶ Most of the other definitions are also similar to the case of discrete chains
- ▶ The chain is irreducible if for all  $i, j$  there is a positive probability of going from  $i$  to  $j$  in some finite time:  
 $P_{ij}(t) > 0$  for some  $t$
- ▶ A recurrent state is positive recurrent if mean time to return is finite:  $E[f_i \mid X(0) = i] < \infty$



# Recap: Irreducibility, positive recurrence

- ▶ Most of the other definitions are also similar to the case of discrete chains
- ▶ The chain is irreducible if for all  $i, j$  there is a positive probability of going from  $i$  to  $j$  in some finite time:  
 $P_{ij}(t) > 0$  for some  $t$
- ▶ A recurrent state is positive recurrent if mean time to return is finite:  $E[f_i \mid X(0) = i] < \infty$   
Otherwise it is null recurrent

# Recap: Irreducibility, positive recurrence

- ▶ Most of the other definitions are also similar to the case of discrete chains
- ▶ The chain is irreducible if for all  $i, j$  there is a positive probability of going from  $i$  to  $j$  in some finite time:  
 $P_{ij}(t) > 0$  for some  $t$
- ▶ A recurrent state is positive recurrent if mean time to return is finite:  $E[f_i \mid X(0) = i] < \infty$   
Otherwise it is null recurrent
- ▶ An irreducible positive recurrent chain would have a unique stationary distribution

# Recap: Irreducibility, positive recurrence

- ▶ Most of the other definitions are also similar to the case of discrete chains
- ▶ The chain is irreducible if for all  $i, j$  there is a positive probability of going from  $i$  to  $j$  in some finite time:  
 $P_{ij}(t) > 0$  for some  $t$
- ▶ A recurrent state is positive recurrent if mean time to return is finite:  $E[f_i \mid X(0) = i] < \infty$   
Otherwise it is null recurrent
- ▶ An irreducible positive recurrent chain would have a unique stationary distribution
- ▶ There is no concept of periodicity in the continuous time case

# Recap: Irreducibility, positive recurrence

- ▶ Most of the other definitions are also similar to the case of discrete chains
- ▶ The chain is irreducible if for all  $i, j$  there is a positive probability of going from  $i$  to  $j$  in some finite time:  
 $P_{ij}(t) > 0$  for some  $t$
- ▶ A recurrent state is positive recurrent if mean time to return is finite:  $E[f_i \mid X(0) = i] < \infty$   
Otherwise it is null recurrent
- ▶ An irreducible positive recurrent chain would have a unique stationary distribution
- ▶ There is no concept of periodicity in the continuous time case
- ▶ An irreducible positive recurrent chain would be called an ergodic chain

# Recap: Distribution of $X(t)$

► Define

$$\pi_j(t) = Pr[X(t) = j] = \sum_i \pi_i(0) P_{ij}(t)$$

# Recap: Distribution of $X(t)$

- ▶ Define

$$\pi_j(t) = Pr[X(t) = j] = \sum_i \pi_i(0) P_{ij}(t)$$

- ▶ The above equation is true for general infinite chains.

# Recap: Distribution of $X(t)$

- ▶ Define

$$\pi_j(t) = Pr[X(t) = j] = \sum_i \pi_i(0) P_{ij}(t)$$

- ▶ The above equation is true for general infinite chains.
- ▶ For a finite chain, taking  $\pi$  as a row vector,

$$\pi(t) = \pi(0) P(t) = \pi(0) e^{Qt}$$

# Recap: Stationary distribution

- ▶ We say  $\pi$  is a stationary distribution if

$$\pi(0) = \pi \quad \Rightarrow \quad \pi(t) = \pi, \quad \forall t$$



# Recap: Stationary distribution

- ▶ We say  $\pi$  is a stationary distribution if

$$\pi(0) = \pi \Rightarrow \pi(t) = \pi, \forall t$$

- ▶ Any stationary distribution  $\pi$  has to satisfy

$$\sum_{k \neq j} q_{kj} \pi_k = \pi_j \sum_{k \neq j} q_{jk}$$

# Recap: Stationary distribution

- ▶ We say  $\pi$  is a stationary distribution if

$$\pi(0) = \pi \Rightarrow \pi(t) = \pi, \forall t$$

- ▶ Any stationary distribution  $\pi$  has to satisfy

$$\sum_{k \neq j} q_{kj} \pi_k = \pi_j \sum_{k \neq j} q_{jk}$$

- ▶ The above equation is known as a balance equation



- ▶ All the processes we considered so far are discrete-state processes.

- ▶ All the processes we considered so far are discrete-state processes.
- ▶ The random variables are discrete random variables.

- ▶ All the processes we considered so far are discrete-state processes.
- ▶ The random variables are discrete random variables.
- ▶ This is so in Markov chains (both discrete and continuous time) and hence in the Poisson process.

- ▶ All the processes we considered so far are discrete-state processes.
- ▶ The random variables are discrete random variables.
- ▶ This is so in Markov chains (both discrete and continuous time) and hence in the Poisson process.
- ▶ We next consider processes where the random variables are continuous type.

- ▶ All the processes we considered so far are discrete-state processes.
- ▶ The random variables are discrete random variables.
- ▶ This is so in Markov chains (both discrete and continuous time) and hence in the Poisson process.
- ▶ We next consider processes where the random variables are continuous type.
- ▶ We consider the Brownian Motion process.



- ▶ All the processes we considered so far are discrete-state processes.
- ▶ The random variables are discrete random variables.
- ▶ This is so in Markov chains (both discrete and continuous time) and hence in the Poisson process.
- ▶ We next consider processes where the random variables are continuous type.
- ▶ We consider the Brownian Motion process.
- ▶ We first start with a simpler process, namely, the random walk.

# Random Walk

- ▶ Let  $Z_i$  be iid with  $Pr[Z_i = +s] = Pr[Z_i = -s] = 0.5$

# Random Walk

- ▶ Let  $Z_i$  be iid with  $Pr[Z_i = +s] = Pr[Z_i = -s] = 0.5$
- ▶ Define (for some fixed  $T$ ) a continuous-time process  $X(t)$

# Random Walk

- ▶ Let  $Z_i$  be iid with  $Pr[Z_i = +s] = Pr[Z_i = -s] = 0.5$
- ▶ Define (for some fixed  $T$ ) a continuous-time process  $X(t)$

$$X(nT) = Z_1 + Z_2 + \cdots + Z_n$$

# Random Walk

- ▶ Let  $Z_i$  be iid with  $Pr[Z_i = +s] = Pr[Z_i = -s] = 0.5$
- ▶ Define (for some fixed  $T$ ) a continuous-time process  $X(t)$

$$X(nT) = Z_1 + Z_2 + \cdots + Z_n$$

$$X(t) = X(nT), \quad \text{for } nT \leq t < (n+1)T$$

# Random Walk

- ▶ Let  $Z_i$  be iid with  $Pr[Z_i = +s] = Pr[Z_i = -s] = 0.5$
- ▶ Define (for some fixed  $T$ ) a continuous-time process  $X(t)$

$$X(nT) = Z_1 + Z_2 + \cdots + Z_n$$

$$X(t) = X(nT), \quad \text{for } nT \leq t < (n+1)T$$

- ▶ Viewed as a discrete-time process,  $X(nT)$ , is a Markov chain.

# Random Walk

- ▶ Let  $Z_i$  be iid with  $Pr[Z_i = +s] = Pr[Z_i = -s] = 0.5$
- ▶ Define (for some fixed  $T$ ) a continuous-time process  $X(t)$

$$\begin{aligned}X(nT) &= Z_1 + Z_2 + \cdots + Z_n \\X(t) &= X(nT), \quad \text{for } nT \leq t < (n+1)T\end{aligned}$$

- ▶ Viewed as a discrete-time process,  $X(nT)$ , is a Markov chain.
- ▶ Called a (one dimensional) random walk

# Random Walk

- ▶ Let  $Z_i$  be iid with  $Pr[Z_i = +s] = Pr[Z_i = -s] = 0.5$
- ▶ Define (for some fixed  $T$ ) a continuous-time process  $X(t)$

$$\begin{aligned}X(nT) &= Z_1 + Z_2 + \cdots + Z_n \\X(t) &= X(nT), \quad \text{for } nT \leq t < (n+1)T\end{aligned}$$

- ▶ Viewed as a discrete-time process,  $X(nT)$ , is a Markov chain.
- ▶ Called a (one dimensional) random walk
- ▶ It is the position after  $n$  random steps



# Random Walk

- ▶ Let  $Z_i$  be iid with  $Pr[Z_i = +s] = Pr[Z_i = -s] = 0.5$
- ▶ Define (for some fixed  $T$ ) a continuous-time process  $X(t)$

$$\begin{aligned}X(nT) &= Z_1 + Z_2 + \cdots + Z_n \\X(t) &= X(nT), \quad \text{for } nT \leq t < (n+1)T\end{aligned}$$

- ▶ Viewed as a discrete-time process,  $X(nT)$ , is a Markov chain.
- ▶ Called a (one dimensional) random walk
- ▶ It is the position after  $n$  random steps
- ▶  $X(t)$  is piece-wise constant interpolation of  $X(nT)$

# Random Walk

- ▶ Let  $Z_i$  be iid with  $Pr[Z_i = +s] = Pr[Z_i = -s] = 0.5$
- ▶ Define (for some fixed  $T$ ) a continuous-time process  $X(t)$

$$\begin{aligned}X(nT) &= Z_1 + Z_2 + \cdots + Z_n \\X(t) &= X(nT), \quad \text{for } nT \leq t < (n+1)T\end{aligned}$$

- ▶ Viewed as a discrete-time process,  $X(nT)$ , is a Markov chain.
- ▶ Called a (one dimensional) random walk
- ▶ It is the position after  $n$  random steps
- ▶  $X(t)$  is piece-wise constant interpolation of  $X(nT)$
- ▶ We could have also use piece-wise linear interpolation



► We have  $EZ_i = 0$  and  $E[Z_i^2] = s^2$

- ▶ We have  $EZ_i = 0$  and  $E[Z_i^2] = s^2$
- ▶ Hence,  $E[X(nT)] = 0$  and  $E[X^2(nT)] = ns^2$

- ▶ We have  $E Z_i = 0$  and  $E[Z_i^2] = s^2$
- ▶ Hence,  $E[X(nT)] = 0$  and  $E[X^2(nT)] = ns^2$
- ▶ For large  $n$ ,  $\frac{X(nT)}{s\sqrt{n}}$  would be Gaussian

- ▶ We have  $EZ_i = 0$  and  $E[Z_i^2] = s^2$
- ▶ Hence,  $E[X(nT)] = 0$  and  $E[X^2(nT)] = ns^2$
- ▶ For large  $n$ ,  $\frac{X(nT)}{s\sqrt{n}}$  would be Gaussian

$$Pr \left[ \frac{X(nT)}{s\sqrt{n}} \leq y \right] \approx \Phi(y)$$

where  $\Phi$  is distribution function of standard Normal

- ▶ We have  $EZ_i = 0$  and  $E[Z_i^2] = s^2$
- ▶ Hence,  $E[X(nT)] = 0$  and  $E[X^2(nT)] = ns^2$
- ▶ For large  $n$ ,  $\frac{X(nT)}{s\sqrt{n}}$  would be Gaussian

$$Pr \left[ \frac{X(nT)}{s\sqrt{n}} \leq y \right] \approx \Phi(y)$$

where  $\Phi$  is distribution function of standard Normal

- ▶ For any  $t$ ,  $X(t)$  is  $X(nT)$  for  $n = [t/T]$ .



- ▶ We have  $EZ_i = 0$  and  $E[Z_i^2] = s^2$
- ▶ Hence,  $E[X(nT)] = 0$  and  $E[X^2(nT)] = ns^2$
- ▶ For large  $n$ ,  $\frac{X(nT)}{s\sqrt{n}}$  would be Gaussian

$$Pr \left[ \frac{X(nT)}{s\sqrt{n}} \leq y \right] \approx \Phi(y)$$

where  $\Phi$  is distribution function of standard Normal

- ▶ For any  $t$ ,  $X(t)$  is  $X(nT)$  for  $n = [t/T]$ .  
Large  $n$  would mean large  $t$ . Hence

$$Pr[X(t) \leq ms] = Pr \left[ \frac{X(t)}{s\sqrt{n}} \leq \frac{ms}{s\sqrt{n}} \right] \approx \Phi \left( \frac{m}{\sqrt{n}} \right), \quad \text{for large } t$$

- ▶ We have  $EZ_i = 0$  and  $E[Z_i^2] = s^2$
- ▶ Hence,  $E[X(nT)] = 0$  and  $E[X^2(nT)] = ns^2$
- ▶ For large  $n$ ,  $\frac{X(nT)}{s\sqrt{n}}$  would be Gaussian

$$Pr \left[ \frac{X(nT)}{s\sqrt{n}} \leq y \right] \approx \Phi(y)$$

where  $\Phi$  is distribution function of standard Normal

- ▶ For any  $t$ ,  $X(t)$  is  $X(nT)$  for  $n = [t/T]$ .  
Large  $n$  would mean large  $t$ . Hence

$$Pr[X(t) \leq ms] = Pr \left[ \frac{X(t)}{s\sqrt{n}} \leq \frac{ms}{s\sqrt{n}} \right] \approx \Phi \left( \frac{m}{\sqrt{n}} \right), \quad \text{for large } t$$

- ▶ We are interested in limit of this process as  $T \rightarrow 0$

► Consider  $t = nT$

► Consider  $t = nT$

$$E[X^2(t)] = ns^2 = s^2 \frac{t}{T}$$

- Consider  $t = nT$

$$E[X^2(t)] = ns^2 = s^2 \frac{t}{T}$$

- If we let  $T \rightarrow 0$  then the variance goes to infinity (the process goes to infinity) unless we let  $s$  also go to zero.

- ▶ Consider  $t = nT$

$$E[X^2(t)] = ns^2 = s^2 \frac{t}{T}$$

- ▶ If we let  $T \rightarrow 0$  then the variance goes to infinity (the process goes to infinity) unless we let  $s$  also to go to zero.
- ▶ We actually need  $s^2$  to go to zero at the same rate as  $T$ .

- ▶ Consider  $t = nT$

$$E[X^2(t)] = ns^2 = s^2 \frac{t}{T}$$

- ▶ If we let  $T \rightarrow 0$  then the variance goes to infinity (the process goes to infinity) unless we let  $s$  also to go to zero.
- ▶ We actually need  $s^2$  to go to zero at the same rate as  $T$ .
- ▶ So, we keep  $s^2 = \alpha T$  and let  $T$  go to zero.

- ▶ Consider  $t = nT$

$$E[X^2(t)] = ns^2 = s^2 \frac{t}{T}$$

- ▶ If we let  $T \rightarrow 0$  then the variance goes to infinity (the process goes to infinity) unless we let  $s$  also to go to zero.
- ▶ We actually need  $s^2$  to go to zero at the same rate as  $T$ .
- ▶ So, we keep  $s^2 = \alpha T$  and let  $T$  go to zero.
- ▶ Define

$$W(t) = \lim_{T \rightarrow 0, s^2 = \alpha T} X(t)$$



- ▶ Consider  $t = nT$

$$E[X^2(t)] = ns^2 = s^2 \frac{t}{T}$$

- ▶ If we let  $T \rightarrow 0$  then the variance goes to infinity (the process goes to infinity) unless we let  $s$  also to go to zero.
- ▶ We actually need  $s^2$  to go to zero at the same rate as  $T$ .
- ▶ So, we keep  $s^2 = \alpha T$  and let  $T$  go to zero.
- ▶ Define

$$W(t) = \lim_{T \rightarrow 0, s^2 = \alpha T} X(t)$$

This is called the Wiener Process or Brownian motion.

- ▶ Consider  $t = nT$

$$E[X^2(t)] = ns^2 = s^2 \frac{t}{T}$$

- ▶ If we let  $T \rightarrow 0$  then the variance goes to infinity (the process goes to infinity) unless we let  $s$  also go to zero.
- ▶ We actually need  $s^2$  to go to zero at the same rate as  $T$ .
- ▶ So, we keep  $s^2 = \alpha T$  and let  $T$  go to zero.
- ▶ Define

$$W(t) = \lim_{T \rightarrow 0, s^2 = \alpha T} X(t)$$

This is called the Wiener Process or Brownian motion.  
This result is known as Donsker's theorem

- ▶ Consider  $t = nT$

$$E[X^2(t)] = ns^2 = s^2 \frac{t}{T}$$

- ▶ If we let  $T \rightarrow 0$  then the variance goes to infinity (the process goes to infinity) unless we let  $s$  also go to zero.
- ▶ We actually need  $s^2$  to go to zero at the same rate as  $T$ .
- ▶ So, we keep  $s^2 = \alpha T$  and let  $T$  go to zero.
- ▶ Define

$$W(t) = \lim_{T \rightarrow 0, s^2 = \alpha T} X(t)$$

This is called the Wiener Process or Brownian motion.

This result is known as Donsker's theorem

- ▶ Let us intuitively see some properties of  $W(t)$

- We have seen that for  $n = \lceil t/T \rceil$ ,

$$\Pr[X(t) \leq ms] \approx \Phi\left(\frac{m}{\sqrt{n}}\right)$$

- ▶ We have seen that for  $n = \lceil t/T \rceil$ ,

$$\Pr[X(t) \leq ms] \approx \Phi\left(\frac{m}{\sqrt{n}}\right)$$

- ▶ Let  $w = ms$  and  $t = nT$ . Then

- We have seen that for  $n = \lceil t/T \rceil$ ,

$$\Pr[X(t) \leq ms] \approx \Phi\left(\frac{m}{\sqrt{n}}\right)$$

- Let  $w = ms$  and  $t = nT$ . Then

$$\frac{m}{\sqrt{n}} = \frac{w/s}{\sqrt{t/T}}$$

- We have seen that for  $n = \lceil t/T \rceil$ ,

$$\Pr[X(t) \leq ms] \approx \Phi\left(\frac{m}{\sqrt{n}}\right)$$

- Let  $w = ms$  and  $t = nT$ . Then

$$\frac{m}{\sqrt{n}} = \frac{w/s}{\sqrt{t/T}} = \frac{w}{\sqrt{t}} \sqrt{\frac{T}{s^2}}$$

- We have seen that for  $n = \lceil t/T \rceil$ ,

$$\Pr[X(t) \leq ms] \approx \Phi\left(\frac{m}{\sqrt{n}}\right)$$

- Let  $w = ms$  and  $t = nT$ . Then

$$\frac{m}{\sqrt{n}} = \frac{w/s}{\sqrt{t/T}} = \frac{w}{\sqrt{t}} \sqrt{\frac{T}{s^2}} = \frac{w}{\sqrt{\alpha t}}$$



- We have seen that for  $n = \lceil t/T \rceil$ ,

$$\Pr[X(t) \leq ms] \approx \Phi\left(\frac{m}{\sqrt{n}}\right)$$

- Let  $w = ms$  and  $t = nT$ . Then

$$\frac{m}{\sqrt{n}} = \frac{w/s}{\sqrt{t/T}} = \frac{w}{\sqrt{t}} \sqrt{\frac{T}{s^2}} = \frac{w}{\sqrt{\alpha t}}$$

- $W(t)$  is limit of  $X(t)$  as  $T$  goes to zero

- ▶ We have seen that for  $n = \lceil t/T \rceil$ ,

$$\Pr[X(t) \leq ms] \approx \Phi\left(\frac{m}{\sqrt{n}}\right)$$

- ▶ Let  $w = ms$  and  $t = nT$ . Then

$$\frac{m}{\sqrt{n}} = \frac{w/s}{\sqrt{t/T}} = \frac{w}{\sqrt{t}} \sqrt{\frac{T}{s^2}} = \frac{w}{\sqrt{\alpha t}}$$

- ▶  $W(t)$  is limit of  $X(t)$  as  $T$  goes to zero
- ▶ As  $T$  goes to zero, any  $t$  is 'large  $n$ '.

- ▶ We have seen that for  $n = \lceil t/T \rceil$ ,

$$Pr[X(t) \leq ms] \approx \Phi\left(\frac{m}{\sqrt{n}}\right)$$

- ▶ Let  $w = ms$  and  $t = nT$ . Then

$$\frac{m}{\sqrt{n}} = \frac{w/s}{\sqrt{t/T}} = \frac{w}{\sqrt{t}} \sqrt{\frac{T}{s^2}} = \frac{w}{\sqrt{\alpha t}}$$

- ▶  $W(t)$  is limit of  $X(t)$  as  $T$  goes to zero
- ▶ As  $T$  goes to zero, any  $t$  is 'large n'.
- ▶ Hence we can expect

$$Pr[W(t) \leq w] = \Phi\left(\frac{w}{\sqrt{\alpha t}}\right)$$

- ▶ We have seen that for  $n = \lfloor t/T \rfloor$ ,

$$Pr[X(t) \leq ms] \approx \Phi\left(\frac{m}{\sqrt{n}}\right)$$

- ▶ Let  $w = ms$  and  $t = nT$ . Then

$$\frac{m}{\sqrt{n}} = \frac{w/s}{\sqrt{t/T}} = \frac{w}{\sqrt{t}} \sqrt{\frac{T}{s^2}} = \frac{w}{\sqrt{\alpha t}}$$

- ▶  $W(t)$  is limit of  $X(t)$  as  $T$  goes to zero
- ▶ As  $T$  goes to zero, any  $t$  is 'large n'.
- ▶ Hence we can expect

$$Pr[W(t) \leq w] = \Phi\left(\frac{w}{\sqrt{\alpha t}}\right)$$

$$\Rightarrow W(t) \sim \mathcal{N}(0, \alpha t)$$

- We had  $Z_i$  iid and defined

$$X(nT) = Z_1 + Z_2 + \cdots + Z_n$$

- ▶ We had  $Z_i$  iid and defined

$$X(nT) = Z_1 + Z_2 + \cdots + Z_n$$

- ▶ Hence we get

$$X((m+n)T) - X(nT) = Z_{n+1} + \cdots + Z_{n+m}$$

- ▶ We had  $Z_i$  iid and defined

$$X(nT) = Z_1 + Z_2 + \cdots + Z_n$$

- ▶ Hence we get

$$X((m+n)T) - X(nT) = Z_{n+1} + \cdots + Z_{n+m}$$

Thus,  $X(nT)$  is independent of  $X((m+n)T) - X(nT)$ .

- ▶ We had  $Z_i$  iid and defined

$$X(nT) = Z_1 + Z_2 + \cdots + Z_n$$

- ▶ Hence we get

$$X((m+n)T) - X(nT) = Z_{n+1} + \cdots + Z_{n+m}$$

Thus,  $X(nT)$  is independent of  $X((m+n)T) - X(nT)$ .

- ▶ Hence the  $X(nT)$  process has independent increments



- ▶ We had  $Z_i$  iid and defined

$$X(nT) = Z_1 + Z_2 + \cdots + Z_n$$

- ▶ Hence we get

$$X((m+n)T) - X(nT) = Z_{n+1} + \cdots + Z_{n+m}$$

Thus,  $X(nT)$  is independent of  $X((m+n)T) - X(nT)$ .

- ▶ Hence the  $X(nT)$  process has independent increments
- ▶ Hence, we can expect  $W(t)$  to be a process with independent increments

► We have

$$X((m+n+k)T) - X((n+k)T) = Z_{n+k+1} + \cdots + Z_{n+k+m}$$

► We have

$$\begin{aligned} X((m+n+k)T) - X((n+k)T) &= Z_{n+k+1} + \cdots + Z_{n+k+m} \\ X((m+n)T) - X(nT) &= Z_{n+1} + \cdots + Z_{n+m} \end{aligned}$$

► We have

$$\begin{aligned}X((m+n+k)T) - X((n+k)T) &= Z_{n+k+1} + \cdots + Z_{n+k+m} \\X((m+n)T) - X(nT) &= Z_{n+1} + \cdots + Z_{n+m}\end{aligned}$$

Both are sums of  $m$  of the iid  $Z_i$ 's

► We have

$$\begin{aligned}X((m+n+k)T) - X((n+k)T) &= Z_{n+k+1} + \cdots + Z_{n+k+m} \\X((m+n)T) - X(nT) &= Z_{n+1} + \cdots + Z_{n+m}\end{aligned}$$

Both are sums of  $m$  of the iid  $Z_i$ 's

► Hence both would have the same distribution

- ▶ We have

$$\begin{aligned}X((m+n+k)T) - X((n+k)T) &= Z_{n+k+1} + \cdots + Z_{n+k+m} \\X((m+n)T) - X(nT) &= Z_{n+1} + \cdots + Z_{n+m}\end{aligned}$$

Both are sums of  $m$  of the iid  $Z_i$ 's

- ▶ Hence both would have the same distribution
- ▶ Thus  $X(nT)$  would also have stationary increments.

- ▶ We have

$$\begin{aligned}X((m+n+k)T) - X((n+k)T) &= Z_{n+k+1} + \cdots + Z_{n+k+m} \\X((m+n)T) - X(nT) &= Z_{n+1} + \cdots + Z_{n+m}\end{aligned}$$

Both are sums of  $m$  of the iid  $Z_i$ 's

- ▶ Hence both would have the same distribution
- ▶ Thus  $X(nT)$  would also have stationary increments.
- ▶ Hence we expect  $W(t)$  to have stationary increments

- We have

$$\begin{aligned}X((m+n+k)T) - X((n+k)T) &= Z_{n+k+1} + \cdots + Z_{n+k+m} \\X((m+n)T) - X(nT) &= Z_{n+1} + \cdots + Z_{n+m}\end{aligned}$$

Both are sums of  $m$  of the iid  $Z_i$ 's

- Hence both would have the same distribution
- Thus  $X(nT)$  would also have stationary increments.
- Hence we expect  $W(t)$  to have stationary increments
- Thus,  $W(t)$  should be a process with stationary and independent increments and for each  $t$ ,  $W(t)$  is Gaussian with zero mean and variance proportional to  $t$



- ▶ We have

$$\begin{aligned}X((m+n+k)T) - X((n+k)T) &= Z_{n+k+1} + \cdots + Z_{n+k+m} \\X((m+n)T) - X(nT) &= Z_{n+1} + \cdots + Z_{n+m}\end{aligned}$$

Both are sums of  $m$  of the iid  $Z_i$ 's

- ▶ Hence both would have the same distribution
- ▶ Thus  $X(nT)$  would also have stationary increments.
- ▶ Hence we expect  $W(t)$  to have stationary increments
- ▶ Thus,  $W(t)$  should be a process with stationary and independent increments and for each  $t$ ,  $W(t)$  is Gaussian with zero mean and variance proportional to  $t$
- ▶ We will now formally define Brownian motion using these properties.

- ▶ Let  $\{X(t), t \geq 0\}$  be a continuous-state continuous-time process

- ▶ Let  $\{X(t), t \geq 0\}$  be a continuous-state continuous-time process

This process is called a Brownian motion if

- ▶ Let  $\{X(t), t \geq 0\}$  be a continuous-state continuous-time process

This process is called a Brownian motion if

1.  $X(0) = 0$

- Let  $\{X(t), t \geq 0\}$  be a continuous-state continuous-time process

This process is called a Brownian motion if

1.  $X(0) = 0$
2. The process has stationary and independent increments

- Let  $\{X(t), t \geq 0\}$  be a continuous-state continuous-time process

This process is called a Brownian motion if

1.  $X(0) = 0$
2. The process has stationary and independent increments
3. For every  $t > 0$ ,  $X(t)$  is Gaussian with mean 0 and variance  $\sigma^2 t$

- ▶ Let  $\{X(t), t \geq 0\}$  be a continuous-state continuous-time process

This process is called a Brownian motion if

1.  $X(0) = 0$
  2. The process has stationary and independent increments
  3. For every  $t > 0$ ,  $X(t)$  is Gaussian with mean 0 and variance  $\sigma^2 t$
- ▶ Let  $B(t) = \frac{X(t)}{\sigma}$ . Then, variance of  $B(t)$  is  $t$

- ▶ Let  $\{X(t), t \geq 0\}$  be a continuous-state continuous-time process

This process is called a Brownian motion if

1.  $X(0) = 0$
  2. The process has stationary and independent increments
  3. For every  $t > 0$ ,  $X(t)$  is Gaussian with mean 0 and variance  $\sigma^2 t$
- ▶ Let  $B(t) = \frac{X(t)}{\sigma}$ . Then, variance of  $B(t)$  is  $t$
  - ▶  $\{B(t), t \geq 0\}$  is called standard Brownian Motion



- ▶ Let  $\{X(t), t \geq 0\}$  be a continuous-state continuous-time process

This process is called a Brownian motion if

1.  $X(0) = 0$
  2. The process has stationary and independent increments
  3. For every  $t > 0$ ,  $X(t)$  is Gaussian with mean 0 and variance  $\sigma^2 t$
- ▶ Let  $B(t) = \frac{X(t)}{\sigma}$ . Then, variance of  $B(t)$  is  $t$
  - ▶  $\{B(t), t \geq 0\}$  is called standard Brownian Motion
  - ▶ Let  $Y(t) = X(t) + \mu t$ . Then  $Y(t)$  has non-zero mean

- ▶ Let  $\{X(t), t \geq 0\}$  be a continuous-state continuous-time process

This process is called a Brownian motion if

1.  $X(0) = 0$
  2. The process has stationary and independent increments
  3. For every  $t > 0$ ,  $X(t)$  is Gaussian with mean 0 and variance  $\sigma^2 t$
- ▶ Let  $B(t) = \frac{X(t)}{\sigma}$ . Then, variance of  $B(t)$  is  $t$
  - ▶  $\{B(t), t \geq 0\}$  is called standard Brownian Motion
  - ▶ Let  $Y(t) = X(t) + \mu t$ . Then  $Y(t)$  has non-zero mean
  - ▶  $\{Y(t), t \geq 0\}$  is called Brownian motion with a drift

- ▶ Let  $\{X(t), t \geq 0\}$  be a continuous-state continuous-time process

This process is called a Brownian motion if

1.  $X(0) = 0$
2. The process has stationary and independent increments
3. For every  $t > 0$ ,  $X(t)$  is Gaussian with mean 0 and variance  $\sigma^2 t$

- ▶ Let  $B(t) = \frac{X(t)}{\sigma}$ . Then, variance of  $B(t)$  is  $t$
- ▶  $\{B(t), t \geq 0\}$  is called standard Brownian Motion
- ▶ Let  $Y(t) = X(t) + \mu t$ . Then  $Y(t)$  has non-zero mean
- ▶  $\{Y(t), t \geq 0\}$  is called Brownian motion with a drift
- ▶ The mean can be a function of time