

Recap

- ▶ X_1, \dots, X_n are continuous rv with joint density

Recap

- ▶ X_1, \dots, X_n are continuous rv with joint density

$$Y_1 = g_1(X_1, \dots, X_n) \quad \dots \quad Y_n = g_n(X_1, \dots, X_n)$$

Recap

- ▶ X_1, \dots, X_n are continuous rv with joint density

$$Y_1 = g_1(X_1, \dots, X_n) \quad \dots \quad Y_n = g_n(X_1, \dots, X_n)$$

- ▶ The transformation is continuous with continuous first partials and is invertible and

$$X_1 = h_1(Y_1, \dots, Y_n) \quad \dots \quad X_n = h_n(Y_1, \dots, Y_n)$$

Recap

- ▶ X_1, \dots, X_n are continuous rv with joint density

$$Y_1 = g_1(X_1, \dots, X_n) \quad \dots \quad Y_n = g_n(X_1, \dots, X_n)$$

- ▶ The transformation is continuous with continuous first partials and is invertible and

$$X_1 = h_1(Y_1, \dots, Y_n) \quad \dots \quad X_n = h_n(Y_1, \dots, Y_n)$$

- ▶ The Jacobian of the inverse transform, J , is non-zero

Recap

- ▶ X_1, \dots, X_n are continuous rv with joint density

$$Y_1 = g_1(X_1, \dots, X_n) \quad \dots \quad Y_n = g_n(X_1, \dots, X_n)$$

- ▶ The transformation is continuous with continuous first partials and is invertible and

$$X_1 = h_1(Y_1, \dots, Y_n) \quad \dots \quad X_n = h_n(Y_1, \dots, Y_n)$$

- ▶ The Jacobian of the inverse transform, J , is non-zero
- ▶ Then the density of \mathbf{Y} is

$$f_{Y_1 \dots Y_n}(y_1, \dots, y_n) = |J| f_{X_1 \dots X_n}(h_1(y_1, \dots, y_n), \dots, h_n(y_1, \dots, y_n))$$

Recap

- ▶ X_1, \dots, X_n are continuous rv with joint density

$$Y_1 = g_1(X_1, \dots, X_n) \quad \dots \quad Y_n = g_n(X_1, \dots, X_n)$$

- ▶ The transformation is continuous with continuous first partials and is invertible and

$$X_1 = h_1(Y_1, \dots, Y_n) \quad \dots \quad X_n = h_n(Y_1, \dots, Y_n)$$

- ▶ The Jacobian of the inverse transform, J , is non-zero
- ▶ Then the density of \mathbf{Y} is

$$f_{Y_1 \dots Y_n}(y_1, \dots, y_n) = |J| f_{X_1 \dots X_n}(h_1(y_1, \dots, y_n), \dots, h_n(y_1, \dots, y_n))$$

- ▶ Called multidimensional change of variable formula

Recap: Densities of some standard functions of rv's

- ▶ One can use the theorem to find densities of sum, difference, product and quotient of random variables.

Recap: Densities of some standard functions of rv's

- ▶ One can use the theorem to find densities of sum, difference, product and quotient of random variables.

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt = \int_{-\infty}^{\infty} f_{XY}(z-t, t) dt$$

Recap: Densities of some standard functions of rv's

- ▶ One can use the theorem to find densities of sum, difference, product and quotient of random variables.

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt = \int_{-\infty}^{\infty} f_{XY}(z-t, t) dt$$

$$f_{X-Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, t-z) dt = \int_{-\infty}^{\infty} f_{XY}(t+z, t) dt$$

Recap: Densities of some standard functions of rv's

- ▶ One can use the theorem to find densities of sum, difference, product and quotient of random variables.

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt = \int_{-\infty}^{\infty} f_{XY}(z-t, t) dt$$

$$f_{X-Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, t-z) dt = \int_{-\infty}^{\infty} f_{XY}(t+z, t) dt$$

$$f_{X*Y}(z) = \int_{-\infty}^{\infty} \left| \frac{1}{t} \right| f_{XY} \left(\frac{z}{t}, t \right) dt = \int_{-\infty}^{\infty} \left| \frac{1}{t} \right| f_{XY} \left(t, \frac{z}{t} \right) dt$$

Recap: Densities of some standard functions of rv's

- One can use the theorem to find densities of sum, difference, product and quotient of random variables.

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt = \int_{-\infty}^{\infty} f_{XY}(z-t, t) dt$$

$$f_{X-Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, t-z) dt = \int_{-\infty}^{\infty} f_{XY}(t+z, t) dt$$

$$f_{X*Y}(z) = \int_{-\infty}^{\infty} \left| \frac{1}{t} \right| f_{XY} \left(\frac{z}{t}, t \right) dt = \int_{-\infty}^{\infty} \left| \frac{1}{t} \right| f_{XY} \left(t, \frac{z}{t} \right) dt$$

$$f_{(X/Y)}(z) = \int_{-\infty}^{\infty} |t| f_{XY}(zt, t) dt = \int_{-\infty}^{\infty} \left| \frac{t}{z^2} \right| f_{XY} \left(t, \frac{t}{z} \right) dt$$

Density of XY

- ▶ Let X, Y have joint density f_{XY} .

Density of XY

- ▶ Let X, Y have joint density f_{XY} .
- ▶ Let $Z = XY$. We want to find density of XY directly

Density of XY

- ▶ Let X, Y have joint density f_{XY} .
- ▶ Let $Z = XY$. We want to find density of XY directly
- ▶ Let $A_z = \{(x, y) \in \mathbb{R}^2 : xy \leq z\} \subset \mathbb{R}^2$.

Density of XY

- ▶ Let X, Y have joint density f_{XY} .
- ▶ Let $Z = XY$. We want to find density of XY directly
- ▶ Let $A_z = \{(x, y) \in \mathbb{R}^2 : xy \leq z\} \subset \mathbb{R}^2$.

$$F_Z(z) = P[XY \leq z] = P[(X, Y) \in A_z]$$

Density of XY

- ▶ Let X, Y have joint density f_{XY} .
- ▶ Let $Z = XY$. We want to find density of XY directly
- ▶ Let $A_z = \{(x, y) \in \mathbb{R}^2 : xy \leq z\} \subset \mathbb{R}^2$.

$$\begin{aligned} F_Z(z) &= P[XY \leq z] = P[(X, Y) \in A_z] \\ &= \int \int_{A_z} f_{XY}(x, y) \, dy \, dx \end{aligned}$$

Density of XY

- ▶ Let X, Y have joint density f_{XY} .
- ▶ Let $Z = XY$. We want to find density of XY directly
- ▶ Let $A_z = \{(x, y) \in \mathbb{R}^2 : xy \leq z\} \subset \mathbb{R}^2$.

$$\begin{aligned} F_Z(z) &= P[XY \leq z] = P[(X, Y) \in A_z] \\ &= \int \int_{A_z} f_{XY}(x, y) \, dy \, dx \end{aligned}$$

- ▶ We need to find limits for integrating over A_z

Density of XY

- ▶ Let X, Y have joint density f_{XY} .
- ▶ Let $Z = XY$. We want to find density of XY directly
- ▶ Let $A_z = \{(x, y) \in \mathbb{R}^2 : xy \leq z\} \subset \mathbb{R}^2$.

$$\begin{aligned} F_Z(z) &= P[XY \leq z] = P[(X, Y) \in A_z] \\ &= \int \int_{A_z} f_{XY}(x, y) \, dy \, dx \end{aligned}$$

- ▶ We need to find limits for integrating over A_z
- ▶ If $x > 0$, then $xy \leq z \Rightarrow y \leq z/x$
If $x < 0$, then $xy \leq z \Rightarrow y \geq z/x$

Density of XY

- ▶ Let X, Y have joint density f_{XY} .
- ▶ Let $Z = XY$. We want to find density of XY directly
- ▶ Let $A_z = \{(x, y) \in \mathbb{R}^2 : xy \leq z\} \subset \mathbb{R}^2$.

$$\begin{aligned} F_Z(z) &= P[XY \leq z] = P[(X, Y) \in A_z] \\ &= \int \int_{A_z} f_{XY}(x, y) \, dy \, dx \end{aligned}$$

- ▶ We need to find limits for integrating over A_z
- ▶ If $x > 0$, then $xy \leq z \Rightarrow y \leq z/x$
If $x < 0$, then $xy \leq z \Rightarrow y \geq z/x$

$$F_Z(z) = \int_{-\infty}^0 \int_{z/x}^{\infty} f_{XY}(x, y) \, dy \, dx + \int_0^{\infty} \int_{-\infty}^{z/x} f_{XY}(x, y) \, dy \, dx$$

$$F_Z(z) = \int_{-\infty}^0 \int_{z/x}^{\infty} f_{XY}(x, y) \, dy \, dx + \int_0^{\infty} \int_{-\infty}^{z/x} f_{XY}(x, y) \, dy \, dx$$

- Change variable from y to t using $t = xy$
 $y = t/x; \quad dy = \frac{1}{x} dt; \quad y = z/x \Rightarrow t = z$

$$F_Z(z) = \int_{-\infty}^0 \int_{z/x}^{\infty} f_{XY}(x, y) \, dy \, dx + \int_0^{\infty} \int_{-\infty}^{z/x} f_{XY}(x, y) \, dy \, dx$$

- Change variable from y to t using $t = xy$
 $y = t/x; \, dy = \frac{1}{x} \, dt; \, y = z/x \Rightarrow t = z$

$$F_Z(z) = \int_{-\infty}^0 \int_z^{-\infty} \frac{1}{x} f_{XY}\left(x, \frac{t}{x}\right) \, dt \, dx + \int_0^{\infty} \int_{-\infty}^z \frac{1}{x} f_{XY}\left(x, \frac{t}{x}\right) \, dt \, dx$$

$$F_Z(z) = \int_{-\infty}^0 \int_{z/x}^{\infty} f_{XY}(x, y) dy dx + \int_0^{\infty} \int_{-\infty}^{z/x} f_{XY}(x, y) dy dx$$

- Change variable from y to t using $t = xy$
 $y = t/x$; $dy = \frac{1}{x} dt$; $y = z/x \Rightarrow t = z$

$$\begin{aligned} F_Z(z) &= \int_{-\infty}^0 \int_z^{-\infty} \frac{1}{x} f_{XY}\left(x, \frac{t}{x}\right) dt dx + \int_0^{\infty} \int_{-\infty}^z \frac{1}{x} f_{XY}\left(x, \frac{t}{x}\right) dt dx \\ &= \int_{-\infty}^0 \int_{-\infty}^z \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dt dx + \int_0^{\infty} \int_{-\infty}^z \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dt dx \end{aligned}$$

$$F_Z(z) = \int_{-\infty}^0 \int_{z/x}^{\infty} f_{XY}(x, y) dy dx + \int_0^{\infty} \int_{-\infty}^{z/x} f_{XY}(x, y) dy dx$$

- Change variable from y to t using $t = xy$
 $y = t/x$; $dy = \frac{1}{x} dt$; $y = z/x \Rightarrow t = z$

$$\begin{aligned} F_Z(z) &= \int_{-\infty}^0 \int_z^{-\infty} \frac{1}{x} f_{XY}\left(x, \frac{t}{x}\right) dt dx + \int_0^{\infty} \int_{-\infty}^z \frac{1}{x} f_{XY}\left(x, \frac{t}{x}\right) dt dx \\ &= \int_{-\infty}^0 \int_{-\infty}^z \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dt dx + \int_0^{\infty} \int_{-\infty}^z \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dt dx \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^z \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dt dx \end{aligned}$$

$$F_Z(z) = \int_{-\infty}^0 \int_{z/x}^{\infty} f_{XY}(x, y) dy dx + \int_0^{\infty} \int_{-\infty}^{z/x} f_{XY}(x, y) dy dx$$

- Change variable from y to t using $t = xy$
 $y = t/x$; $dy = \frac{1}{x} dt$; $y = z/x \Rightarrow t = z$

$$\begin{aligned} F_Z(z) &= \int_{-\infty}^0 \int_z^{-\infty} \frac{1}{x} f_{XY}\left(x, \frac{t}{x}\right) dt dx + \int_0^{\infty} \int_{-\infty}^z \frac{1}{x} f_{XY}\left(x, \frac{t}{x}\right) dt dx \\ &= \int_{-\infty}^0 \int_{-\infty}^z \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dt dx + \int_0^{\infty} \int_{-\infty}^z \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dt dx \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^z \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dt dx \\ &= \int_{-\infty}^z \int_{-\infty}^{\infty} \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dx dt \end{aligned}$$

$$F_Z(z) = \int_{-\infty}^0 \int_{z/x}^{\infty} f_{XY}(x, y) dy dx + \int_0^{\infty} \int_{-\infty}^{z/x} f_{XY}(x, y) dy dx$$

- Change variable from y to t using $t = xy$
 $y = t/x$; $dy = \frac{1}{x} dt$; $y = z/x \Rightarrow t = z$

$$\begin{aligned} F_Z(z) &= \int_{-\infty}^0 \int_z^{-\infty} \frac{1}{x} f_{XY}\left(x, \frac{t}{x}\right) dt dx + \int_0^{\infty} \int_{-\infty}^z \frac{1}{x} f_{XY}\left(x, \frac{t}{x}\right) dt dx \\ &= \int_{-\infty}^0 \int_{-\infty}^z \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dt dx + \int_0^{\infty} \int_{-\infty}^z \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dt dx \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^z \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dt dx \\ &= \int_{-\infty}^z \int_{-\infty}^{\infty} \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{t}{x}\right) dx dt \end{aligned}$$

This shows: $f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{1}{x} \right| f_{XY}\left(x, \frac{z}{x}\right) dx$

Recap: Exchangeable random variables

- ▶ X_1, X_2, \dots, X_n are said to be exchangeable if their joint distribution is same as that of any permutation of them.

Recap: Exchangeable random variables

- ▶ X_1, X_2, \dots, X_n are said to be exchangeable if their joint distribution is same as that of any permutation of them.
- ▶ If the random variables are exchangeable then the joint distribution function remains the same on permutation of arguments.

Recap: Exchangeable random variables

- ▶ X_1, X_2, \dots, X_n are said to be exchangeable if their joint distribution is same as that of any permutation of them.
- ▶ If the random variables are exchangeable then the joint distribution function remains the same on permutation of arguments.
- ▶ Exchangeable random variables are identically distributed but they may not be independent.

Recap: Expectation of function of multiple rv's

- ▶ Let $Z = g(X_1, \dots, X_n) = g(\mathbf{X})$. Then

$$E[Z] = \int_{\mathbb{R}^n} g(\mathbf{x}) dF_{\mathbf{X}}(\mathbf{x})$$

Recap: Expectation of function of multiple rv's

- ▶ Let $Z = g(X_1, \dots, X_n) = g(\mathbf{X})$. Then

$$E[Z] = \int_{\mathbb{R}^n} g(\mathbf{x}) dF_{\mathbf{X}}(\mathbf{x})$$

- ▶ For example, if they have a joint density, then

$$E[Z] = \int_{\mathbb{R}^n} g(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\mathbf{x}$$

Recap: Expectation of function of multiple rv's

- ▶ Let $Z = g(X_1, \dots, X_n) = g(\mathbf{X})$. Then

$$E[Z] = \int_{\mathbb{R}^n} g(\mathbf{x}) dF_{\mathbf{X}}(\mathbf{x})$$

- ▶ For example, if they have a joint density, then

$$E[Z] = \int_{\mathbb{R}^n} g(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\mathbf{x}$$

- ▶ Implies, $E[g_1(\mathbf{X}) + g_2(\mathbf{X})] = E[g_1(\mathbf{X})] + E[g_2(\mathbf{X})]$

Recap: Expectation of function of multiple rv's

- ▶ Let $Z = g(X_1, \dots, X_n) = g(\mathbf{X})$. Then

$$E[Z] = \int_{\mathbb{R}^n} g(\mathbf{x}) dF_{\mathbf{X}}(\mathbf{x})$$

- ▶ For example, if they have a joint density, then

$$E[Z] = \int_{\mathbb{R}^n} g(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\mathbf{x}$$

- ▶ Implies, $E[g_1(\mathbf{X}) + g_2(\mathbf{X})] = E[g_1(\mathbf{X})] + E[g_2(\mathbf{X})]$
- ▶ Specifically, $E[X + Y] = E[X] + E[Y]$

Recap: Covariance

- ▶ The covariance of X, Y is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Recap: Covariance

- ▶ The covariance of X, Y is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Note that $\text{Cov}(X, X) = \text{Var}(X)$

Recap: Covariance

- ▶ The covariance of X, Y is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Note that $\text{Cov}(X, X) = \text{Var}(X)$

- ▶ $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y) + 2\text{Cov}(X, Y)$

Recap: Covariance

- ▶ The covariance of X, Y is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Note that $\text{Cov}(X, X) = \text{Var}(X)$

- ▶ $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y) + 2\text{Cov}(X, Y)$
- ▶ X, Y are called uncorrelated if $\text{Cov}(X, Y) = 0$.

Recap: Covariance

- ▶ The covariance of X, Y is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Note that $\text{Cov}(X, X) = \text{Var}(X)$

- ▶ $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y) + 2\text{Cov}(X, Y)$
- ▶ X, Y are called uncorrelated if $\text{Cov}(X, Y) = 0$.
- ▶ If X, Y are uncorrelated, $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y)$

Recap: Covariance

- ▶ The covariance of X, Y is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Note that $\text{Cov}(X, X) = \text{Var}(X)$

- ▶ $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y) + 2\text{Cov}(X, Y)$
- ▶ X, Y are called uncorrelated if $\text{Cov}(X, Y) = 0$.
- ▶ If X, Y are uncorrelated, $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y)$
- ▶ X, Y independent $\Rightarrow X, Y$ uncorrelated.

Recap: Covariance

- ▶ The covariance of X, Y is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Note that $\text{Cov}(X, X) = \text{Var}(X)$

- ▶ $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y) + 2\text{Cov}(X, Y)$
- ▶ X, Y are called uncorrelated if $\text{Cov}(X, Y) = 0$.
- ▶ If X, Y are uncorrelated, $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y)$
- ▶ X, Y independent $\Rightarrow X, Y$ uncorrelated.
- ▶ Uncorrelated random variables need not necessarily be independent

Recap: Correlation coefficient

- ▶ The correlation coefficient of X, Y is

$$\rho_{XY} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

Recap: Correlation coefficient

- ▶ The correlation coefficient of X, Y is

$$\rho_{XY} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

- ▶ If X, Y are uncorrelated then $\rho_{XY} = 0$.

Recap: Correlation coefficient

- ▶ The correlation coefficient of X, Y is

$$\rho_{XY} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

- ▶ If X, Y are uncorrelated then $\rho_{XY} = 0$.
- ▶ $-1 \leq \rho_{XY} \leq 1, \forall X, Y$

Recap: Correlation coefficient

- ▶ The correlation coefficient of X, Y is

$$\rho_{XY} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

- ▶ If X, Y are uncorrelated then $\rho_{XY} = 0$.
- ▶ $-1 \leq \rho_{XY} \leq 1, \forall X, Y$
- ▶ $|\rho_{XY}| = 1$ iff $X = aY$

Recap: mean square estimation

- ▶ The best mean-square approximation of Y as a 'linear' function of X is

$$Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)} X + \left(EY - \frac{\text{Cov}(X, Y)}{\text{Var}(X)} EX \right)$$

Recap: mean square estimation

- ▶ The best mean-square approximation of Y as a 'linear' function of X is

$$Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)} X + \left(EY - \frac{\text{Cov}(X, Y)}{\text{Var}(X)} EX \right)$$

- ▶ Called the line of regression of Y on X .

Recap: mean square estimation

- ▶ The best mean-square approximation of Y as a 'linear' function of X is

$$Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)} X + \left(EY - \frac{\text{Cov}(X, Y)}{\text{Var}(X)} EX \right)$$

- ▶ Called the line of regression of Y on X .
- ▶ If $\text{cov}(X, Y) = 0$ then this reduces to approximating Y by a constant, EY .

Recap: mean square estimation

- ▶ The best mean-square approximation of Y as a 'linear' function of X is

$$Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)} X + \left(EY - \frac{\text{Cov}(X, Y)}{\text{Var}(X)} EX \right)$$

- ▶ Called the line of regression of Y on X .
- ▶ If $\text{cov}(X, Y) = 0$ then this reduces to approximating Y by a constant, EY .
- ▶ The final mean square error is

$$\text{Var}(Y) (1 - \rho_{XY}^2)$$

Recap: mean square estimation

- ▶ The best mean-square approximation of Y as a 'linear' function of X is

$$Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)} X + \left(EY - \frac{\text{Cov}(X, Y)}{\text{Var}(X)} EX \right)$$

- ▶ Called the line of regression of Y on X .
- ▶ If $\text{cov}(X, Y) = 0$ then this reduces to approximating Y by a constant, EY .
- ▶ The final mean square error is

$$\text{Var}(Y) (1 - \rho_{XY}^2)$$

- ▶ If $\rho_{XY} = \pm 1$ then the error is zero

Recap: mean square estimation

- ▶ The best mean-square approximation of Y as a 'linear' function of X is

$$Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)} X + \left(EY - \frac{\text{Cov}(X, Y)}{\text{Var}(X)} EX \right)$$

- ▶ Called the line of regression of Y on X .
- ▶ If $\text{cov}(X, Y) = 0$ then this reduces to approximating Y by a constant, EY .
- ▶ The final mean square error is

$$\text{Var}(Y) (1 - \rho_{XY}^2)$$

- ▶ If $\rho_{XY} = \pm 1$ then the error is zero
- ▶ If $\rho_{XY} = 0$ the final error is $\text{Var}(Y)$

Recap: Covariance matrix

- For a random vector, $\mathbf{X} = (X_1, \dots, X_n)^T$, the covariance matrix is

$$\Sigma_{\mathbf{X}} = E [(\mathbf{X} - E\mathbf{X})(\mathbf{X} - E\mathbf{X})^T]$$

Recap: Covariance matrix

- For a random vector, $\mathbf{X} = (X_1, \dots, X_n)^T$, the covariance matrix is

$$\Sigma_{\mathbf{X}} = E [(\mathbf{X} - E\mathbf{X})(\mathbf{X} - E\mathbf{X})^T]$$

$$(\Sigma_{\mathbf{X}})_{ij} = E[(X_i - EX_i)(X_j - EX_j)]$$

Recap: Covariance matrix

- ▶ For a random vector, $\mathbf{X} = (X_1, \dots, X_n)^T$, the covariance matrix is

$$\Sigma_{\mathbf{X}} = E [(\mathbf{X} - E\mathbf{X})(\mathbf{X} - E\mathbf{X})^T]$$

$$(\Sigma_{\mathbf{X}})_{ij} = E[(X_i - EX_i)(X_j - EX_j)]$$

- ▶ $\text{Var}(\mathbf{a}^T \mathbf{X}) = \mathbf{a}^T \Sigma_{\mathbf{X}} \mathbf{a}$

Recap: Covariance matrix

- ▶ For a random vector, $\mathbf{X} = (X_1, \dots, X_n)^T$, the covariance matrix is

$$\Sigma_{\mathbf{X}} = E [(\mathbf{X} - E\mathbf{X})(\mathbf{X} - E\mathbf{X})^T]$$

$$(\Sigma_{\mathbf{X}})_{ij} = E[(X_i - EX_i)(X_j - EX_j)]$$

- ▶ $\text{Var}(\mathbf{a}^T \mathbf{X}) = \mathbf{a}^T \Sigma_X \mathbf{a}$
- ▶ Σ_X is a real symmetric and positive semidefinite matrix.

Joint moments

Joint moments

- ▶ Given two random variables, X, Y

Joint moments

- ▶ Given two random variables, X, Y
- ▶ The joint moment of order (i, j) is defined by

$$m_{ij} = E[X^i Y^j]$$

Joint moments

- ▶ Given two random variables, X, Y
- ▶ The joint moment of order (i, j) is defined by

$$m_{ij} = E[X^i Y^j]$$

$m_{10} = EX$, $m_{01} = EY$, $m_{11} = E[XY]$ and so on

Joint moments

- ▶ Given two random variables, X, Y
- ▶ The joint moment of order (i, j) is defined by

$$m_{ij} = E[X^i Y^j]$$

$m_{10} = EX$, $m_{01} = EY$, $m_{11} = E[XY]$ and so on

- ▶ Similarly joint central moments of order (i, j) are defined by

$$s_{ij} = E[(X - EX)^i (Y - EY)^j]$$

Joint moments

- ▶ Given two random variables, X, Y
- ▶ The joint moment of order (i, j) is defined by

$$m_{ij} = E[X^i Y^j]$$

$m_{10} = EX$, $m_{01} = EY$, $m_{11} = E[XY]$ and so on

- ▶ Similarly joint central moments of order (i, j) are defined by

$$s_{ij} = E[(X - EX)^i (Y - EY)^j]$$

$s_{10} = s_{01} = 0$, $s_{11} = \text{Cov}(X, Y)$, $s_{20} = \text{Var}(X)$ and so on

Joint moments

- ▶ Given two random variables, X, Y
- ▶ The joint moment of order (i, j) is defined by

$$m_{ij} = E[X^i Y^j]$$

$m_{10} = EX$, $m_{01} = EY$, $m_{11} = E[XY]$ and so on

- ▶ Similarly joint central moments of order (i, j) are defined by

$$s_{ij} = E[(X - EX)^i (Y - EY)^j]$$

$s_{10} = s_{01} = 0$, $s_{11} = \text{Cov}(X, Y)$, $s_{20} = \text{Var}(X)$ and so on

- ▶ We can similarly define joint moments of multiple random variables

- ▶ We can define moment generating function of X, Y by

$$M_{XY}(s, t) = E \left[e^{sX+tY} \right], \quad s, t \in \Re$$

- ▶ We can define moment generating function of X, Y by

$$M_{XY}(s, t) = E \left[e^{sX+tY} \right], \quad s, t \in \Re$$

- ▶ This is easily generalized to n random variables

$$M_{\mathbf{X}}(\mathbf{s}) = E \left[e^{\mathbf{s}^T \mathbf{X}} \right], \quad \mathbf{s} \in \Re^n$$

- ▶ We can define moment generating function of X, Y by

$$M_{XY}(s, t) = E \left[e^{sX+tY} \right], \quad s, t \in \Re$$

- ▶ This is easily generalized to n random variables

$$M_{\mathbf{X}}(\mathbf{s}) = E \left[e^{\mathbf{s}^T \mathbf{X}} \right], \quad \mathbf{s} \in \Re^n$$

- ▶ Once again, we can get all the moments by differentiating the moment generating function

$$\left. \frac{\partial}{\partial s_i} M_{\mathbf{X}}(\mathbf{s}) \right|_{\mathbf{s}=\mathbf{0}} = EX_i$$

- ▶ We can define moment generating function of X, Y by

$$M_{XY}(s, t) = E \left[e^{sX+tY} \right], \quad s, t \in \mathbb{R}$$

- ▶ This is easily generalized to n random variables

$$M_{\mathbf{X}}(\mathbf{s}) = E \left[e^{\mathbf{s}^T \mathbf{X}} \right], \quad \mathbf{s} \in \mathbb{R}^n$$

- ▶ Once again, we can get all the moments by differentiating the moment generating function

$$\left. \frac{\partial}{\partial s_i} M_{\mathbf{X}}(\mathbf{s}) \right|_{\mathbf{s}=0} = EX_i$$

- ▶ More generally

$$\left. \frac{\partial^{m+n}}{\partial s_i^n \partial s_j^m} M_{\mathbf{X}}(\mathbf{s}) \right|_{\mathbf{s}=0} = EX_i^n X_j^m$$

- We define the characteristic function of \mathbf{X} by

$$\phi_{\mathbf{X}}(\mathbf{s}) = E \left[e^{i\mathbf{s}^T \mathbf{X}} \right], \quad \mathbf{s} \in \Re^n, \quad i = \sqrt{-1}$$

- ▶ We define the characteristic function of \mathbf{X} by

$$\phi_{\mathbf{X}}(\mathbf{s}) = E \left[e^{i\mathbf{s}^T \mathbf{X}} \right], \quad \mathbf{s} \in \Re^n, \quad i = \sqrt{-1}$$

- ▶ Characteristic function exists for all random vectors and it uniquely determines the joint distribution.

- ▶ We define the characteristic function of \mathbf{X} by

$$\phi_{\mathbf{X}}(\mathbf{s}) = E \left[e^{i\mathbf{s}^T \mathbf{X}} \right], \quad \mathbf{s} \in \Re^n, \quad i = \sqrt{-1}$$

- ▶ Characteristic function exists for all random vectors and it uniquely determines the joint distribution.
- ▶ The moment generating function may not exist for all random vectors. When it exists, it uniquely determines the joint distribution.

Conditional Expectation

- ▶ Suppose X, Y have a joint density f_{XY}

Conditional Expectation

- ▶ Suppose X, Y have a joint density f_{XY}
- ▶ Consider the conditional density $f_{X|Y}(x|y)$. This is a density in x for every value of y .

Conditional Expectation

- ▶ Suppose X, Y have a joint density f_{XY}
- ▶ Consider the conditional density $f_{X|Y}(x|y)$. This is a density in x for every value of y .
- ▶ Since it is a density, we can use it in an expectation integral: $\int g(x) f_{X|Y}(x|y) dx$

Conditional Expectation

- ▶ Suppose X, Y have a joint density f_{XY}
- ▶ Consider the conditional density $f_{X|Y}(x|y)$. This is a density in x for every value of y .
- ▶ Since it is a density, we can use it in an expectation integral: $\int g(x) f_{X|Y}(x|y) dx$
- ▶ This is like expectation of $g(X)$ since $f_{X|Y}(x|y)$ is a density in x .

Conditional Expectation

- ▶ Suppose X, Y have a joint density f_{XY}
- ▶ Consider the conditional density $f_{X|Y}(x|y)$. This is a density in x for every value of y .
- ▶ Since it is a density, we can use it in an expectation integral: $\int g(x) f_{X|Y}(x|y) dx$
- ▶ This is like expectation of $g(X)$ since $f_{X|Y}(x|y)$ is a density in x .
- ▶ However, its value would be a function of y .

Conditional Expectation

- ▶ Suppose X, Y have a joint density f_{XY}
- ▶ Consider the conditional density $f_{X|Y}(x|y)$. This is a density in x for every value of y .
- ▶ Since it is a density, we can use it in an expectation integral: $\int g(x) f_{X|Y}(x|y) dx$
- ▶ This is like expectation of $g(X)$ since $f_{X|Y}(x|y)$ is a density in x .
- ▶ However, its value would be a function of y .
- ▶ That is, this is a kind of expectation that is a function of Y (and hence is a random variable)

Conditional Expectation

- ▶ Suppose X, Y have a joint density f_{XY}
- ▶ Consider the conditional density $f_{X|Y}(x|y)$. This is a density in x for every value of y .
- ▶ Since it is a density, we can use it in an expectation integral: $\int g(x) f_{X|Y}(x|y) dx$
- ▶ This is like expectation of $g(X)$ since $f_{X|Y}(x|y)$ is a density in x .
- ▶ However, its value would be a function of y .
- ▶ That is, this is a kind of expectation that is a function of Y (and hence is a random variable)
- ▶ It is called conditional expectation.

- ▶ Let X, Y be discrete random variables (on the same probability space).

- ▶ Let X, Y be discrete random variables (on the same probability space).
- ▶ The conditional expectation of $h(X)$ conditioned on Y is a function of Y , and its value for any y is defined by

- ▶ Let X, Y be discrete random variables (on the same probability space).
- ▶ The conditional expectation of $h(X)$ conditioned on Y is a function of Y , and its value for any y is defined by

$$E[h(X)|Y = y] = \sum_x h(x) f_{X|Y}(x|y)$$

- ▶ Let X, Y be discrete random variables (on the same probability space).
- ▶ The conditional expectation of $h(X)$ conditioned on Y is a function of Y , and its value for any y is defined by

$$\begin{aligned} E[h(X)|Y = y] &= \sum_x h(x) f_{X|Y}(x|y) \\ &= \sum_x h(x) P[X = x|Y = y] \end{aligned}$$

- ▶ Let X, Y be discrete random variables (on the same probability space).
- ▶ The conditional expectation of $h(X)$ conditioned on Y is a function of Y , and its value for any y is defined by

$$\begin{aligned} E[h(X)|Y = y] &= \sum_x h(x) f_{X|Y}(x|y) \\ &= \sum_x h(x) P[X = x|Y = y] \end{aligned}$$

- ▶ That is, we define $E[h(X)|Y] = g(Y)$ where

$$g(y) = \sum_x h(x) f_{X|Y}(x|y)$$

- ▶ Let X, Y be discrete random variables (on the same probability space).
- ▶ The conditional expectation of $h(X)$ conditioned on Y is a function of Y , and its value for any y is defined by

$$\begin{aligned} E[h(X)|Y = y] &= \sum_x h(x) f_{X|Y}(x|y) \\ &= \sum_x h(x) P[X = x|Y = y] \end{aligned}$$

- ▶ That is, we define $E[h(X)|Y] = g(Y)$ where

$$E[h(X)|Y = y] = g(y) = \sum_x h(x) f_{X|Y}(x|y)$$

- ▶ Let X, Y be discrete random variables (on the same probability space).
- ▶ The conditional expectation of $h(X)$ conditioned on Y is a function of Y , and its value for any y is defined by

$$\begin{aligned} E[h(X)|Y = y] &= \sum_x h(x) f_{X|Y}(x|y) \\ &= \sum_x h(x) P[X = x|Y = y] \end{aligned}$$

- ▶ That is, we define $E[h(X)|Y] = g(Y)$ where

$$E[h(X)|Y = y] = g(y) = \sum_x h(x) f_{X|Y}(x|y)$$

- ▶ Thus, $E[h(X)|Y]$ is a random variable

- ▶ Let X, Y have joint density f_{XY} .

- ▶ Let X, Y have joint density f_{XY} .
- ▶ The conditional expectation of $h(X)$ conditioned on Y is a function of Y , and its value for any y is defined by

$$E[h(X)|Y = y] = \int_{-\infty}^{\infty} h(x) f_{X|Y}(x|y) dx$$

- ▶ Let X, Y have joint density f_{XY} .
- ▶ The conditional expectation of $h(X)$ conditioned on Y is a function of Y , and its value for any y is defined by

$$E[h(X)|Y = y] = \int_{-\infty}^{\infty} h(x) f_{X|Y}(x|y) dx$$

- ▶ Once again, what this means is that $E[h(X)|Y] = g(Y)$ where

$$g(y) = \int_{-\infty}^{\infty} h(x) f_{X|Y}(x|y) dx$$

A simple example

- Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

A simple example

- Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- We calculated the conditional densities earlier

$$f_{X|Y}(x|y) = \frac{1}{y}, \quad f_{Y|X}(y|x) = \frac{1}{1-x}, \quad 0 < x < y < 1$$

A simple example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We calculated the conditional densities earlier

$$f_{X|Y}(x|y) = \frac{1}{y}, \quad f_{Y|X}(y|x) = \frac{1}{1-x}, \quad 0 < x < y < 1$$

- ▶ Now we can calculate the conditional expectation

A simple example

- Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- We calculated the conditional densities earlier

$$f_{X|Y}(x|y) = \frac{1}{y}, \quad f_{Y|X}(y|x) = \frac{1}{1-x}, \quad 0 < x < y < 1$$

- Now we can calculate the conditional expectation

$$E[X|Y = y] = \int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx$$

A simple example

- Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- We calculated the conditional densities earlier

$$f_{X|Y}(x|y) = \frac{1}{y}, \quad f_{Y|X}(y|x) = \frac{1}{1-x}, \quad 0 < x < y < 1$$

- Now we can calculate the conditional expectation

$$\begin{aligned} E[X|Y = y] &= \int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx \\ &= \int_0^y x \frac{1}{y} dx \end{aligned}$$

A simple example

- Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- We calculated the conditional densities earlier

$$f_{X|Y}(x|y) = \frac{1}{y}, \quad f_{Y|X}(y|x) = \frac{1}{1-x}, \quad 0 < x < y < 1$$

- Now we can calculate the conditional expectation

$$\begin{aligned} E[X|Y = y] &= \int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx \\ &= \int_0^y x \frac{1}{y} dx = \frac{1}{y} \left. \frac{x^2}{2} \right|_0^y = \frac{y}{2} \end{aligned}$$

A simple example

- Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- We calculated the conditional densities earlier

$$f_{X|Y}(x|y) = \frac{1}{y}, \quad f_{Y|X}(y|x) = \frac{1}{1-x}, \quad 0 < x < y < 1$$

- Now we can calculate the conditional expectation

$$\begin{aligned} E[X|Y = y] &= \int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx \\ &= \int_0^y x \frac{1}{y} dx = \frac{1}{y} \left. \frac{x^2}{2} \right|_0^y = \frac{y}{2} \end{aligned}$$

- This gives: $E[X|Y] = \frac{Y}{2}$

A simple example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We calculated the conditional densities earlier

$$f_{X|Y}(x|y) = \frac{1}{y}, \quad f_{Y|X}(y|x) = \frac{1}{1-x}, \quad 0 < x < y < 1$$

- ▶ Now we can calculate the conditional expectation

$$\begin{aligned} E[X|Y = y] &= \int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx \\ &= \int_0^y x \frac{1}{y} dx = \frac{1}{y} \left. \frac{x^2}{2} \right|_0^y = \frac{y}{2} \end{aligned}$$

- ▶ This gives: $E[X|Y] = \frac{Y}{2}$
- ▶ We can show $E[Y|X] =$

A simple example

- Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- We calculated the conditional densities earlier

$$f_{X|Y}(x|y) = \frac{1}{y}, \quad f_{Y|X}(y|x) = \frac{1}{1-x}, \quad 0 < x < y < 1$$

- Now we can calculate the conditional expectation

$$\begin{aligned} E[X|Y = y] &= \int_{-\infty}^{\infty} x f_{X|Y}(x|y) dx \\ &= \int_0^y x \frac{1}{y} dx = \frac{1}{y} \left. \frac{x^2}{2} \right|_0^y = \frac{y}{2} \end{aligned}$$

- This gives: $E[X|Y] = \frac{Y}{2}$
- We can show $E[Y|X] = \frac{1+X}{2}$

- ▶ The conditional expectation is defined by

- The conditional expectation is defined by

$$E[h(X)|Y = y] = \sum_x h(x) f_{X|Y}(x|y), \quad X, Y \text{ are discrete}$$

► The conditional expectation is defined by

$$E[h(X)|Y = y] = \sum_x h(x) f_{X|Y}(x|y), \quad X, Y \text{ are discrete}$$

$$E[h(X)|Y = y] = \int_{-\infty}^{\infty} h(x) f_{X|Y}(x|y) dx, \quad X, Y \text{ have joint density}$$

- ▶ The conditional expectation is defined by

$$E[h(X)|Y = y] = \sum_x h(x) f_{X|Y}(x|y), \quad X, Y \text{ are discrete}$$

$$E[h(X)|Y = y] = \int_{-\infty}^{\infty} h(x); f_{X|Y}(x|y) dx, \quad X, Y \text{ have joint density}$$

- ▶ We can actually define $E[h(X, Y)|Y]$ also as above.
That is,

$$E[h(X, Y)|Y = y] = \int_{-\infty}^{\infty} h(x, y) f_{X|Y}(x|y) dx$$

- ▶ The conditional expectation is defined by

$$E[h(X)|Y = y] = \sum_x h(x) f_{X|Y}(x|y), \quad X, Y \text{ are discrete}$$

$$E[h(X)|Y = y] = \int_{-\infty}^{\infty} h(x); f_{X|Y}(x|y) dx, \quad X, Y \text{ have joint density}$$

- ▶ We can actually define $E[h(X, Y)|Y]$ also as above.
That is,

$$E[h(X, Y)|Y = y] = \int_{-\infty}^{\infty} h(x, y) f_{X|Y}(x|y) dx$$

- ▶ It has all the properties of expectation:

- ▶ The conditional expectation is defined by

$$E[h(X)|Y = y] = \sum_x h(x) f_{X|Y}(x|y), \quad X, Y \text{ are discrete}$$

$$E[h(X)|Y = y] = \int_{-\infty}^{\infty} h(x); f_{X|Y}(x|y) dx, \quad X, Y \text{ have joint density}$$

- ▶ We can actually define $E[h(X, Y)|Y]$ also as above.
That is,

$$E[h(X, Y)|Y = y] = \int_{-\infty}^{\infty} h(x, y) f_{X|Y}(x|y) dx$$

- ▶ It has all the properties of expectation:

1. $E[a|Y] = a$ where a is a constant

- ▶ The conditional expectation is defined by

$$E[h(X)|Y = y] = \sum_x h(x) f_{X|Y}(x|y), \quad X, Y \text{ are discrete}$$

$$E[h(X)|Y = y] = \int_{-\infty}^{\infty} h(x); f_{X|Y}(x|y) dx, \quad X, Y \text{ have joint density}$$

- ▶ We can actually define $E[h(X, Y)|Y]$ also as above.
That is,

$$E[h(X, Y)|Y = y] = \int_{-\infty}^{\infty} h(x, y) f_{X|Y}(x|y) dx$$

- ▶ It has all the properties of expectation:

1. $E[a|Y] = a$ where a is a constant
2. $E[ah_1(X) + bh_2(X)|Y] = aE[h_1(X)|Y] + bE[h_2(X)|Y]$

- ▶ The conditional expectation is defined by

$$E[h(X)|Y = y] = \sum_x h(x) f_{X|Y}(x|y), \quad X, Y \text{ are discrete}$$

$$E[h(X)|Y = y] = \int_{-\infty}^{\infty} h(x); f_{X|Y}(x|y) dx, \quad X, Y \text{ have joint density}$$

- ▶ We can actually define $E[h(X, Y)|Y]$ also as above.
That is,

$$E[h(X, Y)|Y = y] = \int_{-\infty}^{\infty} h(x, y) f_{X|Y}(x|y) dx$$

- ▶ It has all the properties of expectation:

1. $E[a|Y] = a$ where a is a constant
2. $E[ah_1(X) + bh_2(X)|Y] = aE[h_1(X)|Y] + bE[h_2(X)|Y]$
3. $h_1(X) \geq h_2(X) \Rightarrow E[h_1(X)|Y] \geq E[h_2(X)|Y]$

- ▶ Conditional expectation also has some extra properties which are very important

- ▶ Conditional expectation also has some extra properties which are very important
 - ▶ $E[E[h(X)|Y]] = E[h(X)]$

- ▶ Conditional expectation also has some extra properties which are very important
 - ▶ $E[E[h(X)|Y]] = E[h(X)]$
 - ▶ $E[h_1(X)h_2(Y)|Y] = h_2(Y)E[h_1(X)|Y]$

- ▶ Conditional expectation also has some extra properties which are very important

- ▶ $E[E[h(X)|Y]] = E[h(X)]$
- ▶ $E[h_1(X)h_2(Y)|Y] = h_2(Y)E[h_1(X)|Y]$
- ▶ $E[h(X, Y)|Y = y] = E[h(X, y)|Y = y]$

- ▶ Conditional expectation also has some extra properties which are very important
 - ▶ $E[E[h(X)|Y]] = E[h(X)]$
 - ▶ $E[h_1(X)h_2(Y)|Y] = h_2(Y)E[h_1(X)|Y]$
 - ▶ $E[h(X, Y)|Y = y] = E[h(X, y)|Y = y]$
- ▶ We will justify each of these.

- ▶ Conditional expectation also has some extra properties which are very important
 - ▶ $E[E[h(X)|Y]] = E[h(X)]$
 - ▶ $E[h_1(X)h_2(Y)|Y] = h_2(Y)E[h_1(X)|Y]$
 - ▶ $E[h(X, Y)|Y = y] = E[h(X, y)|Y = y]$
- ▶ We will justify each of these.
- ▶ The last property above follows directly from the definition.

- ▶ Expectation of a conditional expectation is the unconditional expectation

- Expectation of a conditional expectation is the unconditional expectation

$$E [E[h(X)|Y]] = E[h(X)]$$

- Expectation of a conditional expectation is the unconditional expectation

$$E [E[h(X)|Y]] = E[h(X)]$$

In the above, LHS is expectation of a function of Y .

- ▶ Expectation of a conditional expectation is the unconditional expectation

$$E [E[h(X)|Y]] = E[h(X)]$$

In the above, LHS is expectation of a function of Y .

- ▶ Let us denote $g(Y) = E[h(X)|Y]$. Then

- ▶ Expectation of a conditional expectation is the unconditional expectation

$$E [E[h(X)|Y]] = E[h(X)]$$

In the above, LHS is expectation of a function of Y .

- ▶ Let us denote $g(Y) = E[h(X)|Y]$. Then

$$E [E[h(X)|Y]] = E[g(Y)]$$

- Expectation of a conditional expectation is the unconditional expectation

$$E [E[h(X)|Y]] = E[h(X)]$$

In the above, LHS is expectation of a function of Y .

- Let us denote $g(Y) = E[h(X)|Y]$. Then

$$\begin{aligned} E [E[h(X)|Y]] &= E[g(Y)] \\ &= \int_{-\infty}^{\infty} g(y) f_Y(y) dy \end{aligned}$$

- Expectation of a conditional expectation is the unconditional expectation

$$E [E[h(X)|Y]] = E[h(X)]$$

In the above, LHS is expectation of a function of Y .

- Let us denote $g(Y) = E[h(X)|Y]$. Then

$$\begin{aligned} E [E[h(X)|Y]] &= E[g(Y)] \\ &= \int_{-\infty}^{\infty} g(y) f_Y(y) dy \\ &= \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} h(x) f_{X|Y}(x|y) dx \right) f_Y(y) dy \end{aligned}$$

- Expectation of a conditional expectation is the unconditional expectation

$$E [E[h(X)|Y]] = E[h(X)]$$

In the above, LHS is expectation of a function of Y .

- Let us denote $g(Y) = E[h(X)|Y]$. Then

$$\begin{aligned} E [E[h(X)|Y]] &= E[g(Y)] \\ &= \int_{-\infty}^{\infty} g(y) f_Y(y) dy \\ &= \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} h(x) f_{X|Y}(x|y) dx \right) f_Y(y) dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} h(x) f_{XY}(x, y) dy dx \end{aligned}$$

- Expectation of a conditional expectation is the unconditional expectation

$$E [E[h(X)|Y]] = E[h(X)]$$

In the above, LHS is expectation of a function of Y .

- Let us denote $g(Y) = E[h(X)|Y]$. Then

$$\begin{aligned} E [E[h(X)|Y]] &= E[g(Y)] \\ &= \int_{-\infty}^{\infty} g(y) f_Y(y) dy \\ &= \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} h(x) f_{X|Y}(x|y) dx \right) f_Y(y) dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} h(x) f_{XY}(x, y) dy dx \\ &= \int_{-\infty}^{\infty} h(x) f_X(x) dx \end{aligned}$$

- Expectation of a conditional expectation is the unconditional expectation

$$E [E[h(X)|Y]] = E[h(X)]$$

In the above, LHS is expectation of a function of Y .

- Let us denote $g(Y) = E[h(X)|Y]$. Then

$$\begin{aligned} E [E[h(X)|Y]] &= E[g(Y)] \\ &= \int_{-\infty}^{\infty} g(y) f_Y(y) dy \\ &= \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} h(x) f_{X|Y}(x|y) dx \right) f_Y(y) dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} h(x) f_{XY}(x, y) dy dx \\ &= \int_{-\infty}^{\infty} h(x) f_X(x) dx \\ &= E[h(X)] \end{aligned}$$

- ▶ Any factor that depends only on the conditioning variable behaves like a constant inside a conditional expectation

- ▶ Any factor that depends only on the conditioning variable behaves like a constant inside a conditional expectation

$$E[h_1(X) h_2(Y)|Y] = h_2(Y)E[h_1(X)|Y]$$

- ▶ Any factor that depends only on the conditioning variable behaves like a constant inside a conditional expectation

$$E[h_1(X) h_2(Y)|Y] = h_2(Y)E[h_1(X)|Y]$$

- ▶ Let us denote $g(Y) = E[h_1(X) h_2(Y)|Y]$

- ▶ Any factor that depends only on the conditioning variable behaves like a constant inside a conditional expectation

$$E[h_1(X) h_2(Y)|Y] = h_2(Y)E[h_1(X)|Y]$$

- ▶ Let us denote $g(Y) = E[h_1(X) h_2(Y)|Y]$

$$g(y) = E[h_1(X) h_2(Y)|Y = y]$$

- ▶ Any factor that depends only on the conditioning variable behaves like a constant inside a conditional expectation

$$E[h_1(X) h_2(Y)|Y] = h_2(Y)E[h_1(X)|Y]$$

- ▶ Let us denote $g(Y) = E[h_1(X) h_2(Y)|Y]$

$$\begin{aligned} g(y) &= E[h_1(X) h_2(Y)|Y = y] \\ &= \int_{-\infty}^{\infty} h_1(x) h_2(y) f_{X|Y}(x|y) dx \end{aligned}$$

- ▶ Any factor that depends only on the conditioning variable behaves like a constant inside a conditional expectation

$$E[h_1(X) h_2(Y)|Y] = h_2(Y)E[h_1(X)|Y]$$

- ▶ Let us denote $g(Y) = E[h_1(X) h_2(Y)|Y]$

$$\begin{aligned} g(y) &= E[h_1(X) h_2(Y)|Y = y] \\ &= \int_{-\infty}^{\infty} h_1(x) h_2(y) f_{X|Y}(x|y) dx \\ &= h_2(y) \int_{-\infty}^{\infty} h_1(x) f_{X|Y}(x|y) dx \end{aligned}$$

- ▶ Any factor that depends only on the conditioning variable behaves like a constant inside a conditional expectation

$$E[h_1(X) h_2(Y)|Y] = h_2(Y)E[h_1(X)|Y]$$

- ▶ Let us denote $g(Y) = E[h_1(X) h_2(Y)|Y]$

$$\begin{aligned} g(y) &= E[h_1(X) h_2(Y)|Y = y] \\ &= \int_{-\infty}^{\infty} h_1(x) h_2(y) f_{X|Y}(x|y) dx \\ &= h_2(y) \int_{-\infty}^{\infty} h_1(x) f_{X|Y}(x|y) dx \\ &= h_2(y) E[h_1(X)|Y = y] \end{aligned}$$

- ▶ Any factor that depends only on the conditioning variable behaves like a constant inside a conditional expectation

$$E[h_1(X) h_2(Y)|Y] = h_2(Y)E[h_1(X)|Y]$$

- ▶ Let us denote $g(Y) = E[h_1(X) h_2(Y)|Y]$

$$\begin{aligned} g(y) &= E[h_1(X) h_2(Y)|Y = y] \\ &= \int_{-\infty}^{\infty} h_1(x) h_2(y) f_{X|Y}(x|y) dx \\ &= h_2(y) \int_{-\infty}^{\infty} h_1(x) f_{X|Y}(x|y) dx \\ &= h_2(y) E[h_1(X)|Y = y] \\ \Rightarrow E[h_1(X) h_2(Y)|Y] &= g(Y) \end{aligned}$$

- ▶ Any factor that depends only on the conditioning variable behaves like a constant inside a conditional expectation

$$E[h_1(X) h_2(Y)|Y] = h_2(Y)E[h_1(X)|Y]$$

- ▶ Let us denote $g(Y) = E[h_1(X) h_2(Y)|Y]$

$$\begin{aligned} g(y) &= E[h_1(X) h_2(Y)|Y = y] \\ &= \int_{-\infty}^{\infty} h_1(x) h_2(y) f_{X|Y}(x|y) dx \\ &= h_2(y) \int_{-\infty}^{\infty} h_1(x) f_{X|Y}(x|y) dx \\ &= h_2(y) E[h_1(X)|Y = y] \\ \Rightarrow E[h_1(X) h_2(Y)|Y] &= g(Y) = h_2(Y)E[h_1(X)|Y] \end{aligned}$$

- ▶ A very useful property of conditional expectation is
 $E[E[X|Y]] = E[X]$

- ▶ A very useful property of conditional expectation is $E[E[X|Y]] = E[X]$ (Assuming all expectations exist)

- ▶ A very useful property of conditional expectation is $E[E[X|Y]] = E[X]$ (Assuming all expectations exist)
- ▶ We can see this in our earlier example.

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ A very useful property of conditional expectation is $E[E[X|Y]] = E[X]$ (Assuming all expectations exist)
- ▶ We can see this in our earlier example.

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We calculated: $EX = \frac{1}{3}$ and $EY = \frac{2}{3}$

- ▶ A very useful property of conditional expectation is $E[E[X|Y]] = E[X]$ (Assuming all expectations exist)
- ▶ We can see this in our earlier example.

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We calculated: $EX = \frac{1}{3}$ and $EY = \frac{2}{3}$
- ▶ We also showed $E[X|Y] = \frac{Y}{2}$

- ▶ A very useful property of conditional expectation is $E[E[X|Y]] = E[X]$ (Assuming all expectations exist)
- ▶ We can see this in our earlier example.

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We calculated: $EX = \frac{1}{3}$ and $EY = \frac{2}{3}$
- ▶ We also showed $E[X|Y] = \frac{Y}{2}$

$$E[E[X|Y]] = E \left[\frac{Y}{2} \right] = \frac{1}{3} = E[X]$$

- ▶ A very useful property of conditional expectation is $E[E[X|Y]] = E[X]$ (Assuming all expectations exist)
- ▶ We can see this in our earlier example.

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We calculated: $EX = \frac{1}{3}$ and $EY = \frac{2}{3}$
- ▶ We also showed $E[X|Y] = \frac{Y}{2}$

$$E[E[X|Y]] = E \left[\frac{Y}{2} \right] = \frac{1}{3} = E[X]$$

- ▶ Similarly

$$E[E[Y|X]] = E \left[\frac{1+X}{2} \right] = \frac{2}{3} = E[Y]$$

Example

- ▶ Let X, Y be random variables with joint density given by

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

Example

- ▶ Let X, Y be random variables with joint density given by

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- ▶ The marginal densities are:

Example

- ▶ Let X, Y be random variables with joint density given by

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- ▶ The marginal densities are:

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy$$

Example

- ▶ Let X, Y be random variables with joint density given by

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- ▶ The marginal densities are:

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_x^{\infty} e^{-y} dy$$

Example

- ▶ Let X, Y be random variables with joint density given by

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- ▶ The marginal densities are:

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_x^{\infty} e^{-y} dy = e^{-x}, \quad x > 0$$

Example

- ▶ Let X, Y be random variables with joint density given by

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- ▶ The marginal densities are:

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_x^{\infty} e^{-y} dy = e^{-x}, \quad x > 0$$

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx$$

Example

- ▶ Let X, Y be random variables with joint density given by

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- ▶ The marginal densities are:

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_x^{\infty} e^{-y} dy = e^{-x}, \quad x > 0$$

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx = \int_0^y e^{-y} dx$$

Example

- ▶ Let X, Y be random variables with joint density given by

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- ▶ The marginal densities are:

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_x^{\infty} e^{-y} dy = e^{-x}, \quad x > 0$$

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx = \int_0^y e^{-y} dx = y e^{-y}, \quad y > 0$$

Example

- ▶ Let X, Y be random variables with joint density given by

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- ▶ The marginal densities are:

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_x^{\infty} e^{-y} dy = e^{-x}, \quad x > 0$$

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx = \int_0^y e^{-y} dx = y e^{-y}, \quad y > 0$$

Thus, X is exponential and Y is gamma.

Example

- ▶ Let X, Y be random variables with joint density given by

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- ▶ The marginal densities are:

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_x^{\infty} e^{-y} dy = e^{-x}, \quad x > 0$$

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx = \int_0^y e^{-y} dx = y e^{-y}, \quad y > 0$$

Thus, X is exponential and Y is gamma.

- ▶ Hence we have

$$EX = 1; \quad \text{Var}(X) = 1;$$

Example

- ▶ Let X, Y be random variables with joint density given by

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- ▶ The marginal densities are:

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_x^{\infty} e^{-y} dy = e^{-x}, \quad x > 0$$

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx = \int_0^y e^{-y} dx = y e^{-y}, \quad y > 0$$

Thus, X is exponential and Y is gamma.

- ▶ Hence we have

$$EX = 1; \quad \text{Var}(X) = 1; \quad EY = 2; \quad \text{Var}(Y) = 2$$

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- Let us calculate covariance of X and Y

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- Let us calculate covariance of X and Y

$$E[XY] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{XY}(x, y) dx dy$$

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- Let us calculate covariance of X and Y

$$\begin{aligned} E[XY] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{XY}(x, y) dx dy \\ &= \int_0^{\infty} \int_0^y xye^{-y} dx dy \end{aligned}$$

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- Let us calculate covariance of X and Y

$$\begin{aligned} E[XY] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{XY}(x, y) dx dy \\ &= \int_0^{\infty} \int_0^y xye^{-y} dx dy = \int_0^{\infty} \frac{1}{2} y^3 e^{-y} dy = 3 \end{aligned}$$

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- Let us calculate covariance of X and Y

$$\begin{aligned} E[XY] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{XY}(x, y) dx dy \\ &= \int_0^{\infty} \int_0^y xye^{-y} dx dy = \int_0^{\infty} \frac{1}{2} y^3 e^{-y} dy = 3 \end{aligned}$$

- Hence, $\text{Cov}(X, Y) = E[XY] - EX EY = 3 - 2 = 1$.

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

- Let us calculate covariance of X and Y

$$\begin{aligned} E[XY] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{XY}(x, y) dx dy \\ &= \int_0^{\infty} \int_0^y xye^{-y} dx dy = \int_0^{\infty} \frac{1}{2} y^3 e^{-y} dy = 3 \end{aligned}$$

- Hence, $\text{Cov}(X, Y) = E[XY] - EX EY = 3 - 2 = 1$.
- $\rho_{XY} = \frac{1}{\sqrt{2}}$

- Recall the joint and marginal densities

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

$$f_X(x) = e^{-x}, \quad x > 0; \quad f_Y(y) = ye^{-y}, \quad y > 0$$

- Recall the joint and marginal densities

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

$$f_X(x) = e^{-x}, \quad x > 0; \quad f_Y(y) = ye^{-y}, \quad y > 0$$

- The conditional densities will be

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)}$$

- Recall the joint and marginal densities

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

$$f_X(x) = e^{-x}, \quad x > 0; \quad f_Y(y) = ye^{-y}, \quad y > 0$$

- The conditional densities will be

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{e^{-y}}{ye^{-y}}$$

- Recall the joint and marginal densities

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

$$f_X(x) = e^{-x}, \quad x > 0; \quad f_Y(y) = ye^{-y}, \quad y > 0$$

- The conditional densities will be

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{e^{-y}}{ye^{-y}} = \frac{1}{y}, \quad 0 < x < y < \infty$$

- Recall the joint and marginal densities

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

$$f_X(x) = e^{-x}, \quad x > 0; \quad f_Y(y) = ye^{-y}, \quad y > 0$$

- The conditional densities will be

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{e^{-y}}{ye^{-y}} = \frac{1}{y}, \quad 0 < x < y < \infty$$

$$f_{Y|X}(y|x) = \frac{f_{XY}(x, y)}{f_X(x)}$$

- Recall the joint and marginal densities

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

$$f_X(x) = e^{-x}, \quad x > 0; \quad f_Y(y) = ye^{-y}, \quad y > 0$$

- The conditional densities will be

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{e^{-y}}{ye^{-y}} = \frac{1}{y}, \quad 0 < x < y < \infty$$

$$f_{Y|X}(y|x) = \frac{f_{XY}(x, y)}{f_X(x)} = \frac{e^{-y}}{e^{-x}}$$

- Recall the joint and marginal densities

$$f_{XY}(x, y) = e^{-y}, \quad 0 < x < y < \infty$$

$$f_X(x) = e^{-x}, \quad x > 0; \quad f_Y(y) = ye^{-y}, \quad y > 0$$

- The conditional densities will be

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{e^{-y}}{ye^{-y}} = \frac{1}{y}, \quad 0 < x < y < \infty$$

$$f_{Y|X}(y|x) = \frac{f_{XY}(x, y)}{f_X(x)} = \frac{e^{-y}}{e^{-x}} = e^{-(y-x)}, \quad 0 < x < y < \infty$$

- The conditional densities are

$$f_{X|Y}(x|y) = \frac{1}{y}; \quad f_{Y|X}(y|x) = e^{-(y-x)}, \quad 0 < x < y < \infty$$

- ▶ The conditional densities are

$$f_{X|Y}(x|y) = \frac{1}{y}; \quad f_{Y|X}(y|x) = e^{-(y-x)}, \quad 0 < x < y < \infty$$

- ▶ We can now calculate the conditional expectation

$$E[X|Y = y] = \int x f_{X|Y}(x|y) dx$$

- ▶ The conditional densities are

$$f_{X|Y}(x|y) = \frac{1}{y}; \quad f_{Y|X}(y|x) = e^{-(y-x)}, \quad 0 < x < y < \infty$$

- ▶ We can now calculate the conditional expectation

$$E[X|Y = y] = \int x f_{X|Y}(x|y) dx = \int_0^y x \frac{1}{y} dx = \frac{y}{2}$$

- ▶ The conditional densities are

$$f_{X|Y}(x|y) = \frac{1}{y}; \quad f_{Y|X}(y|x) = e^{-(y-x)}, \quad 0 < x < y < \infty$$

- ▶ We can now calculate the conditional expectation

$$E[X|Y = y] = \int x f_{X|Y}(x|y) dx = \int_0^y x \frac{1}{y} dx = \frac{y}{2}$$

Thus $E[X|Y] = \frac{Y}{2}$

- ▶ The conditional densities are

$$f_{X|Y}(x|y) = \frac{1}{y}; \quad f_{Y|X}(y|x) = e^{-(y-x)}, \quad 0 < x < y < \infty$$

- ▶ We can now calculate the conditional expectation

$$E[X|Y = y] = \int x f_{X|Y}(x|y) dx = \int_0^y x \frac{1}{y} dx = \frac{y}{2}$$

$$\text{Thus } E[X|Y] = \frac{Y}{2}$$

$$E[Y|X = x] = \int y f_{Y|X}(y|x) dy$$

- ▶ The conditional densities are

$$f_{X|Y}(x|y) = \frac{1}{y}; \quad f_{Y|X}(y|x) = e^{-(y-x)}, \quad 0 < x < y < \infty$$

- ▶ We can now calculate the conditional expectation

$$E[X|Y = y] = \int x f_{X|Y}(x|y) dx = \int_0^y x \frac{1}{y} dx = \frac{y}{2}$$

Thus $E[X|Y] = \frac{Y}{2}$

$$E[Y|X = x] = \int y f_{Y|X}(y|x) dy = \int_x^\infty y e^{-(y-x)} dy$$

- The conditional densities are

$$f_{X|Y}(x|y) = \frac{1}{y}; \quad f_{Y|X}(y|x) = e^{-(y-x)}, \quad 0 < x < y < \infty$$

- We can now calculate the conditional expectation

$$E[X|Y = y] = \int x f_{X|Y}(x|y) dx = \int_0^y x \frac{1}{y} dx = \frac{y}{2}$$

Thus $E[X|Y] = \frac{Y}{2}$

$$\begin{aligned} E[Y|X = x] &= \int y f_{Y|X}(y|x) dy = \int_x^\infty y e^{-(y-x)} dy \\ &= e^x \left(-ye^{-y} \Big|_x^\infty + \int_x^\infty e^{-y} dy \right) \end{aligned}$$

- ▶ The conditional densities are

$$f_{X|Y}(x|y) = \frac{1}{y}; \quad f_{Y|X}(y|x) = e^{-(y-x)}, \quad 0 < x < y < \infty$$

- ▶ We can now calculate the conditional expectation

$$E[X|Y = y] = \int x f_{X|Y}(x|y) dx = \int_0^y x \frac{1}{y} dx = \frac{y}{2}$$

Thus $E[X|Y] = \frac{Y}{2}$

$$\begin{aligned} E[Y|X = x] &= \int y f_{Y|X}(y|x) dy = \int_x^\infty y e^{-(y-x)} dy \\ &= e^x \left(-ye^{-y} \Big|_x^\infty + \int_x^\infty e^{-y} dy \right) \\ &= e^x (xe^{-x} + e^{-x}) = 1 + x \end{aligned}$$

- ▶ The conditional densities are

$$f_{X|Y}(x|y) = \frac{1}{y}; \quad f_{Y|X}(y|x) = e^{-(y-x)}, \quad 0 < x < y < \infty$$

- ▶ We can now calculate the conditional expectation

$$E[X|Y = y] = \int x f_{X|Y}(x|y) dx = \int_0^y x \frac{1}{y} dx = \frac{y}{2}$$

$$\text{Thus } E[X|Y] = \frac{Y}{2}$$

$$\begin{aligned} E[Y|X = x] &= \int y f_{Y|X}(y|x) dy = \int_x^\infty y e^{-(y-x)} dy \\ &= e^x \left(-ye^{-y} \Big|_x^\infty + \int_x^\infty e^{-y} dy \right) \\ &= e^x (xe^{-x} + e^{-x}) = 1 + x \end{aligned}$$

$$\text{Thus, } E[Y|X] = 1 + X$$

► We got

$$E[X|Y] = \frac{Y}{2}; \quad E[Y|X] = 1 + X$$

- ▶ We got

$$E[X|Y] = \frac{Y}{2}; \quad E[Y|X] = 1 + X$$

- ▶ Using this we can verify:

$$E[E[X|Y]] = E \left[\frac{Y}{2} \right]$$

- ▶ We got

$$E[X|Y] = \frac{Y}{2}; \quad E[Y|X] = 1 + X$$

- ▶ Using this we can verify:

$$E[E[X|Y]] = E \left[\frac{Y}{2} \right] = \frac{EY}{2}$$

- ▶ We got

$$E[X|Y] = \frac{Y}{2}; \quad E[Y|X] = 1 + X$$

- ▶ Using this we can verify:

$$E[E[X|Y]] = E \left[\frac{Y}{2} \right] = \frac{EY}{2} = \frac{2}{2} = 1$$

- We got

$$E[X|Y] = \frac{Y}{2}; \quad E[Y|X] = 1 + X$$

- Using this we can verify:

$$E[E[X|Y]] = E \left[\frac{Y}{2} \right] = \frac{EY}{2} = \frac{2}{2} = 1 = EX$$

- We got

$$E[X|Y] = \frac{Y}{2}; \quad E[Y|X] = 1 + X$$

- Using this we can verify:

$$E[E[X|Y]] = E \left[\frac{Y}{2} \right] = \frac{EY}{2} = \frac{2}{2} = 1 = EX$$

$$E[E[Y|X]] = E[1 + X]$$

- ▶ We got

$$E[X|Y] = \frac{Y}{2}; \quad E[Y|X] = 1 + X$$

- ▶ Using this we can verify:

$$E[E[X|Y]] = E \left[\frac{Y}{2} \right] = \frac{EY}{2} = \frac{2}{2} = 1 = EX$$

$$E[E[Y|X]] = E[1 + X] = 1 + 1 = 2$$

- ▶ We got

$$E[X|Y] = \frac{Y}{2}; \quad E[Y|X] = 1 + X$$

- ▶ Using this we can verify:

$$E[E[X|Y]] = E\left[\frac{Y}{2}\right] = \frac{EY}{2} = \frac{2}{2} = 1 = EX$$

$$E[E[Y|X]] = E[1 + X] = 1 + 1 = 2 = EY$$

- ▶ A property of conditional expectation is

$$E[E[X|Y]] = E[X]$$

- ▶ A property of conditional expectation is

$$E[E[X|Y]] = E[X]$$

- ▶ We assume that all three expectations exist.

- ▶ A property of conditional expectation is

$$E[E[X|Y]] = E[X]$$

- ▶ We assume that all three expectations exist.
- ▶ Very useful in calculating expectations

- ▶ A property of conditional expectation is

$$E[E[X|Y]] = E[X]$$

- ▶ We assume that all three expectations exist.
- ▶ Very useful in calculating expectations

$$EX = \sum_y E[X|Y = y] f_Y(y)$$

- ▶ A property of conditional expectation is

$$E[E[X|Y]] = E[X]$$

- ▶ We assume that all three expectations exist.
- ▶ Very useful in calculating expectations

$$EX = \sum_y E[X|Y = y] f_Y(y) \quad \text{or} \quad \int E[X|Y = y] f_Y(y) dy$$

- ▶ A property of conditional expectation is

$$E[E[X|Y]] = E[X]$$

- ▶ We assume that all three expectations exist.
- ▶ Very useful in calculating expectations

$$EX = \sum_y E[X|Y = y] f_Y(y) \quad \text{or} \quad \int E[X|Y = y] f_Y(y) dy$$

- ▶ Can be used to calculate probabilities of events too

- ▶ A property of conditional expectation is

$$E[E[X|Y]] = E[X]$$

- ▶ We assume that all three expectations exist.
- ▶ Very useful in calculating expectations

$$EX = \sum_y E[X|Y = y] f_Y(y) \quad \text{or} \quad \int E[X|Y = y] f_Y(y) dy$$

- ▶ Can be used to calculate probabilities of events too

$$P(A) = E[I_A] = E[E[I_A|Y]]$$

- ▶ Let X be geometric and we want EX .

- ▶ Let X be geometric and we want EX .
- ▶ X is number of tosses needed to get head

- ▶ Let X be geometric and we want EX .
- ▶ X is number of tosses needed to get head
- ▶ Let $Y \in \{0, 1\}$ be outcome of first toss.

- ▶ Let X be geometric and we want EX .
- ▶ X is number of tosses needed to get head
- ▶ Let $Y \in \{0, 1\}$ be outcome of first toss. (1 for head)

- ▶ Let X be geometric and we want EX .
- ▶ X is number of tosses needed to get head
- ▶ Let $Y \in \{0, 1\}$ be outcome of first toss. (1 for head)

$$E[X] = E[E[X|Y]]$$

- ▶ Let X be geometric and we want EX .
- ▶ X is number of tosses needed to get head
- ▶ Let $Y \in \{0, 1\}$ be outcome of first toss. (1 for head)

$$\begin{aligned} E[X] &= E[E[X|Y]] \\ &= E[X|Y = 1] P[Y = 1] + E[X|Y = 0] P[Y = 0] \end{aligned}$$

- ▶ Let X be geometric and we want EX .
- ▶ X is number of tosses needed to get head
- ▶ Let $Y \in \{0, 1\}$ be outcome of first toss. (1 for head)

$$\begin{aligned} E[X] &= E[E[X|Y]] \\ &= E[X|Y = 1] P[Y = 1] + E[X|Y = 0] P[Y = 0] \\ &= E[X|Y = 1] p + E[X|Y = 0] (1 - p) \end{aligned}$$

- ▶ Let X be geometric and we want EX .
- ▶ X is number of tosses needed to get head
- ▶ Let $Y \in \{0, 1\}$ be outcome of first toss. (1 for head)

$$\begin{aligned} E[X] &= E[E[X|Y]] \\ &= E[X|Y = 1] P[Y = 1] + E[X|Y = 0] P[Y = 0] \\ &= E[X|Y = 1] p + E[X|Y = 0] (1 - p) \\ &= 1 p + \end{aligned}$$

- ▶ Let X be geometric and we want EX .
- ▶ X is number of tosses needed to get head
- ▶ Let $Y \in \{0, 1\}$ be outcome of first toss. (1 for head)

$$\begin{aligned} E[X] &= E[E[X|Y]] \\ &= E[X|Y = 1] P[Y = 1] + E[X|Y = 0] P[Y = 0] \\ &= E[X|Y = 1] p + E[X|Y = 0] (1 - p) \\ &= 1 p + (1 + EX)(1 - p) \end{aligned}$$

- ▶ Let X be geometric and we want EX .
- ▶ X is number of tosses needed to get head
- ▶ Let $Y \in \{0, 1\}$ be outcome of first toss. (1 for head)

$$\begin{aligned} E[X] &= E[E[X|Y]] \\ &= E[X|Y = 1] P[Y = 1] + E[X|Y = 0] P[Y = 0] \\ &= E[X|Y = 1] p + E[X|Y = 0] (1 - p) \\ &= 1 p + (1 + EX)(1 - p) \\ \Rightarrow EX (1 - (1 - p)) &= p + (1 - p) \end{aligned}$$

- ▶ Let X be geometric and we want EX .
- ▶ X is number of tosses needed to get head
- ▶ Let $Y \in \{0, 1\}$ be outcome of first toss. (1 for head)

$$\begin{aligned} E[X] &= E[E[X|Y]] \\ &= E[X|Y = 1] P[Y = 1] + E[X|Y = 0] P[Y = 0] \\ &= E[X|Y = 1] p + E[X|Y = 0] (1 - p) \\ &= 1 p + (1 + EX)(1 - p) \\ \Rightarrow EX (1 - (1 - p)) &= p + (1 - p) \\ \Rightarrow EX p &= 1 \end{aligned}$$

- ▶ Let X be geometric and we want EX .
- ▶ X is number of tosses needed to get head
- ▶ Let $Y \in \{0, 1\}$ be outcome of first toss. (1 for head)

$$\begin{aligned} E[X] &= E[E[X|Y]] \\ &= E[X|Y = 1] P[Y = 1] + E[X|Y = 0] P[Y = 0] \\ &= E[X|Y = 1] p + E[X|Y = 0] (1 - p) \\ &= 1 p + (1 + EX)(1 - p) \\ \Rightarrow EX (1 - (1 - p)) &= p + (1 - p) \\ \Rightarrow EX p &= 1 \\ \Rightarrow EX &= \frac{1}{p} \end{aligned}$$

- ▶ $P[X = k|Y = 1] = 1$ if $k = 1$ (otherwise it is zero) and hence $E[X|Y = 1] = 1$

- $P[X = k|Y = 1] = 1$ if $k = 1$ (otherwise it is zero) and hence $E[X|Y = 1] = 1$

$$P[X = k|Y = 0] = \begin{cases} 0 & \text{if } k = 1 \\ \frac{(1-p)^{k-1}p}{(1-p)} & \text{if } k \geq 2 \end{cases}$$

- $P[X = k|Y = 1] = 1$ if $k = 1$ (otherwise it is zero) and hence $E[X|Y = 1] = 1$

$$P[X = k|Y = 0] = \begin{cases} 0 & \text{if } k = 1 \\ \frac{(1-p)^{k-1}p}{(1-p)} & \text{if } k \geq 2 \end{cases}$$

Hence

- $P[X = k|Y = 1] = 1$ if $k = 1$ (otherwise it is zero) and hence $E[X|Y = 1] = 1$

$$P[X = k|Y = 0] = \begin{cases} 0 & \text{if } k = 1 \\ \frac{(1-p)^{k-1}p}{(1-p)} & \text{if } k \geq 2 \end{cases}$$

Hence

$$E[X|Y = 0] = \sum_{k=2}^{\infty} k (1-p)^{k-2} p$$

- $P[X = k|Y = 1] = 1$ if $k = 1$ (otherwise it is zero) and hence $E[X|Y = 1] = 1$

$$P[X = k|Y = 0] = \begin{cases} 0 & \text{if } k = 1 \\ \frac{(1-p)^{k-1}p}{(1-p)} & \text{if } k \geq 2 \end{cases}$$

Hence

$$\begin{aligned} E[X|Y = 0] &= \sum_{k=2}^{\infty} k (1-p)^{k-2} p \\ &= \sum_{k=2}^{\infty} (k-1) (1-p)^{k-2} p + \sum_{k=2}^{\infty} (1-p)^{k-2} p \end{aligned}$$

- $P[X = k|Y = 1] = 1$ if $k = 1$ (otherwise it is zero) and hence $E[X|Y = 1] = 1$

$$P[X = k|Y = 0] = \begin{cases} 0 & \text{if } k = 1 \\ \frac{(1-p)^{k-1}p}{(1-p)} & \text{if } k \geq 2 \end{cases}$$

Hence

$$\begin{aligned} E[X|Y = 0] &= \sum_{k=2}^{\infty} k (1-p)^{k-2} p \\ &= \sum_{k=2}^{\infty} (k-1) (1-p)^{k-2} p + \sum_{k=2}^{\infty} (1-p)^{k-2} p \\ &= \sum_{k'=1}^{\infty} k' (1-p)^{k'-1} p + \sum_{k'=1}^{\infty} (1-p)^{k'-1} p \end{aligned}$$

- $P[X = k|Y = 1] = 1$ if $k = 1$ (otherwise it is zero) and hence $E[X|Y = 1] = 1$

$$P[X = k|Y = 0] = \begin{cases} 0 & \text{if } k = 1 \\ \frac{(1-p)^{k-1}p}{(1-p)} & \text{if } k \geq 2 \end{cases}$$

Hence

$$\begin{aligned} E[X|Y = 0] &= \sum_{k=2}^{\infty} k (1-p)^{k-2} p \\ &= \sum_{k=2}^{\infty} (k-1) (1-p)^{k-2} p + \sum_{k=2}^{\infty} (1-p)^{k-2} p \\ &= \sum_{k'=1}^{\infty} k' (1-p)^{k'-1} p + \sum_{k'=1}^{\infty} (1-p)^{k'-1} p \\ &= EX + 1 \end{aligned}$$

Another example

- ▶ Example: multiple rounds of the party game

Another example

- ▶ Example: multiple rounds of the party game
- ▶ Let R_n denote number of rounds when you start with n people.

Another example

- ▶ Example: multiple rounds of the party game
- ▶ Let R_n denote number of rounds when you start with n people.
- ▶ We want $\bar{R}_n = E[R_n]$.

Another example

- ▶ Example: multiple rounds of the party game
- ▶ Let R_n denote number of rounds when you start with n people.
- ▶ We want $\bar{R}_n = E[R_n]$.
- ▶ We want to use $E[R_n] = E[E[R_n|X_n]]$

Another example

- ▶ Example: multiple rounds of the party game
- ▶ Let R_n denote number of rounds when you start with n people.
- ▶ We want $\bar{R}_n = E[R_n]$.
- ▶ We want to use $E[R_n] = E[E[R_n|X_n]]$
- ▶ We need to think of a useful X_n .

Another example

- ▶ Example: multiple rounds of the party game
- ▶ Let R_n denote number of rounds when you start with n people.
- ▶ We want $\bar{R}_n = E[R_n]$.
- ▶ We want to use $E[R_n] = E[E[R_n|X_n]]$
- ▶ We need to think of a useful X_n .
- ▶ Let X_n be the number of people who got their own hat in the first round with n people.

- ▶ R_n – number of rounds when you start with n people.

- ▶ R_n – number of rounds when you start with n people.
- ▶ X_n – number of people who got their own hat in the first round

- ▶ R_n – number of rounds when you start with n people.
- ▶ X_n – number of people who got their own hat in the first round

$$E[R_n] = E[E[R_n|X_n]]$$

- ▶ R_n – number of rounds when you start with n people.
- ▶ X_n – number of people who got their own hat in the first round

$$\begin{aligned} E[R_n] &= E[E[R_n | X_n]] \\ &= \sum_{i=0}^n E[R_n | X_n = i] P[X_n = i] \end{aligned}$$

- ▶ R_n – number of rounds when you start with n people.
- ▶ X_n – number of people who got their own hat in the first round

$$\begin{aligned}
 E[R_n] &= E[E[R_n | X_n]] \\
 &= \sum_{i=0}^n E[R_n | X_n = i] P[X_n = i] \\
 &= \sum_{i=0}^n (1 + E[R_{n-i}]) P[X_n = i]
 \end{aligned}$$

- ▶ R_n – number of rounds when you start with n people.
- ▶ X_n – number of people who got their own hat in the first round

$$\begin{aligned}
 E[R_n] &= E[E[R_n | X_n]] \\
 &= \sum_{i=0}^n E[R_n | X_n = i] P[X_n = i] \\
 &= \sum_{i=0}^n (1 + E[R_{n-i}]) P[X_n = i] \\
 &= \sum_{i=0}^n P[X_n = i] + \sum_{i=0}^n E[R_{n-i}] P[X_n = i]
 \end{aligned}$$

- ▶ R_n – number of rounds when you start with n people.
- ▶ X_n – number of people who got their own hat in the first round

$$\begin{aligned}
 E[R_n] &= E[E[R_n | X_n]] \\
 &= \sum_{i=0}^n E[R_n | X_n = i] P[X_n = i] \\
 &= \sum_{i=0}^n (1 + E[R_{n-i}]) P[X_n = i] \\
 &= \sum_{i=0}^n P[X_n = i] + \sum_{i=0}^n E[R_{n-i}] P[X_n = i]
 \end{aligned}$$

If we can guess value of $E[R_n]$ then we can prove it using mathematical induction

► What would be $E[X_n]$?

- ▶ What would be $E[X_n]$?
- ▶ Let $Y_i \in \{0, 1\}$ denote whether or not i^{th} person got his own hat.

- ▶ What would be $E[X_n]$?
- ▶ Let $Y_i \in \{0, 1\}$ denote whether or not i^{th} person got his own hat.
- ▶ We know

$$E[Y_i] = P[Y_i = 1] = \frac{(n-1)!}{n!} = \frac{1}{n}$$

- ▶ What would be $E[X_n]$?
- ▶ Let $Y_i \in \{0, 1\}$ denote whether or not i^{th} person got his own hat.
- ▶ We know

$$E[Y_i] = P[Y_i = 1] = \frac{(n-1)!}{n!} = \frac{1}{n}$$

Now,
$$X_n = \sum_{i=1}^n Y_i$$

- ▶ What would be $E[X_n]$?
- ▶ Let $Y_i \in \{0, 1\}$ denote whether or not i^{th} person got his own hat.
- ▶ We know

$$E[Y_i] = P[Y_i = 1] = \frac{(n-1)!}{n!} = \frac{1}{n}$$

$$\text{Now, } X_n = \sum_{i=1}^n Y_i \text{ and hence } EX_n = \sum_{i=1}^n E[Y_i] = 1$$

- ▶ What would be $E[X_n]$?
- ▶ Let $Y_i \in \{0, 1\}$ denote whether or not i^{th} person got his own hat.
- ▶ We know

$$E[Y_i] = P[Y_i = 1] = \frac{(n-1)!}{n!} = \frac{1}{n}$$

$$\text{Now, } X_n = \sum_{i=1}^n Y_i \text{ and hence } EX_n = \sum_{i=1}^n E[Y_i] = 1$$

- ▶ Hence a good guess is $E[R_n] = n$.

- ▶ What would be $E[X_n]$?
- ▶ Let $Y_i \in \{0, 1\}$ denote whether or not i^{th} person got his own hat.
- ▶ We know

$$E[Y_i] = P[Y_i = 1] = \frac{(n-1)!}{n!} = \frac{1}{n}$$

$$\text{Now, } X_n = \sum_{i=1}^n Y_i \text{ and hence } EX_n = \sum_{i=1}^n E[Y_i] = 1$$

- ▶ Hence a good guess is $E[R_n] = n$.
- ▶ We verify it using mathematical induction. We know $E[R_1] = 1$

► Assume: $E[R_k] = k, \quad 1 \leq k \leq n-1$

► Assume: $E[R_k] = k, \quad 1 \leq k \leq n-1$

$$E[R_n] = \sum_{i=0}^n P[X_n = i] + \sum_{i=0}^n E[R_{n-i}] P[X_n = i]$$

► Assume: $E[R_k] = k, \quad 1 \leq k \leq n-1$

$$\begin{aligned} E[R_n] &= \sum_{i=0}^n P[X_n = i] + \sum_{i=0}^n E[R_{n-i}] P[X_n = i] \\ &= 1 + E[R_n] P[X_n = 0] + \sum_{i=1}^n E[R_{n-i}] P[X_n = i] \end{aligned}$$

► Assume: $E[R_k] = k, \quad 1 \leq k \leq n-1$

$$\begin{aligned} E[R_n] &= \sum_{i=0}^n P[X_n = i] + \sum_{i=0}^n E[R_{n-i}] P[X_n = i] \\ &= 1 + E[R_n] P[X_n = 0] + \sum_{i=1}^n E[R_{n-i}] P[X_n = i] \\ &= 1 + E[R_n] P[X_n = 0] + \sum_{i=1}^n (n-i) P[X_n = i] \end{aligned}$$

► Assume: $E[R_k] = k, \quad 1 \leq k \leq n-1$

$$\begin{aligned} E[R_n] &= \sum_{i=0}^n P[X_n = i] + \sum_{i=0}^n E[R_{n-i}] P[X_n = i] \\ &= 1 + E[R_n] P[X_n = 0] + \sum_{i=1}^n E[R_{n-i}] P[X_n = i] \\ &= 1 + E[R_n] P[X_n = 0] + \sum_{i=1}^n (n-i) P[X_n = i] \\ E[R_n] (1 - P[X_n = 0]) &= 1 + n(1 - P[X_n = 0]) - \sum_{i=1}^n i P[X_n = i] \end{aligned}$$

► Assume: $E[R_k] = k, \quad 1 \leq k \leq n-1$

$$\begin{aligned} E[R_n] &= \sum_{i=0}^n P[X_n = i] + \sum_{i=0}^n E[R_{n-i}] P[X_n = i] \\ &= 1 + E[R_n] P[X_n = 0] + \sum_{i=1}^n E[R_{n-i}] P[X_n = i] \\ &= 1 + E[R_n] P[X_n = 0] + \sum_{i=1}^n (n-i) P[X_n = i] \\ E[R_n] (1 - P[X_n = 0]) &= 1 + n(1 - P[X_n = 0]) - \sum_{i=1}^n i P[X_n = i] \\ &= 1 + n(1 - P[X_n = 0]) - E[X_n] \end{aligned}$$

► Assume: $E[R_k] = k, \quad 1 \leq k \leq n-1$

$$\begin{aligned} E[R_n] &= \sum_{i=0}^n P[X_n = i] + \sum_{i=0}^n E[R_{n-i}] P[X_n = i] \\ &= 1 + E[R_n] P[X_n = 0] + \sum_{i=1}^n E[R_{n-i}] P[X_n = i] \\ &= 1 + E[R_n] P[X_n = 0] + \sum_{i=1}^n (n-i) P[X_n = i] \\ E[R_n] (1 - P[X_n = 0]) &= 1 + n(1 - P[X_n = 0]) - \sum_{i=1}^n i P[X_n = i] \\ &= 1 + n(1 - P[X_n = 0]) - E[X_n] \\ &= 1 + n(1 - P[X_n = 0]) - 1 \end{aligned}$$

► Assume: $E[R_k] = k, \quad 1 \leq k \leq n-1$

$$\begin{aligned} E[R_n] &= \sum_{i=0}^n P[X_n = i] + \sum_{i=0}^n E[R_{n-i}] P[X_n = i] \\ &= 1 + E[R_n] P[X_n = 0] + \sum_{i=1}^n E[R_{n-i}] P[X_n = i] \\ &= 1 + E[R_n] P[X_n = 0] + \sum_{i=1}^n (n-i) P[X_n = i] \\ E[R_n] (1 - P[X_n = 0]) &= 1 + n(1 - P[X_n = 0]) - \sum_{i=1}^n i P[X_n = i] \\ &= 1 + n(1 - P[X_n = 0]) - E[X_n] \\ &= 1 + n(1 - P[X_n = 0]) - 1 \\ \Rightarrow E[R_n] &= n \end{aligned}$$

Analysis of Quicksort

- ▶ Given n numbers we want to sort them. Many algorithms.

Analysis of Quicksort

- ▶ Given n numbers we want to sort them. Many algorithms.
- ▶ Complexity – order of the number of comparisons needed

Analysis of Quicksort

- ▶ Given n numbers we want to sort them. Many algorithms.
- ▶ Complexity – order of the number of comparisons needed
- ▶ Quicksort: Choose a pivot. Separate numbers into two parts – less and greater than pivot, do recursively

Analysis of Quicksort

- ▶ Given n numbers we want to sort them. Many algorithms.
- ▶ Complexity – order of the number of comparisons needed
- ▶ Quicksort: Choose a pivot. Separate numbers into two parts – less and greater than pivot, do recursively
- ▶ Separating into two parts takes $n - 1$ comparisons.

Analysis of Quicksort

- ▶ Given n numbers we want to sort them. Many algorithms.
- ▶ Complexity – order of the number of comparisons needed
- ▶ Quicksort: Choose a pivot. Separate numbers into two parts – less and greater than pivot, do recursively
- ▶ Separating into two parts takes $n - 1$ comparisons.
- ▶ Suppose the two parts contain m and $n - m - 1$. Comparisons needed to Separate each of them into two parts depends on m

Analysis of Quicksort

- ▶ Given n numbers we want to sort them. Many algorithms.
- ▶ Complexity – order of the number of comparisons needed
- ▶ Quicksort: Choose a pivot. Separate numbers into two parts – less and greater than pivot, do recursively
- ▶ Separating into two parts takes $n - 1$ comparisons.
- ▶ Suppose the two parts contain m and $n - m - 1$. Comparisons needed to Separate each of them into two parts depends on m
- ▶ So, final number of comparisons depends on the ‘number of rounds’

quicksort details

- ▶ Given $\{x_1, \dots, x_n\}$.

quicksort details

- ▶ Given $\{x_1, \dots, x_n\}$.
- ▶ Choose first as pivot

$$\{x_{j_1}, x_{j_2}, \dots, x_{j_m}\} x_1 \{x_{k_1}, x_{k_2}, \dots, x_{k_{n-1-m}}\}$$

quicksort details

- ▶ Given $\{x_1, \dots, x_n\}$.
- ▶ Choose first as pivot

$$\{x_{j_1}, x_{j_2}, \dots, x_{j_m}\} x_1 \{x_{k_1}, x_{k_2}, \dots, x_{k_{n-1-m}}\}$$

- ▶ Suppose r_n is the number of comparisons. If we get (roughly) equal parts, then

$$r_n \approx n + 2r_{n/2} = n + 2(n/2 + 2r_{n/4}) = n + n + 4r_{n/4} = \dots = n \log_2(n)$$

quicksort details

- ▶ Given $\{x_1, \dots, x_n\}$.
- ▶ Choose first as pivot

$$\{x_{j_1}, x_{j_2}, \dots, x_{j_m}\} x_1 \{x_{k_1}, x_{k_2}, \dots, x_{k_{n-1-m}}\}$$

- ▶ Suppose r_n is the number of comparisons. If we get (roughly) equal parts, then

$$r_n \approx n + 2r_{n/2} = n + 2(n/2 + 2r_{n/4}) = n + n + 4r_{n/4} = \dots = n \log_2(n)$$

- ▶ If all the rest go into one part, then

$$r_n = n + r_{n-1} = n + (n-1) + r_{n-2} = \dots = \frac{n(n+1)}{2}$$

quicksort details

- ▶ Given $\{x_1, \dots, x_n\}$.
- ▶ Choose first as pivot

$$\{x_{j_1}, x_{j_2}, \dots, x_{j_m}\} x_1 \{x_{k_1}, x_{k_2}, \dots, x_{k_{n-1-m}}\}$$

- ▶ Suppose r_n is the number of comparisons. If we get (roughly) equal parts, then

$$r_n \approx n + 2r_{n/2} = n + 2(n/2 + 2r_{n/4}) = n + n + 4r_{n/4} = \dots = n \log_2(n)$$

- ▶ If all the rest go into one part, then

$$r_n = n + r_{n-1} = n + (n-1) + r_{n-2} = \dots = \frac{n(n+1)}{2}$$

- ▶ If you are lucky, $O(n \log(n))$ comparisons.

quicksort details

- ▶ Given $\{x_1, \dots, x_n\}$.
- ▶ Choose first as pivot

$$\{x_{j_1}, x_{j_2}, \dots, x_{j_m}\} x_1 \{x_{k_1}, x_{k_2}, \dots, x_{k_{n-1-m}}\}$$

- ▶ Suppose r_n is the number of comparisons. If we get (roughly) equal parts, then

$$r_n \approx n + 2r_{n/2} = n + 2(n/2 + 2r_{n/4}) = n + n + 4r_{n/4} = \dots = n \log_2(n)$$

- ▶ If all the rest go into one part, then

$$r_n = n + r_{n-1} = n + (n-1) + r_{n-2} = \dots = \frac{n(n+1)}{2}$$

- ▶ If you are lucky, $O(n \log(n))$ comparisons.
- ▶ If unlucky, in the worst case, $O(n^2)$ comparisons

quicksort details

- ▶ Given $\{x_1, \dots, x_n\}$.
- ▶ Choose first as pivot

$$\{x_{j_1}, x_{j_2}, \dots, x_{j_m}\} x_1 \{x_{k_1}, x_{k_2}, \dots, x_{k_{n-1-m}}\}$$

- ▶ Suppose r_n is the number of comparisons. If we get (roughly) equal parts, then

$$r_n \approx n + 2r_{n/2} = n + 2(n/2 + 2r_{n/4}) = n + n + 4r_{n/4} = \dots = n \log_2(n)$$

- ▶ If all the rest go into one part, then

$$r_n = n + r_{n-1} = n + (n-1) + r_{n-2} = \dots = \frac{n(n+1)}{2}$$

- ▶ If you are lucky, $O(n \log(n))$ comparisons.
- ▶ If unlucky, in the worst case, $O(n^2)$ comparisons
- ▶ Question: 'on the average' how many comparisons?

Average case complexity of quicksort

Average case complexity of quicksort

- ▶ Assume pivot is equally likely to be the smallest or second smallest or m^{th} smallest.

Average case complexity of quicksort

- ▶ Assume pivot is equally likely to be the smallest or second smallest or m^{th} smallest.
- ▶ M_n – number of comparisons.

Average case complexity of quicksort

- ▶ Assume pivot is equally likely to be the smallest or second smallest or m^{th} smallest.
- ▶ M_n – number of comparisons.
- ▶ Define: $X = j$ if pivot is j^{th} smallest

Average case complexity of quicksort

- ▶ Assume pivot is equally likely to be the smallest or second smallest or m^{th} smallest.
- ▶ M_n – number of comparisons.
- ▶ Define: $X = j$ if pivot is j^{th} smallest
- ▶ Given $X = j$ we know $M_n = (n - 1) + M_{j-1} + M_{n-j}$.

Average case complexity of quicksort

- ▶ Assume pivot is equally likely to be the smallest or second smallest or m^{th} smallest.
- ▶ M_n – number of comparisons.
- ▶ Define: $X = j$ if pivot is j^{th} smallest
- ▶ Given $X = j$ we know $M_n = (n - 1) + M_{j-1} + M_{n-j}$.

$$E[M_n] = E[E[M_n|X]]$$

Average case complexity of quicksort

- ▶ Assume pivot is equally likely to be the smallest or second smallest or m^{th} smallest.
- ▶ M_n – number of comparisons.
- ▶ Define: $X = j$ if pivot is j^{th} smallest
- ▶ Given $X = j$ we know $M_n = (n - 1) + M_{j-1} + M_{n-j}$.

$$E[M_n] = E[E[M_n|X]] = \sum_{j=1}^n E[M_n|X = j] P[X = j]$$

Average case complexity of quicksort

- ▶ Assume pivot is equally likely to be the smallest or second smallest or m^{th} smallest.
- ▶ M_n – number of comparisons.
- ▶ Define: $X = j$ if pivot is j^{th} smallest
- ▶ Given $X = j$ we know $M_n = (n - 1) + M_{j-1} + M_{n-j}$.

$$\begin{aligned} E[M_n] &= E[E[M_n|X]] = \sum_{j=1}^n E[M_n|X = j] P[X = j] \\ &= \sum_{j=1}^n E[(n - 1) + M_{j-1} + M_{n-j}] \frac{1}{n} \end{aligned}$$

Average case complexity of quicksort

- ▶ Assume pivot is equally likely to be the smallest or second smallest or m^{th} smallest.
- ▶ M_n – number of comparisons.
- ▶ Define: $X = j$ if pivot is j^{th} smallest
- ▶ Given $X = j$ we know $M_n = (n - 1) + M_{j-1} + M_{n-j}$.

$$\begin{aligned} E[M_n] &= E[E[M_n|X]] = \sum_{j=1}^n E[M_n|X = j] P[X = j] \\ &= \sum_{j=1}^n E[(n - 1) + M_{j-1} + M_{n-j}] \frac{1}{n} \\ &= (n - 1) + \frac{2}{n} \sum_{k=1}^{n-1} E[M_k], \quad (\text{taking } M_0 = 0) \end{aligned}$$

Average case complexity of quicksort

- ▶ Assume pivot is equally likely to be the smallest or second smallest or m^{th} smallest.
- ▶ M_n – number of comparisons.
- ▶ Define: $X = j$ if pivot is j^{th} smallest
- ▶ Given $X = j$ we know $M_n = (n - 1) + M_{j-1} + M_{n-j}$.

$$\begin{aligned} E[M_n] &= E[E[M_n|X]] = \sum_{j=1}^n E[M_n|X = j] P[X = j] \\ &= \sum_{j=1}^n E[(n - 1) + M_{j-1} + M_{n-j}] \frac{1}{n} \\ &= (n - 1) + \frac{2}{n} \sum_{k=1}^{n-1} E[M_k], \quad (\text{taking } M_0 = 0) \end{aligned}$$

- ▶ This is a recurrence relation. (A little complicated to solve)

Least squares estimation

- ▶ We want to estimate Y as a function of X .

Least squares estimation

- ▶ We want to estimate Y as a function of X .
- ▶ We want an estimate with minimum mean square error.

Least squares estimation

- ▶ We want to estimate Y as a function of X .
- ▶ We want an estimate with minimum mean square error.
- ▶ We want to solve (the min is over all functions g)

$$\min_g E (Y - g(X))^2$$

Least squares estimation

- ▶ We want to estimate Y as a function of X .
- ▶ We want an estimate with minimum mean square error.
- ▶ We want to solve (the min is over all functions g)

$$\min_g E (Y - g(X))^2$$

- ▶ Earlier we considered only linear functions:
 $g(X) = aX + b$

Least squares estimation

- ▶ We want to estimate Y as a function of X .
- ▶ We want an estimate with minimum mean square error.
- ▶ We want to solve (the min is over all functions g)

$$\min_g E (Y - g(X))^2$$

- ▶ Earlier we considered only linear functions:
 $g(X) = aX + b$
- ▶ Now we want the 'best' function (linear or nonlinear)

Least squares estimation

- ▶ We want to estimate Y as a function of X .
- ▶ We want an estimate with minimum mean square error.
- ▶ We want to solve (the min is over all functions g)

$$\min_g E (Y - g(X))^2$$

- ▶ Earlier we considered only linear functions:
 $g(X) = aX + b$
- ▶ Now we want the 'best' function (linear or nonlinear)
- ▶ The solution now turns out to be

$$g^*(X) = E[Y|X]$$

Least squares estimation

- ▶ We want to estimate Y as a function of X .
- ▶ We want an estimate with minimum mean square error.
- ▶ We want to solve (the min is over all functions g)

$$\min_g E(Y - g(X))^2$$

- ▶ Earlier we considered only linear functions:
 $g(X) = aX + b$
- ▶ Now we want the 'best' function (linear or nonlinear)
- ▶ The solution now turns out to be

$$g^*(X) = E[Y|X]$$

- ▶ Let us prove this.

- We want to show that for all g

$$E \left[(E[Y | X] - Y)^2 \right] \leq E \left[(g(X) - Y)^2 \right]$$

- ▶ We want to show that for all g

$$E \left[(E[Y | X] - Y)^2 \right] \leq E \left[(g(X) - Y)^2 \right]$$

- ▶ We have

$$(g(X) - Y)^2 = \left[(g(X) - E[Y | X]) + (E[Y | X] - Y) \right]^2$$

- We want to show that for all g

$$E \left[(E[Y | X] - Y)^2 \right] \leq E \left[(g(X) - Y)^2 \right]$$

- We have

$$\begin{aligned} (g(X) - Y)^2 &= \left[(g(X) - E[Y | X]) + (E[Y | X] - Y) \right]^2 \\ &= (g(X) - E[Y | X])^2 + (E[Y | X] - Y)^2 \\ &\quad + 2(g(X) - E[Y | X])(E[Y | X] - Y) \end{aligned}$$

- We want to show that for all g

$$E \left[(E[Y | X] - Y)^2 \right] \leq E \left[(g(X) - Y)^2 \right]$$

- We have

$$\begin{aligned} (g(X) - Y)^2 &= \left[(g(X) - E[Y | X]) + (E[Y | X] - Y) \right]^2 \\ &= (g(X) - E[Y | X])^2 + (E[Y | X] - Y)^2 \\ &\quad + 2(g(X) - E[Y | X])(E[Y | X] - Y) \end{aligned}$$

- Now we can take expectation on both sides.

- We want to show that for all g

$$E[(E[Y | X] - Y)^2] \leq E[(g(X) - Y)^2]$$

- We have

$$\begin{aligned}(g(X) - Y)^2 &= [(g(X) - E[Y | X]) + (E[Y | X] - Y)]^2 \\&= (g(X) - E[Y | X])^2 + (E[Y | X] - Y)^2 \\&\quad + 2(g(X) - E[Y | X])(E[Y | X] - Y)\end{aligned}$$

- Now we can take expectation on both sides.
- We first show that expectation of last term on RHS above is zero.

First consider the last term

$$E[(g(X) - E[Y | X])(E[Y | X] - Y)]$$

First consider the last term

$$\begin{aligned} & E \left[(g(X) - E[Y | X])(E[Y | X] - Y) \right] \\ = & E \left[E \left\{ (g(X) - E[Y | X])(E[Y | X] - Y) \mid X \right\} \right] \end{aligned}$$

First consider the last term

$$\begin{aligned} & E \left[(g(X) - E[Y | X])(E[Y | X] - Y) \right] \\ = & E \left[E \left\{ (g(X) - E[Y | X])(E[Y | X] - Y) \mid X \right\} \right] \\ & \text{because } E[Z] = E[E[Z|X]] \end{aligned}$$

First consider the last term

$$\begin{aligned} & E[(g(X) - E[Y | X])(E[Y | X] - Y)] \\ = & E[E\{(g(X) - E[Y | X])(E[Y | X] - Y) | X\}] \\ = & E[(g(X) - E[Y | X]) E\{(E[Y | X] - Y) | X\}] \end{aligned}$$

First consider the last term

$$\begin{aligned} & E \left[(g(X) - E[Y | X])(E[Y | X] - Y) \right] \\ = & E \left[E \left\{ (g(X) - E[Y | X])(E[Y | X] - Y) \mid X \right\} \right] \\ = & E \left[(g(X) - E[Y | X]) E \left\{ (E[Y | X] - Y) \mid X \right\} \right] \\ & \text{because } E[h_1(X)h_2(Z)|X] = h_1(X) E[h_2(Z)|X] \end{aligned}$$

First consider the last term

$$\begin{aligned} & E[(g(X) - E[Y | X])(E[Y | X] - Y)] \\ = & E[E\{(g(X) - E[Y | X])(E[Y | X] - Y) | X\}] \\ = & E[(g(X) - E[Y | X]) E\{(E[Y | X] - Y) | X\}] \\ = & E[(g(X) - E[Y | X]) (E\{E[Y | X] | X\} - E\{Y | X\})] \end{aligned}$$

First consider the last term

$$\begin{aligned} & E[(g(X) - E[Y | X])(E[Y | X] - Y)] \\ = & E[E\{(g(X) - E[Y | X])(E[Y | X] - Y) | X\}] \\ = & E[(g(X) - E[Y | X]) E\{(E[Y | X] - Y) | X\}] \\ = & E[(g(X) - E[Y | X]) (E\{E[Y | X] | X\} - E\{Y | X\})] \\ = & E[(g(X) - E[Y | X]) (E[Y | X] - E[Y | X])] \end{aligned}$$

First consider the last term

$$\begin{aligned} & E[(g(X) - E[Y | X])(E[Y | X] - Y)] \\ = & E[E\{(g(X) - E[Y | X])(E[Y | X] - Y) | X\}] \\ = & E[(g(X) - E[Y | X]) E\{(E[Y | X] - Y) | X\}] \\ = & E[(g(X) - E[Y | X]) (E\{E[Y | X] | X\} - E\{Y | X\})] \\ = & E[(g(X) - E[Y | X]) (E[Y | X] - E[Y | X])] \\ = & 0 \end{aligned}$$

► We earlier got

► We earlier got

$$\begin{aligned}(g(X) - Y)^2 &= (g(X) - E[Y | X])^2 + (E[Y | X] - Y)^2 \\ &\quad + 2(g(X) - E[Y | X])(E[Y | X] - Y)\end{aligned}$$

► We earlier got

$$\begin{aligned}(g(X) - Y)^2 &= (g(X) - E[Y | X])^2 + (E[Y | X] - Y)^2 \\ &\quad + 2(g(X) - E[Y | X])(E[Y | X] - Y)\end{aligned}$$

► Hence we get

► We earlier got

$$\begin{aligned}(g(X) - Y)^2 &= (g(X) - E[Y | X])^2 + (E[Y | X] - Y)^2 \\ &\quad + 2(g(X) - E[Y | X])(E[Y | X] - Y)\end{aligned}$$

► Hence we get

$$\begin{aligned}E [(g(X) - Y)^2] &= E [(g(X) - E[Y | X])^2] \\ &\quad + E [(E[Y | X] - Y)^2]\end{aligned}$$

► We earlier got

$$\begin{aligned}(g(X) - Y)^2 &= (g(X) - E[Y | X])^2 + (E[Y | X] - Y)^2 \\ &\quad + 2(g(X) - E[Y | X])(E[Y | X] - Y)\end{aligned}$$

► Hence we get

$$\begin{aligned}E \left[(g(X) - Y)^2 \right] &= E \left[(g(X) - E[Y | X])^2 \right] \\ &\quad + E \left[(E[Y | X] - Y)^2 \right] \\ &\geq E \left[(E[Y | X] - Y)^2 \right]\end{aligned}$$

- We earlier got

$$\begin{aligned}(g(X) - Y)^2 &= (g(X) - E[Y | X])^2 + (E[Y | X] - Y)^2 \\ &\quad + 2(g(X) - E[Y | X])(E[Y | X] - Y)\end{aligned}$$

- Hence we get

$$\begin{aligned}E[(g(X) - Y)^2] &= E[(g(X) - E[Y | X])^2] \\ &\quad + E[(E[Y | X] - Y)^2] \\ &\geq E[(E[Y | X] - Y)^2]\end{aligned}$$

- Since the above is true for all functions g , we get

$$g^*(X) = E[Y | X]$$