

Recap: Brownian Motion

- ▶ $\{X(t), t \geq 0\}$ is called a Brownian motion if
 1. $X(0) = 0$
 2. The process has stationary and independent increments
 3. For every $t > 0$, $X(t)$ is Gaussian with mean 0 and variance $\sigma^2 t$

Recap: Brownian Motion

- ▶ $\{X(t), t \geq 0\}$ is called a Brownian motion if
 1. $X(0) = 0$
 2. The process has stationary and independent increments
 3. For every $t > 0$, $X(t)$ is Gaussian with mean 0 and variance $\sigma^2 t$
- ▶ If $\sigma^2 = 1$, it is called standard Brownian Motion

Recap: Brownian Motion

- ▶ $\{X(t), t \geq 0\}$ is called a Brownian motion if
 1. $X(0) = 0$
 2. The process has stationary and independent increments
 3. For every $t > 0$, $X(t)$ is Gaussian with mean 0 and variance $\sigma^2 t$
- ▶ If $\sigma^2 = 1$, it is called standard Brownian Motion
- ▶ $Y(t) = X(t) + \mu t$ is called Brownian motion with a drift

Recap: Some properties

- ▶ An important result is that Brownian motion paths are continuous

Recap: Some properties

- ▶ An important result is that Brownian motion paths are continuous
- ▶ The autocorrelation and autocovariance of the process is

$$R(t_1, t_2) = \sigma^2 \min(t_1, t_2)$$

Recap: Some properties

- ▶ An important result is that Brownian motion paths are continuous
- ▶ The autocorrelation and autocovariance of the process is

$$R(t_1, t_2) = \sigma^2 \min(t_1, t_2) = \text{Cov}(X(t_1), X(t_2))$$

Recap: Some properties

- ▶ An important result is that Brownian motion paths are continuous
- ▶ The autocorrelation and autocovariance of the process is

$$R(t_1, t_2) = \sigma^2 \min(t_1, t_2) = \text{Cov}(X(t_1), X(t_2))$$

- ▶ All n^{th} order distributions are Gaussian:

Recap: Some properties

- ▶ An important result is that Brownian motion paths are continuous
- ▶ The autocorrelation and autocovariance of the process is

$$R(t_1, t_2) = \sigma^2 \min(t_1, t_2) = \text{Cov}(X(t_1), X(t_2))$$

- ▶ All n^{th} order distributions are Gaussian:
 $X(t_1), \dots, X(t_n)$ are jointly Gaussian.

Recap: Gaussian Processes

- ▶ A continuous-time continuous-state process $\{X(t), t \geq 0\}$ is said to be a Gaussian process if for all n and all t_1, t_2, \dots, t_n , we have that $X(t_1), \dots, X(t_n)$ are jointly Gaussian.

Recap: Gaussian Processes

- ▶ A continuous-time continuous-state process $\{X(t), t \geq 0\}$ is said to be a Gaussian process if for all n and all t_1, t_2, \dots, t_n , we have that $X(t_1), \dots, X(t_n)$ are jointly Gaussian.
- ▶ A general Gaussian process is specified by the mean function and the variance and covariance functions

Recap: Gaussian Processes

- ▶ A continuous-time continuous-state process $\{X(t), t \geq 0\}$ is said to be a Gaussian process if for all n and all t_1, t_2, \dots, t_n , we have that $X(t_1), \dots, X(t_n)$ are jointly Gaussian.
- ▶ A general Gaussian process is specified by the mean function and the variance and covariance functions
- ▶ The Brownian motion is an example of a Gaussian Process

Recap: Gaussian Processes

- ▶ A continuous-time continuous-state process $\{X(t), t \geq 0\}$ is said to be a Gaussian process if for all n and all t_1, t_2, \dots, t_n , we have that $X(t_1), \dots, X(t_n)$ are jointly Gaussian.
- ▶ A general Gaussian process is specified by the mean function and the variance and covariance functions
- ▶ The Brownian motion is an example of a Gaussian Process
- ▶ The Brownian motion is a Gaussian process with

$$E[X(t)] = 0, \quad \text{Var}(X(t)) = \sigma^2 t, \quad \text{Cov}(X(s), X(t)) = \sigma^2 \min(s, t)$$

Recap: Conditional Densities

- For $s < t$, $f_{X(s)|X(t)}$ is Gaussian and

$$E[X(s)|X(t)] = \frac{s}{t} X(t); \quad \text{Var}(X(s)|X(t)) = \frac{s}{t} (t - s)$$

Recap: Conditional Densities

- For $s < t$, $f_{X(s)|X(t)}$ is Gaussian and

$$E[X(s)|X(t)] = \frac{s}{t} X(t); \quad \text{Var}(X(s)|X(t)) = \frac{s}{t} (t - s)$$

- For $s < t$, we also showed $f_{X(t)|X(s)}$ is Gaussian with

$$E[X(t) | X(s)] = X(s); \quad \text{Var}(X(t)|X(s)) = (t - s)$$

Recap: Hitting Times

- ▶ Let T_a denote the first time Brownian motion hits a .

$$Pr[T_a \leq t] = \frac{2}{\sqrt{2\pi}} \int_{|a|/\sqrt{t}}^{\infty} e^{-\frac{y^2}{2}} dy$$

Recap: Hitting Times

- ▶ Let T_a denote the first time Brownian motion hits a .

$$Pr[T_a \leq t] = \frac{2}{\sqrt{2\pi}} \int_{|a|/\sqrt{t}}^{\infty} e^{-\frac{y^2}{2}} dy$$

- ▶ By continuity of sample paths,

$$Pr[\max_{0 \leq s \leq t} X(s) \geq a] = Pr[T_a \leq t] = 2Pr[X(t) \geq a]$$

Recap: Geometric Brownian Motion

- ▶ Let $\{Y(t), t \geq 0\}$ be a Brownian motion with drift.
Define

$$X(t) = e^{Y(t)}$$

Recap: Geometric Brownian Motion

- ▶ Let $\{Y(t), t \geq 0\}$ be a Brownian motion with drift. Define

$$X(t) = e^{Y(t)}$$

- ▶ Then, $\{X(t), t \geq 0\}$ is called geometric Brownian motion. It is useful in mathematical finance

- ▶ We have considered three random processes

- ▶ We have considered three random processes
- ▶ (discrete-time) Markov Chain

- ▶ We have considered three random processes
- ▶ (discrete-time) Markov Chain
 - Example of Discrete-time discrete-state process

- ▶ We have considered three random processes
- ▶ (discrete-time) Markov Chain
 - Example of Discrete-time discrete-state process
- ▶ Continuous time Markov Chains (e.g., Poisson Process)

- ▶ We have considered three random processes
- ▶ (discrete-time) Markov Chain
 - Example of Discrete-time discrete-state process
- ▶ Continuous time Markov Chains (e.g., Poisson Process)
 - Example of continuous-time discrete-state process

- ▶ We have considered three random processes
- ▶ (discrete-time) Markov Chain
 - Example of Discrete-time discrete-state process
- ▶ Continuous time Markov Chains (e.g., Poisson Process)
 - Example of continuous-time discrete-state process
- ▶ Brownian Motion

- ▶ We have considered three random processes
- ▶ (discrete-time) Markov Chain
 - Example of Discrete-time discrete-state process
- ▶ Continuous time Markov Chains (e.g., Poisson Process)
 - Example of continuous-time discrete-state process
- ▶ Brownian Motion
 - Example of continuous-time continuous-state process

- ▶ We have considered three random processes
- ▶ (discrete-time) Markov Chain
 - Example of Discrete-time discrete-state process
- ▶ Continuous time Markov Chains (e.g., Poisson Process)
 - Example of continuous-time discrete-state process
- ▶ Brownian Motion
 - Example of continuous-time continuous-state process
- ▶ We need an example of discrete-time continuous-state process!

- ▶ We have considered three random processes
- ▶ (discrete-time) Markov Chain
 - Example of Discrete-time discrete-state process
- ▶ Continuous time Markov Chains (e.g., Poisson Process)
 - Example of continuous-time discrete-state process
- ▶ Brownian Motion
 - Example of continuous-time continuous-state process
- ▶ We need an example of discrete-time continuous-state process!
- ▶ Any sequence of continuous random variables would be a discrete-time continuous-state process

- ▶ In general, any 'stochastic' algorithm would generate discrete-state continuous time process.

- ▶ In general, any ‘stochastic’ algorithm would generate discrete-state continuous time process.
- ▶ If an algorithm uses a random step, then the algorithm would be like

$$X(n+1) = X(n) + \eta_n G(X(n), \xi(n))$$

where $\xi(n)$ would be some random variable which may be dependent on $X(n)$.

- ▶ In general, any ‘stochastic’ algorithm would generate discrete-state continuous time process.
- ▶ If an algorithm uses a random step, then the algorithm would be like

$$X(n+1) = X(n) + \eta_n G(X(n), \xi(n))$$

where $\xi(n)$ would be some random variable which may be dependent on $X(n)$.

- ▶ Many algorithms can be written in this general form.

- ▶ In general, any ‘stochastic’ algorithm would generate discrete-state continuous time process.
- ▶ If an algorithm uses a random step, then the algorithm would be like

$$X(n+1) = X(n) + \eta_n G(X(n), \xi(n))$$

where $\xi(n)$ would be some random variable which may be dependent on $X(n)$.

- ▶ Many algorithms can be written in this general form.
- ▶ The $X(n), n = 0, 1, \dots$ would be a discrete-time continuous-state stochastic process

- ▶ In general, any ‘stochastic’ algorithm would generate discrete-state continuous time process.
- ▶ If an algorithm uses a random step, then the algorithm would be like

$$X(n+1) = X(n) + \eta_n G(X(n), \xi(n))$$

where $\xi(n)$ would be some random variable which may be dependent on $X(n)$.

- ▶ Many algorithms can be written in this general form.
- ▶ The $X(n), n = 0, 1, \dots$ would be a discrete-time continuous-state stochastic process
- ▶ We consider an important class of such processes

- ▶ Let $\{X_n, n = 0, 1, \dots\}$ be a discrete-time continuous-state process.

- ▶ Let $\{X_n, n = 0, 1, \dots\}$ be a discrete-time continuous-state process.
- ▶ It is called a **martingale** if $E|X_n| < \infty, \forall n$ and

$$E[X_{n+1} \mid X_n, \dots, X_0] = X_n, \forall n$$

- ▶ Let $\{X_n, n = 0, 1, \dots\}$ be a discrete-time continuous-state process.
- ▶ It is called a **martingale** if $E|X_n| < \infty, \forall n$ and

$$E[X_{n+1} \mid X_n, \dots, X_0] = X_n, \forall n$$

- ▶ One can think of Martingale as a general fair gambling game.

- ▶ Let $\{X_n, n = 0, 1, \dots\}$ be a discrete-time continuous-state process.
- ▶ It is called a **martingale** if $E|X_n| < \infty, \forall n$ and

$$E[X_{n+1} \mid X_n, \dots, X_0] = X_n, \forall n$$

- ▶ One can think of Martingale as a general fair gambling game.
- ▶ When X_n is a martingale, we have

$$E[X_{n+1}] = E[X_n], \forall n$$

Example

- ▶ Sequence of sums of independent random variables forms a Martingale

Example

- ▶ Sequence of sums of independent random variables forms a Martingale
- ▶ Suppose Z_i are iid with
 $Pr[Z_i = +1] = Pr[Z_i = -1] = 0.5$.

Example

- ▶ Sequence of sums of independent random variables forms a Martingale
- ▶ Suppose Z_i are iid with $Pr[Z_i = +1] = Pr[Z_i = -1] = 0.5$. Let

$$X_n = \sum_{i=1}^n Z_i$$

Example

- ▶ Sequence of sums of independent random variables forms a Martingale
- ▶ Suppose Z_i are iid with $Pr[Z_i = +1] = Pr[Z_i = -1] = 0.5$. Let

$$X_n = \sum_{i=1}^n Z_i \quad \Rightarrow \quad X_{n+1} = X_n + Z_{n+1}$$

Example

- ▶ Sequence of sums of independent random variables forms a Martingale
- ▶ Suppose Z_i are iid with $Pr[Z_i = +1] = Pr[Z_i = -1] = 0.5$. Let

$$X_n = \sum_{i=1}^n Z_i \quad \Rightarrow \quad X_{n+1} = X_n + Z_{n+1}$$

- ▶ Since $EZ_i = 0$, $\forall i$,

Example

- ▶ Sequence of sums of independent random variables forms a Martingale
- ▶ Suppose Z_i are iid with $Pr[Z_i = +1] = Pr[Z_i = -1] = 0.5$. Let

$$X_n = \sum_{i=1}^n Z_i \quad \Rightarrow \quad X_{n+1} = X_n + Z_{n+1}$$

- ▶ Since $EZ_i = 0, \forall i$,

$$E[X_{n+1} \mid X_n, \dots] = E[X_n + Z_{n+1} \mid X_n, \dots]$$

Example

- ▶ Sequence of sums of independent random variables forms a Martingale
- ▶ Suppose Z_i are iid with $Pr[Z_i = +1] = Pr[Z_i = -1] = 0.5$. Let

$$X_n = \sum_{i=1}^n Z_i \quad \Rightarrow \quad X_{n+1} = X_n + Z_{n+1}$$

- ▶ Since $EZ_i = 0, \forall i$,

$$E[X_{n+1} \mid X_n, \dots] = E[X_n + Z_{n+1} \mid X_n, \dots] = X_n + E[Z_{n+1} \mid X_n, \dots] = X_n$$

Example

- ▶ Sequence of sums of independent random variables forms a Martingale
- ▶ Suppose Z_i are iid with $Pr[Z_i = +1] = Pr[Z_i = -1] = 0.5$. Let

$$X_n = \sum_{i=1}^n Z_i \quad \Rightarrow \quad X_{n+1} = X_n + Z_{n+1}$$

- ▶ Since $EZ_i = 0, \forall i$,

$$E[X_{n+1} \mid X_n, \dots] = E[X_n + Z_{n+1} \mid X_n, \dots] = X_n + E[Z_{n+1} \mid X_n, \dots] = X_n$$

- ▶ Hence, X_n is a martingale.

Example

- ▶ Suppose X_1, X_2, \dots are iid with $E[X_i] = 1$.

Example

- ▶ Suppose X_1, X_2, \dots are iid with $E[X_i] = 1$.
- ▶ Let $Z_n = \prod_{i=1}^n X_i$.

Example

- ▶ Suppose X_1, X_2, \dots are iid with $E[X_i] = 1$.
- ▶ Let $Z_n = \prod_{i=1}^n X_i$.
- ▶ Since $Z_{n+1} = Z_n X_{n+1}$,

Example

- ▶ Suppose X_1, X_2, \dots are iid with $E[X_i] = 1$.
- ▶ Let $Z_n = \prod_{i=1}^n X_i$.
- ▶ Since $Z_{n+1} = Z_n X_{n+1}$,

$$E[Z_{n+1} | Z_n, \dots, Z_1] = E[Z_n X_{n+1} | Z_n, \dots, Z_1]$$

Example

- ▶ Suppose X_1, X_2, \dots are iid with $E[X_i] = 1$.
- ▶ Let $Z_n = \prod_{i=1}^n X_i$.
- ▶ Since $Z_{n+1} = Z_n X_{n+1}$,

$$\begin{aligned} E[Z_{n+1} | Z_n, \dots, Z_1] &= E[Z_n X_{n+1} | Z_n, \dots, Z_1] \\ &= Z_n E[X_{n+1} | Z_n, \dots, Z_1] \end{aligned}$$

Example

- ▶ Suppose X_1, X_2, \dots are iid with $E[X_i] = 1$.
- ▶ Let $Z_n = \prod_{i=1}^n X_i$.
- ▶ Since $Z_{n+1} = Z_n X_{n+1}$,

$$\begin{aligned} E[Z_{n+1} | Z_n, \dots, Z_1] &= E[Z_n X_{n+1} | Z_n, \dots, Z_1] \\ &= Z_n E[X_{n+1} | Z_n, \dots, Z_1] \\ &= Z_n, \text{ since } X_i \text{ iid, } E[X_i] = 1 \end{aligned}$$

Example

- ▶ Suppose X_1, X_2, \dots are iid with $E[X_i] = 1$.
- ▶ Let $Z_n = \prod_{i=1}^n X_i$.
- ▶ Since $Z_{n+1} = Z_n X_{n+1}$,

$$\begin{aligned} E[Z_{n+1} | Z_n, \dots, Z_1] &= E[Z_n X_{n+1} | Z_n, \dots, Z_1] \\ &= Z_n E[X_{n+1} | Z_n, \dots, Z_1] \\ &= Z_n, \text{ since } X_i \text{ iid, } E[X_i] = 1 \end{aligned}$$

- ▶ Thus, Z_n is a martingale

- Given $\{X_n, n = 1, 2, \dots\}$ and $E|X_n| < \infty, \forall n$, It is called a martingale if

$$E[X_{n+1} \mid X_n, \dots, X_1] = X_n, \forall n$$

- ▶ Given $\{X_n, n = 1, 2, \dots\}$ and $E|X_n| < \infty, \forall n$, It is called a martingale if

$$E[X_{n+1} \mid X_n, \dots, X_1] = X_n, \forall n$$

- ▶ We may often actually show

$$E[X_{n+1} \mid Y_n, \dots, Y_1] = X_n, \forall n$$

Where Y_1, \dots, Y_n determine X_1, \dots, X_n for all n .

- ▶ Given $\{X_n, n = 1, 2, \dots\}$ and $E|X_n| < \infty, \forall n$, It is called a martingale if

$$E[X_{n+1} \mid X_n, \dots, X_1] = X_n, \forall n$$

- ▶ We may often actually show

$$E[X_{n+1} \mid Y_n, \dots, Y_1] = X_n, \forall n$$

Where Y_1, \dots, Y_n determine X_1, \dots, X_n for all n .

- ▶ This is justified by the following two identities

- Given $\{X_n, n = 1, 2, \dots\}$ and $E|X_n| < \infty, \forall n$, It is called a martingale if

$$E[X_{n+1} \mid X_n, \dots, X_1] = X_n, \forall n$$

- We may often actually show

$$E[X_{n+1} \mid Y_n, \dots, Y_1] = X_n, \forall n$$

Where Y_1, \dots, Y_n determine X_1, \dots, X_n for all n .

- This is justified by the following two identities

$$\begin{aligned} E[E[X_{n+1} \mid X_n, \dots, X_1, Y_n, \dots, Y_1] \mid X_1, \dots, X_n] \\ = E[X_{n+1} \mid X_n, \dots, X_1] \end{aligned}$$

for any X_i, Y_i by properties of conditional expectation

- Given $\{X_n, n = 1, 2, \dots\}$ and $E|X_n| < \infty, \forall n$, It is called a martingale if

$$E[X_{n+1} \mid X_n, \dots, X_1] = X_n, \forall n$$

- We may often actually show

$$E[X_{n+1} \mid Y_n, \dots, Y_1] = X_n, \forall n$$

Where Y_1, \dots, Y_n determine X_1, \dots, X_n for all n .

- This is justified by the following two identities

$$\begin{aligned} E[E[X_{n+1} \mid X_n, \dots, X_1, Y_n, \dots, Y_1] \mid X_1, \dots, X_n] \\ = E[X_{n+1} \mid X_n, \dots, X_1] \end{aligned}$$

for any X_i, Y_i by properties of conditional expectation

$$E[X_{n+1} \mid X_n, \dots, X_1, Y_n, \dots, Y_1] = E[X_{n+1} \mid Y_n, \dots, Y_1]$$

when Y_1, \dots, Y_n determine X_1, \dots, X_n for all n

A property of Conditional Expectation

- ▶ Conditional expectation satisfies

$$E[E[h(X)|Y, Z] | Y] = E[h(X)|Y]$$

A property of Conditional Expectation

- ▶ Conditional expectation satisfies

$$E[E[h(X)|Y, Z] | Y] = E[h(X)|Y]$$

Note that all these can be random vectors.

A property of Conditional Expectation

- ▶ Conditional expectation satisfies

$$E[E[h(X)|Y, Z] | Y] = E[h(X)|Y]$$

Note that all these can be random vectors.

- ▶ Let

$$\begin{aligned} g_1(Y, Z) &= E[h(X)|Y, Z] \\ g_2(Y) &= E[g_1(Y, Z)|Y] \end{aligned}$$

A property of Conditional Expectation

- ▶ Conditional expectation satisfies

$$E[E[h(X)|Y, Z] | Y] = E[h(X)|Y]$$

Note that all these can be random vectors.

- ▶ Let

$$\begin{aligned} g_1(Y, Z) &= E[h(X)|Y, Z] \\ g_2(Y) &= E[g_1(Y, Z)|Y] \end{aligned}$$

We want to show $g_2(Y) = E[h(X)|Y]$

► Recall: $g_1(Y, Z) = E[h(X)|Y, Z]$, $g_2(Y) = E[g_1(Y, Z)|Y]$

► Recall: $g_1(Y, Z) = E[h(X)|Y, Z]$, $g_2(Y) = E[g_1(Y, Z)|Y]$

$$g_2(y) = \int g_1(y, z) f_{Z|Y}(z|y) dz$$

► Recall: $g_1(Y, Z) = E[h(X)|Y, Z]$, $g_2(Y) = E[g_1(Y, Z)|Y]$

$$\begin{aligned} g_2(y) &= \int g_1(y, z) f_{Z|Y}(z|y) dz \\ &= \int \left[\int h(x) f_{X|YZ}(x|y, z) dx \right] f_{Z|Y}(z|y) dz \end{aligned}$$

► Recall: $g_1(Y, Z) = E[h(X)|Y, Z]$, $g_2(Y) = E[g_1(Y, Z)|Y]$

$$\begin{aligned} g_2(y) &= \int g_1(y, z) f_{Z|Y}(z|y) dz \\ &= \int \left[\int h(x) f_{X|YZ}(x|y, z) dx \right] f_{Z|Y}(z|y) dz \\ &= \int h(x) \left[\int f_{X|YZ}(x|y, z) f_{Z|Y}(z|y) dz \right] dx \end{aligned}$$

► Recall: $g_1(Y, Z) = E[h(X)|Y, Z]$, $g_2(Y) = E[g_1(Y, Z)|Y]$

$$\begin{aligned} g_2(y) &= \int g_1(y, z) f_{Z|Y}(z|y) dz \\ &= \int \left[\int h(x) f_{X|YZ}(x|y, z) dx \right] f_{Z|Y}(z|y) dz \\ &= \int h(x) \left[\int f_{X|YZ}(x|y, z) f_{Z|Y}(z|y) dz \right] dx \\ &= \int h(x) \left[\int f_{XZ|Y}(x, z|y) dz \right] dx \end{aligned}$$

► Recall: $g_1(Y, Z) = E[h(X)|Y, Z]$, $g_2(Y) = E[g_1(Y, Z)|Y]$

$$\begin{aligned} g_2(y) &= \int g_1(y, z) f_{Z|Y}(z|y) dz \\ &= \int \left[\int h(x) f_{X|YZ}(x|y, z) dx \right] f_{Z|Y}(z|y) dz \\ &= \int h(x) \left[\int f_{X|YZ}(x|y, z) f_{Z|Y}(z|y) dz \right] dx \\ &= \int h(x) \left[\int f_{XZ|Y}(x, z|y) dz \right] dx \\ &= \int h(x) f_{X|Y}(x|y) dx \end{aligned}$$

- Recall: $g_1(Y, Z) = E[h(X)|Y, Z]$, $g_2(Y) = E[g_1(Y, Z)|Y]$

$$\begin{aligned} g_2(y) &= \int g_1(y, z) f_{Z|Y}(z|y) dz \\ &= \int \left[\int h(x) f_{X|YZ}(x|y, z) dx \right] f_{Z|Y}(z|y) dz \\ &= \int h(x) \left[\int f_{X|YZ}(x|y, z) f_{Z|Y}(z|y) dz \right] dx \\ &= \int h(x) \left[\int f_{XZ|Y}(x, z|y) dz \right] dx \\ &= \int h(x) f_{X|Y}(x|y) dx \end{aligned}$$

- Thus we get

$$E[E[h(X)|Y, Z] | Y] = E[h(X)|Y]$$

Example

- ▶ We saw that partial sums of iid X_i with $E[X_i] = 0$ is a martingale.

Example

- ▶ We saw that partial sums of iid X_i with $E[X_i] = 0$ is a martingale.
- ▶ We can generalize this.

Example

- ▶ We saw that partial sums of iid X_i with $E[X_i] = 0$ is a martingale.
- ▶ We can generalize this.
- ▶ Let X_1, X_2, \dots any sequence of random variables.

Example

- ▶ We saw that partial sums of iid X_i with $E[X_i] = 0$ is a martingale.
- ▶ We can generalize this.
- ▶ Let X_1, X_2, \dots any sequence of random variables.
- ▶ Let $W_i = X_i - E[X_i \mid X_1, \dots, X_{i-1}]$.

Example

- ▶ We saw that partial sums of iid X_i with $E[X_i] = 0$ is a martingale.
- ▶ We can generalize this.
- ▶ Let X_1, X_2, \dots any sequence of random variables.
- ▶ Let $W_i = X_i - E[X_i \mid X_1, \dots, X_{i-1}]$.
- ▶ Now $E[W_i] = 0, \forall i$ though W_i may not be independent.

Example

- ▶ We saw that partial sums of iid X_i with $E[X_i] = 0$ is a martingale.
- ▶ We can generalize this.
- ▶ Let X_1, X_2, \dots any sequence of random variables.
- ▶ Let $W_i = X_i - E[X_i \mid X_1, \dots, X_{i-1}]$.
- ▶ Now $E[W_i] = 0, \forall i$ though W_i may not be independent.
- ▶ Let $Z_n = \sum_{i=1}^n W_i$.

Example

- ▶ We saw that partial sums of iid X_i with $E[X_i] = 0$ is a martingale.
- ▶ We can generalize this.
- ▶ Let X_1, X_2, \dots any sequence of random variables.
- ▶ Let $W_i = X_i - E[X_i \mid X_1, \dots, X_{i-1}]$.
- ▶ Now $E[W_i] = 0, \forall i$ though W_i may not be independent.
- ▶ Let $Z_n = \sum_{i=1}^n W_i$.
- ▶ We can show that $Z_n, n = 1, 2, \dots$ is a martingale, assuming that $E[|Z_n|] < \infty$.

► We have

$$Z_n = \sum_{i=1}^n W_i = \sum_{i=1}^n X_i - E[X_i \mid X_1, \dots, X_{i-1}]$$

- ▶ We have

$$Z_n = \sum_{i=1}^n W_i = \sum_{i=1}^n X_i - E[X_i \mid X_1, \dots, X_{i-1}]$$

- ▶ Note that X_1, \dots, X_n determine $Z_1, \dots, Z_n, \forall n$

- ▶ We have

$$Z_n = \sum_{i=1}^n W_i = \sum_{i=1}^n X_i - E[X_i | X_1, \dots, X_{i-1}]$$

- ▶ Note that X_1, \dots, X_n determine $Z_1, \dots, Z_n, \forall n$
- ▶ We have, $Z_{n+1} = Z_n + X_{n+1} - E[X_{n+1} | X_1, \dots, X_n]$.

- ▶ We have

$$Z_n = \sum_{i=1}^n W_i = \sum_{i=1}^n X_i - E[X_i | X_1, \dots, X_{i-1}]$$

- ▶ Note that X_1, \dots, X_n determine $Z_1, \dots, Z_n, \forall n$
- ▶ We have, $Z_{n+1} = Z_n + X_{n+1} - E[X_{n+1} | X_1, \dots, X_n]$.

$$E[Z_{n+1} | X_1, \dots, X_n] = E[Z_n + W_{n+1} | X_1, \dots, X_n]$$

- ▶ We have

$$Z_n = \sum_{i=1}^n W_i = \sum_{i=1}^n X_i - E[X_i | X_1, \dots, X_{i-1}]$$

- ▶ Note that X_1, \dots, X_n determine $Z_1, \dots, Z_n, \forall n$
- ▶ We have, $Z_{n+1} = Z_n + X_{n+1} - E[X_{n+1} | X_1, \dots, X_n]$.

$$\begin{aligned} E[Z_{n+1} | X_1, \dots, X_n] &= E[Z_n + W_{n+1} | X_1, \dots, X_n] \\ &= Z_n + \end{aligned}$$

- ▶ We have

$$Z_n = \sum_{i=1}^n W_i = \sum_{i=1}^n X_i - E[X_i | X_1, \dots, X_{i-1}]$$

- ▶ Note that X_1, \dots, X_n determine $Z_1, \dots, Z_n, \forall n$
- ▶ We have, $Z_{n+1} = Z_n + X_{n+1} - E[X_{n+1} | X_1, \dots, X_n]$.

$$\begin{aligned} E[Z_{n+1} | X_1, \dots, X_n] &= E[Z_n + W_{n+1} | X_1, \dots, X_n] \\ &= Z_n + E[X_{n+1} | X_1, \dots, X_n] - \end{aligned}$$

- We have

$$Z_n = \sum_{i=1}^n W_i = \sum_{i=1}^n X_i - E[X_i | X_1, \dots, X_{i-1}]$$

- Note that X_1, \dots, X_n determine $Z_1, \dots, Z_n, \forall n$
- We have, $Z_{n+1} = Z_n + X_{n+1} - E[X_{n+1} | X_1, \dots, X_n]$.

$$\begin{aligned} E[Z_{n+1} | X_1, \dots, X_n] &= E[Z_n + W_{n+1} | X_1, \dots, X_n] \\ &= Z_n + E[X_{n+1} | X_1, \dots, X_n] - \\ &\quad E[X_{n+1} | X_1, \dots, X_n] \end{aligned}$$

- ▶ We have

$$Z_n = \sum_{i=1}^n W_i = \sum_{i=1}^n X_i - E[X_i | X_1, \dots, X_{i-1}]$$

- ▶ Note that X_1, \dots, X_n determine $Z_1, \dots, Z_n, \forall n$
- ▶ We have, $Z_{n+1} = Z_n + X_{n+1} - E[X_{n+1} | X_1, \dots, X_n]$.

$$\begin{aligned} E[Z_{n+1} | X_1, \dots, X_n] &= E[Z_n + W_{n+1} | X_1, \dots, X_n] \\ &= Z_n + E[X_{n+1} | X_1, \dots, X_n] - \\ &\quad E[X_{n+1} | X_1, \dots, X_n] \\ &= Z_n \end{aligned}$$

- ▶ This shows that Z_n is a martingale

- ▶ Let N be a positive integer valued random variable with $P[N < \infty] = 1$.

- ▶ Let N be a positive integer valued random variable with $P[N < \infty] = 1$.
- ▶ N is said to be a stopping time for the process Z_1, Z_2, \dots if
the event $[N = n]$ is determined by the random variable Z_1, \dots, Z_n .

- ▶ Let N be a positive integer valued random variable with $P[N < \infty] = 1$.
- ▶ N is said to be a stopping time for the process Z_1, Z_2, \dots if
the event $[N = n]$ is determined by the random variable Z_1, \dots, Z_n .
- ▶ If we know the values of Z_1, \dots, Z_n , then we can say whether or not $N = n$.

- ▶ Let N be a positive integer valued random variable with $P[N < \infty] = 1$.
- ▶ N is said to be a stopping time for the process Z_1, Z_2, \dots if
the event $[N = n]$ is determined by the random variable Z_1, \dots, Z_n .
- ▶ If we know the values of Z_1, \dots, Z_n , then we can say whether or not $N = n$.
- ▶ The idea is that we can decide to stop the process at N and the decision to stop cannot anticipate the future

- ▶ Let N be a stopping time for the process $\{Z_n, n \geq 1\}$.

- ▶ Let N be a stopping time for the process $\{Z_n, n \geq 1\}$.
- ▶ Define

$$\bar{Z}_n = \begin{cases} Z_n & \text{if } n \leq N \\ Z_N & \text{if } n > N \end{cases}$$

- ▶ Let N be a stopping time for the process $\{Z_n, n \geq 1\}$.
- ▶ Define

$$\bar{Z}_n = \begin{cases} Z_n & \text{if } n \leq N \\ Z_N & \text{if } n > N \end{cases}$$

- ▶ We call $\{\bar{Z}_n, n \geq 1\}$ the stopped process

- ▶ Let $\{Z_n, n \geq 1\}$ be a martingale and let N a stopping time for it.

- ▶ Let $\{Z_n, n \geq 1\}$ be a martingale and let N a stopping time for it.
- ▶ **Theorem:** The stopped Process is a martingale

- ▶ Let $\{Z_n, n \geq 1\}$ be a martingale and let N a stopping time for it.
- ▶ **Theorem:** The stopped Process is a martingale
That is, a stopped martingale is a martingale

- ▶ Let $\{Z_n, n \geq 1\}$ be a martingale and let N a stopping time for it.
- ▶ **Theorem:** The stopped Process is a martingale
That is, a stopped martingale is a martingale
- ▶ Note that, by definition, $\bar{Z}_1 = Z_1$.

- ▶ Let $\{Z_n, n \geq 1\}$ be a martingale and let N a stopping time for it.
- ▶ **Theorem:** The stopped Process is a martingale
That is, a stopped martingale is a martingale
- ▶ Note that, by definition, $\bar{Z}_1 = Z_1$.
- ▶ So, if the stopped process is a martingale, then,

$$E[\bar{Z}_n] = E[\bar{Z}_1] = E[Z_1], \forall n$$

- ▶ Let $\{Z_n, n \geq 1\}$ be a martingale and let N a stopping time for it.
- ▶ **Theorem:** The stopped Process is a martingale
That is, a stopped martingale is a martingale
- ▶ Note that, by definition, $\bar{Z}_1 = Z_1$.
- ▶ So, if the stopped process is a martingale, then,

$$E[\bar{Z}_n] = E[\bar{Z}_1] = E[Z_1], \forall n$$

- ▶ Recall that for a martingale, $E[Z_n] = E[Z_1], \forall n$

- ▶ Let $\{Z_n, n \geq 1\}$ be a martingale and let N a stopping time for it.
- ▶ **Theorem:** The stopped Process is a martingale
That is, a stopped martingale is a martingale
- ▶ Note that, by definition, $\bar{Z}_1 = Z_1$.
- ▶ So, if the stopped process is a martingale, then,

$$E[\bar{Z}_n] = E[\bar{Z}_1] = E[Z_1], \forall n$$

- ▶ Recall that for a martingale, $E[Z_n] = E[Z_1], \forall n$
- ▶ The theorem essentially says there is no strategy to have a positive expectation from a fair gambling game.

- $\{X_n, n = 0, 1, \dots\}$ and $E|X_n| < \infty, \forall n$ is called a martingale if

$$E[X_{n+1} \mid X_n, \dots, X_0] = X_n, \forall n$$

- $\{X_n, n = 0, 1, \dots\}$ and $E|X_n| < \infty, \forall n$ is called a martingale if

$$E[X_{n+1} \mid X_n, \dots, X_0] = X_n, \forall n$$

- It is called a supermartingale if

$$E[X_{n+1} \mid X_n, \dots, X_0] \leq X_n, \forall n$$

- ▶ $\{X_n, n = 0, 1, \dots\}$ and $E|X_n| < \infty, \forall n$ is called a martingale if

$$E[X_{n+1} \mid X_n, \dots, X_0] = X_n, \forall n$$

- ▶ It is called a supermartingale if

$$E[X_{n+1} \mid X_n, \dots, X_0] \leq X_n, \forall n$$

- ▶ It is called a submartingale if

$$E[X_{n+1} \mid X_n, \dots, X_0] \geq X_n, \forall n$$

- ▶ $\{X_n, n = 0, 1, \dots\}$ and $E|X_n| < \infty, \forall n$ is called a martingale if

$$E[X_{n+1} \mid X_n, \dots, X_0] = X_n, \forall n$$

- ▶ It is called a supermartingale if

$$E[X_{n+1} \mid X_n, \dots, X_0] \leq X_n, \forall n$$

- ▶ It is called a submartingale if

$$E[X_{n+1} \mid X_n, \dots, X_0] \geq X_n, \forall n$$

- ▶ Note that $E[X_{n+1}] \leq E[X_n]$ for supermartingales and $E[X_{n+1}] \geq E[X_n]$ for submartingales.

- ▶ $\{X_n, n = 0, 1, \dots\}$ and $E|X_n| < \infty, \forall n$ is called a martingale if

$$E[X_{n+1} \mid X_n, \dots, X_0] = X_n, \forall n$$

- ▶ It is called a supermartingale if

$$E[X_{n+1} \mid X_n, \dots, X_0] \leq X_n, \forall n$$

- ▶ It is called a submartingale if

$$E[X_{n+1} \mid X_n, \dots, X_0] \geq X_n, \forall n$$

- ▶ Note that $E[X_{n+1}] \leq E[X_n]$ for supermartingales and $E[X_{n+1}] \geq E[X_n]$ for submartingales.
- ▶ In the above, the conditioning random variables can be another sequence Y_i if Y_1, \dots, Y_n determine X_1, \dots, X_n

- ▶ A useful result is the martingale convergence theorem.

- ▶ A useful result is the martingale convergence theorem.

Martingale convergence theorem: If X_n is a martingale with $\sup_n E|X_n| < \infty$ then X_n converges almost surely to a rv X which will have finite expectation. A positive supermartingale also converges almost surely

- ▶ A useful result is the martingale convergence theorem.

Martingale convergence theorem: If X_n is a martingale with $\sup_n E|X_n| < \infty$ then X_n converges almost surely to a rv X which will have finite expectation. A positive supermartingale also converges almost surely

- ▶ If X_n are bounded, then the condition is always true and the almost sure convergence implies convergence in the mean.

- ▶ A useful result is the martingale convergence theorem.

Martingale convergence theorem: If X_n is a martingale with $\sup_n E|X_n| < \infty$ then X_n converges almost surely to a rv X which will have finite expectation. A positive supermartingale also converges almost surely

- ▶ If X_n are bounded, then the condition is always true and the almost sure convergence implies convergence in the mean.
- ▶ This is often useful in dealing with many sequences of random variables such as a stochastic algorithm.

Example

- ▶ We consider a simple algorithm for a two armed bandit problem.

Example

- ▶ We consider a simple algorithm for a two armed bandit problem.
- ▶ 2 arms. Response is binary (1 for reward).

Example

- ▶ We consider a simple algorithm for a two armed bandit problem.
- ▶ 2 arms. Response is binary (1 for reward).
- ▶ d_i - prob of reward for arm- i . Do not know d_i

Example

- ▶ We consider a simple algorithm for a two armed bandit problem.
- ▶ 2 arms. Response is binary (1 for reward).
- ▶ d_i - prob of reward for arm- i . Do not know d_i
- ▶ need to play and find which is better arm.

Example

- ▶ We consider a simple algorithm for a two armed bandit problem.
- ▶ 2 arms. Response is binary (1 for reward).
- ▶ d_i - prob of reward for arm- i . Do not know d_i
- ▶ need to play and find which is better arm.
- ▶ We choose arm-1 with prob $p(k)$ (and hence arm-2 with prob $(1 - p(k))$) at iteration k and update $p(k)$ based on the outcome.

- ▶ $p(k)$ denotes probability of choosing arm-1 at k and $b(k)$ denotes the response.

- ▶ $p(k)$ denotes probability of choosing arm-1 at k and $b(k)$ denotes the response.
- ▶ Our algorithm for the 2-armed bandit problem is

- ▶ $p(k)$ denotes probability of choosing arm-1 at k and $b(k)$ denotes the response.
- ▶ Our algorithm for the 2-armed bandit problem is

$$p(k+1) = p(k) + \lambda(1 - p(k)) \quad \text{if arm 1 chosen, } b(k) = 1$$

- ▶ $p(k)$ denotes probability of choosing arm-1 at k and $b(k)$ denotes the response.
- ▶ Our algorithm for the 2-armed bandit problem is

$$\begin{aligned} p(k+1) &= p(k) + \lambda(1 - p(k)) \quad \text{if arm 1 chosen, } b(k) = 1 \\ &= p(k) - \lambda p(k) \quad \text{if arm 2 is chosen and } b(k) = 1 \end{aligned}$$

- ▶ $p(k)$ denotes probability of choosing arm-1 at k and $b(k)$ denotes the response.
- ▶ Our algorithm for the 2-armed bandit problem is

$$\begin{aligned} p(k+1) &= p(k) + \lambda(1 - p(k)) \quad \text{if arm 1 chosen, } b(k) = 1 \\ &= p(k) - \lambda p(k) \quad \text{if arm 2 is chosen and } b(k) = 1 \\ &= p(k) \quad \text{if } b(k) = 0 \end{aligned}$$

- ▶ $p(k)$ denotes probability of choosing arm-1 at k and $b(k)$ denotes the response.
- ▶ Our algorithm for the 2-armed bandit problem is

$$\begin{aligned} p(k+1) &= p(k) + \lambda(1 - p(k)) \quad \text{if arm 1 chosen, } b(k) = 1 \\ &= p(k) - \lambda p(k) \quad \text{if arm 2 is chosen and } b(k) = 1 \\ &= p(k) \quad \text{if } b(k) = 0 \end{aligned}$$

- ▶ We want to know whether the algorithm converges.

► The algorithm is

$$\begin{aligned} p(k+1) &= p(k) + \lambda(1 - p(k)) \quad \text{if arm 1 chosen, } b(k) = 1 \\ &= p(k) - \lambda p(k) \quad \text{if arm 2 is chosen and } b(k) = 1 \\ &= p(k) \quad \text{if } b(k) = 0 \end{aligned}$$

- ▶ The algorithm is

$$\begin{aligned} p(k+1) &= p(k) + \lambda(1 - p(k)) \quad \text{if arm 1 chosen, } b(k) = 1 \\ &= p(k) - \lambda p(k) \quad \text{if arm 2 is chosen and } b(k) = 1 \\ &= p(k) \quad \text{if } b(k) = 0 \end{aligned}$$

- ▶ We get

- The algorithm is

$$\begin{aligned} p(k+1) &= p(k) + \lambda(1 - p(k)) \quad \text{if arm 1 chosen, } b(k) = 1 \\ &= p(k) - \lambda p(k) \quad \text{if arm 2 is chosen and } b(k) = 1 \\ &= p(k) \quad \text{if } b(k) = 0 \end{aligned}$$

- We get

$$E[p(k+1) - p(k) | p(k)]$$

- The algorithm is

$$\begin{aligned} p(k+1) &= p(k) + \lambda(1 - p(k)) \quad \text{if arm 1 chosen, } b(k) = 1 \\ &= p(k) - \lambda p(k) \quad \text{if arm 2 is chosen and } b(k) = 1 \\ &= p(k) \quad \text{if } b(k) = 0 \end{aligned}$$

- We get

$$\begin{aligned} E[p(k+1) - p(k) | p(k)] \\ = \lambda(1 - p(k)) \Pr[b(k) = 1, \text{arm 1} \mid p(k)] \end{aligned}$$

- The algorithm is

$$\begin{aligned} p(k+1) &= p(k) + \lambda(1 - p(k)) \quad \text{if arm 1 chosen, } b(k) = 1 \\ &= p(k) - \lambda p(k) \quad \text{if arm 2 is chosen and } b(k) = 1 \\ &= p(k) \quad \text{if } b(k) = 0 \end{aligned}$$

- We get

$$\begin{aligned} E[p(k+1) - p(k) | p(k)] \\ &= \lambda(1 - p(k)) \Pr[b(k) = 1, \text{arm 1} \mid p(k)] \\ &\quad - \lambda p(k) \Pr[b(k) = 1, \text{arm 2} \mid p(k)] \end{aligned}$$

- The algorithm is

$$\begin{aligned} p(k+1) &= p(k) + \lambda(1 - p(k)) \quad \text{if arm 1 chosen, } b(k) = 1 \\ &= p(k) - \lambda p(k) \quad \text{if arm 2 is chosen and } b(k) = 1 \\ &= p(k) \quad \text{if } b(k) = 0 \end{aligned}$$

- We get

$$\begin{aligned} &E[p(k+1) - p(k) | p(k)] \\ &= \lambda(1 - p(k)) Pr[b(k) = 1, \text{arm 1} | p(k)] \\ &\quad - \lambda p(k) Pr[b(k) = 1, \text{arm 2} | p(k)] \\ &= \lambda(1 - p(k)) Pr[b(k) = 1 | \text{arm 1}, p(k)] Pr[\text{arm 1} | p(k)] \end{aligned}$$

- The algorithm is

$$\begin{aligned} p(k+1) &= p(k) + \lambda(1 - p(k)) \text{ if arm 1 chosen, } b(k) = 1 \\ &= p(k) - \lambda p(k) \text{ if arm 2 is chosen and } b(k) = 1 \\ &= p(k) \text{ if } b(k) = 0 \end{aligned}$$

- We get

$$\begin{aligned} &E[p(k+1) - p(k) | p(k)] \\ &= \lambda(1 - p(k)) Pr[b(k) = 1, \text{arm 1} | p(k)] \\ &\quad - \lambda p(k) Pr[b(k) = 1, \text{arm 2} | p(k)] \\ &= \lambda(1 - p(k)) Pr[b(k) = 1 | \text{arm 1}, p(k)] Pr[\text{arm 1} | p(k)] \\ &\quad - \lambda p(k) Pr[b(k) = 1 | \text{arm 2}, p(k)] Pr[\text{arm 2} | p(k)] \end{aligned}$$

► This gives us

► This gives us

$$\begin{aligned} E[p(k+1) - p(k) | p(k)] &= \lambda(1 - p(k)) d_1 p(k) \\ &\quad - \lambda p(k) d_2 (1 - p(k)) \end{aligned}$$

► This gives us

$$\begin{aligned} E[p(k+1) - p(k)|p(k)] &= \lambda(1 - p(k)) d_1 p(k) \\ &\quad - \lambda p(k) d_2 (1 - p(k)) \\ &= \lambda p(k)(1 - p(k)) (d_1 - d_2) \end{aligned}$$

► This gives us

$$\begin{aligned} E[p(k+1) - p(k)|p(k)] &= \lambda(1 - p(k)) d_1 p(k) \\ &\quad - \lambda p(k) d_2 (1 - p(k)) \\ &= \lambda p(k)(1 - p(k)) (d_1 - d_2) \\ &\geq 0, \quad \text{if } d_1 > d_2 \end{aligned}$$

► This gives us

$$\begin{aligned} E[p(k+1) - p(k) | p(k)] &= \lambda(1 - p(k)) d_1 p(k) \\ &\quad - \lambda p(k) d_2 (1 - p(k)) \\ &= \lambda p(k)(1 - p(k)) (d_1 - d_2) \\ &\geq 0, \quad \text{if } d_1 > d_2 \end{aligned}$$

$$\Rightarrow E[p(k+1) | p(k)] \geq p(k)$$

► This gives us

$$\begin{aligned} E[p(k+1) - p(k) | p(k)] &= \lambda(1 - p(k)) d_1 p(k) \\ &\quad - \lambda p(k) d_2 (1 - p(k)) \\ &= \lambda p(k)(1 - p(k)) (d_1 - d_2) \\ &\geq 0, \quad \text{if } d_1 > d_2 \end{aligned}$$

$$\Rightarrow E[p(k+1) | p(k)] \geq p(k) \Rightarrow E[p(k+1)] \geq E[p(k)], \forall k$$

- This gives us

$$\begin{aligned} E[p(k+1) - p(k) | p(k)] &= \lambda(1 - p(k)) d_1 p(k) \\ &\quad - \lambda p(k) d_2 (1 - p(k)) \\ &= \lambda p(k)(1 - p(k)) (d_1 - d_2) \\ &\geq 0, \quad \text{if } d_1 > d_2 \end{aligned}$$

$$\Rightarrow E[p(k+1) | p(k)] \geq p(k) \Rightarrow E[p(k+1)] \geq E[p(k)], \forall k$$

- This also shows $p(k)$ is a submartingale.

- This gives us

$$\begin{aligned} E[p(k+1) - p(k) | p(k)] &= \lambda(1 - p(k)) d_1 p(k) \\ &\quad - \lambda p(k) d_2 (1 - p(k)) \\ &= \lambda p(k)(1 - p(k)) (d_1 - d_2) \\ &\geq 0, \quad \text{if } d_1 > d_2 \end{aligned}$$

$$\Rightarrow E[p(k+1) | p(k)] \geq p(k) \Rightarrow E[p(k+1)] \geq E[p(k)], \forall k$$

- This also shows $p(k)$ is a submartingale.
- Here, $p(k)$ is bounded and $1 - p(k)$ is a supermartingale.

- This gives us

$$\begin{aligned} E[p(k+1) - p(k) | p(k)] &= \lambda(1 - p(k)) d_1 p(k) \\ &\quad - \lambda p(k) d_2 (1 - p(k)) \\ &= \lambda p(k)(1 - p(k)) (d_1 - d_2) \\ &\geq 0, \quad \text{if } d_1 > d_2 \end{aligned}$$

$$\Rightarrow E[p(k+1) | p(k)] \geq p(k) \Rightarrow E[p(k+1)] \geq E[p(k)], \forall k$$

- This also shows $p(k)$ is a submartingale.
- Here, $p(k)$ is bounded and $1 - p(k)$ is a supermartingale.
- So, we can conclude, the algorithm converges almost surely

- ▶ We considered martingales as an example of discrete-time continuous-state processes

- ▶ We considered martingales as an example of discrete-time continuous-state processes
- ▶ Stochastic iterative algorithms generate such processes.

- ▶ We considered martingales as an example of discrete-time continuous-state processes
- ▶ Stochastic iterative algorithms generate such processes.
- ▶ Martingales can be useful in analyzing convergence of many stochastic algorithms

- ▶ We considered martingales as an example of discrete-time continuous-state processes
- ▶ Stochastic iterative algorithms generate such processes.
- ▶ Martingales can be useful in analyzing convergence of many stochastic algorithms
- ▶ While we mentioned only discrete-time martingales, one can similarly have continuous-time martingales that satisfy

$$E[X(t)|(X(s'), 0 \leq s' \leq s < t] = X(s)$$