

Recap: Random Variables

- ▶ Given a probability space (Ω, \mathcal{F}, P) , a random variable is a real-valued function on Ω .

Recap: Random Variables

- ▶ Given a probability space (Ω, \mathcal{F}, P) , a random variable is a real-valued function on Ω .
- ▶ It essentially results in an induced probability space

$$(\Omega, \mathcal{F}, P) \xrightarrow{X} (\mathbb{R}, \mathcal{B}, P_X)$$

where \mathcal{B} is the Borel σ -algebra and

$$P_X(B) = P[X \in B] = P(\{\omega \in \Omega : X(\omega) \in B\})$$

Recap: Distribution function of a random variable

- ▶ Let X be a random variable. Its distribution function, $F_X : \mathfrak{R} \rightarrow \mathfrak{R}$, is defined by

$$F_X(x) = P[X \leq x] = P(\{\omega \in \Omega : X(\omega) \leq x\})$$

Recap: Distribution function of a random variable

- ▶ Let X be a random variable. Its distribution function, $F_X : \Re \rightarrow \Re$, is defined by

$$F_X(x) = P[X \leq x] = P(\{\omega \in \Omega : X(\omega) \leq x\})$$

- ▶ The distribution function, F_X , completely specifies the probability measure, P_X .

Recap: Properties of distribution function

► The distribution function satisfies

1. $0 \leq F_X(x) \leq 1, \forall x$
2. $F_X(-\infty) = 0; F_X(\infty) = 1$
3. F_X is non-decreasing: $x_1 \leq x_2 \Rightarrow F_X(x_1) \leq F_X(x_2)$
4. F_X is right continuous and has left-hand limits.

Recap: Properties of distribution function

- ▶ The distribution function satisfies
 1. $0 \leq F_X(x) \leq 1, \forall x$
 2. $F_X(-\infty) = 0; F_X(\infty) = 1$
 3. F_X is non-decreasing: $x_1 \leq x_2 \Rightarrow F_X(x_1) \leq F_X(x_2)$
 4. F_X is right continuous and has left-hand limits.
- ▶ Any real-valued function of a real variable satisfying the above four properties would be a distribution function of some random variable.

Recap: Properties of distribution function

- ▶ The distribution function satisfies
 1. $0 \leq F_X(x) \leq 1, \forall x$
 2. $F_X(-\infty) = 0; F_X(\infty) = 1$
 3. F_X is non-decreasing: $x_1 \leq x_2 \Rightarrow F_X(x_1) \leq F_X(x_2)$
 4. F_X is right continuous and has left-hand limits.
- ▶ Any real-valued function of a real variable satisfying the above four properties would be a distribution function of some random variable.
- ▶ We also have
$$F_X(x^+) - F_X(x^-) = F_X(x) - F_X(x^-) = P[X = x]$$

Recap: Properties of distribution function

- ▶ The distribution function satisfies
 1. $0 \leq F_X(x) \leq 1, \forall x$
 2. $F_X(-\infty) = 0; F_X(\infty) = 1$
 3. F_X is non-decreasing: $x_1 \leq x_2 \Rightarrow F_X(x_1) \leq F_X(x_2)$
 4. F_X is right continuous and has left-hand limits.
- ▶ Any real-valued function of a real variable satisfying the above four properties would be a distribution function of some random variable.
- ▶ We also have
$$F_X(x^+) - F_X(x^-) = F_X(x) - F_X(x^-) = P[X = x]$$
$$P[a < X \leq b] = F_X(b) - F_X(a).$$

Recap: Discrete Random Variable

- ▶ A random variable X is said to be discrete if it takes only finitely many or countably infinitely many distinct values.

Recap: Discrete Random Variable

- ▶ A random variable X is said to be discrete if it takes only finitely many or countably infinitely many distinct values.
- ▶ Let $X \in \{x_1, x_2, \dots\}$

Recap: Discrete Random Variable

- ▶ A random variable X is said to be discrete if it takes only finitely many or countably infinitely many distinct values.
- ▶ Let $X \in \{x_1, x_2, \dots\}$
- ▶ Its distribution function, F_X is a stair-case function with jump discontinuities at each x_i and the magnitude of the jump at x_i is equal to $P[X = x_i]$

Recap: probability mass function

- ▶ Let $X \in \{x_1, x_2, \dots\}$.

Recap: probability mass function

- ▶ Let $X \in \{x_1, x_2, \dots\}$.
- ▶ The probability mass function (pmf) of X is defined by

$$f_X(x_i) = P[X = x_i]; \quad f_X(x) = 0, \quad \text{for all other } x$$

Recap: probability mass function

- ▶ Let $X \in \{x_1, x_2, \dots\}$.
- ▶ The probability mass function (pmf) of X is defined by

$$f_X(x_i) = P[X = x_i]; \quad f_X(x) = 0, \quad \text{for all other } x$$

- ▶ It satisfies
 1. $f_X(x) \geq 0, \forall x$ and $f_X(x) = 0$ if $x \neq x_i$ for some i
 2. $\sum_i f_X(x_i) = 1$

Recap: probability mass function

- ▶ Let $X \in \{x_1, x_2, \dots\}$.
- ▶ The probability mass function (pmf) of X is defined by

$$f_X(x_i) = P[X = x_i]; \quad f_X(x) = 0, \quad \text{for all other } x$$

- ▶ It satisfies
 1. $f_X(x) \geq 0, \forall x$ and $f_X(x) = 0$ if $x \neq x_i$ for some i
 2. $\sum_i f_X(x_i) = 1$

- ▶ We have

$$F_X(x) = \sum_{i: x_i \leq x} f_X(x_i)$$
$$f_X(x) = F_X(x) - F_X(x^-)$$

Recap: probability mass function

- ▶ Let $X \in \{x_1, x_2, \dots\}$.
- ▶ The probability mass function (pmf) of X is defined by

$$f_X(x_i) = P[X = x_i]; \quad f_X(x) = 0, \quad \text{for all other } x$$

- ▶ It satisfies
 1. $f_X(x) \geq 0, \forall x$ and $f_X(x) = 0$ if $x \neq x_i$ for some i
 2. $\sum_i f_X(x_i) = 1$

- ▶ We have

$$F_X(x) = \sum_{i: x_i \leq x} f_X(x_i)$$
$$f_X(x) = F_X(x) - F_X(x^-)$$

- ▶ We can calculate the probability of any event as

$$P[X \in B] = \sum_{\substack{i: \\ x_i \in B}} f_X(x_i)$$

Recap: continuous random variable

- ▶ X is said to be a continuous random variable if there exists a function $f_X : \Re \rightarrow \Re$ satisfying

$$F_X(x) = \int_{-\infty}^x f_X(x) \, dx$$

The f_X is called the probability density function.

Recap: continuous random variable

- ▶ X is said to be a continuous random variable if there exists a function $f_X : \Re \rightarrow \Re$ satisfying

$$F_X(x) = \int_{-\infty}^x f_X(x) \, dx$$

The f_X is called the probability density function.

- ▶ Same as saying F_X is absolutely continuous.

Recap: continuous random variable

- ▶ X is said to be a continuous random variable if there exists a function $f_X : \Re \rightarrow \Re$ satisfying

$$F_X(x) = \int_{-\infty}^x f_X(x) dx$$

The f_X is called the probability density function.

- ▶ Same as saying F_X is absolutely continuous.
- ▶ Since F_X is continuous here, we have

$$P[X = x] = F_X(x) - F_X(x^-) = 0, \quad \forall x$$

Recap: continuous random variable

- ▶ X is said to be a continuous random variable if there exists a function $f_X : \Re \rightarrow \Re$ satisfying

$$F_X(x) = \int_{-\infty}^x f_X(x) dx$$

The f_X is called the probability density function.

- ▶ Same as saying F_X is absolutely continuous.
- ▶ Since F_X is continuous here, we have

$$P[X = x] = F_X(x) - F_X(x^-) = 0, \forall x$$

- ▶ A continuous rv takes uncountably many distinct values. However, not every rv that takes uncountably many values is a continuous rv

Recap: probability density function

- ▶ The pdf of a continuous rv is defined to be the f_X that satisfies

$$F_X(x) = \int_{-\infty}^x f_X(t) dt, \quad \forall x$$

Recap: probability density function

- ▶ The pdf of a continuous rv is defined to be the f_X that satisfies

$$F_X(x) = \int_{-\infty}^x f_X(t) dt, \quad \forall x$$

- ▶ It satisfies

1. $f_X(x) \geq 0, \quad \forall x$
2. $\int_{-\infty}^{\infty} f_X(t) dt = 1$

Recap: probability density function

- ▶ The pdf of a continuous rv is defined to be the f_X that satisfies

$$F_X(x) = \int_{-\infty}^x f_X(t) dt, \quad \forall x$$

- ▶ It satisfies

1. $f_X(x) \geq 0, \quad \forall x$

2. $\int_{-\infty}^{\infty} f_X(t) dt = 1$

- ▶ We can, in principle, compute probability of any event as

$$P[X \in B] = \int_B f_X(t) dt, \quad \forall B \in \mathcal{B}$$

Recap: probability density function

- ▶ The pdf of a continuous rv is defined to be the f_X that satisfies

$$F_X(x) = \int_{-\infty}^x f_X(t) dt, \quad \forall x$$

- ▶ It satisfies

1. $f_X(x) \geq 0, \quad \forall x$

2. $\int_{-\infty}^{\infty} f_X(t) dt = 1$

- ▶ We can, in principle, compute probability of any event as

$$P[X \in B] = \int_B f_X(t) dt, \quad \forall B \in \mathcal{B}$$

- ▶ In particular,

$$P[a \leq X \leq b] = \int_a^b f_X(t) dt$$

Recap: Function of a random variable

- ▶ If X is a random variable and $g : \mathfrak{R} \rightarrow \mathfrak{R}$ is a function, then $Y = g(X)$ is a random variable.

Recap: Function of a random variable

- ▶ If X is a random variable and $g : \mathfrak{R} \rightarrow \mathfrak{R}$ is a function, then $Y = g(X)$ is a random variable.
- ▶ More formally, Y is a random variable if g is a Borel measurable function.

Recap: Function of a random variable

- ▶ If X is a random variable and $g : \mathfrak{R} \rightarrow \mathfrak{R}$ is a function, then $Y = g(X)$ is a random variable.
- ▶ More formally, Y is a random variable if g is a Borel measurable function.
- ▶ We can determine distribution of Y given the function g and the distribution of X

Recap

- ▶ Let X be a rv and let $Y = g(X)$.

Recap

- ▶ Let X be a rv and let $Y = g(X)$.
- ▶ The distribution function of Y is given by

$$\begin{aligned} F_Y(y) &= P[g(X) \leq y] \\ &= P[X \in \{z : g(z) \leq y\}] \end{aligned}$$

Recap

- ▶ Let X be a rv and let $Y = g(X)$.
- ▶ The distribution function of Y is given by

$$\begin{aligned}F_Y(y) &= P[g(X) \leq y] \\&= P[X \in \{z : g(z) \leq y\}]\end{aligned}$$

- ▶ This probability can be obtained from distribution of X .

Recap

- ▶ Suppose X is a discrete rv with $X \in \{x_1, x_2, \dots\}$.

Recap

- ▶ Suppose X is a discrete rv with $X \in \{x_1, x_2, \dots\}$.
- ▶ Suppose $Y = g(X)$.

Recap

- ▶ Suppose X is a discrete rv with $X \in \{x_1, x_2, \dots\}$.
- ▶ Suppose $Y = g(X)$.
- ▶ Then Y is also discrete and $Y \in \{g(x_1), g(x_2), \dots\}$.

Recap

- ▶ Suppose X is a discrete rv with $X \in \{x_1, x_2, \dots\}$.
- ▶ Suppose $Y = g(X)$.
- ▶ Then Y is also discrete and $Y \in \{g(x_1), g(x_2), \dots\}$.
- ▶ We can find the pmf of Y as

$$\begin{aligned} f_Y(y) &= p[Y = y] = P[g(X) = y] \\ &= P[X \in \{x_i : g(x_i) = y\}] \\ &= \sum_{\substack{i: \\ g(x_i)=y}} f_X(x_i) \end{aligned}$$

Recap

- ▶ Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable with $g'(x) > 0, \forall x$ or $g'(x) < 0, \forall x$.

Recap

- ▶ Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be differentiable with $g'(x) > 0, \forall x$ or $g'(x) < 0, \forall x$.
- ▶ Let X be a continuous rv and let $Y = g(X)$.

Recap

- ▶ Let $g : \Re \rightarrow \Re$ be differentiable with $g'(x) > 0, \forall x$ or $g'(x) < 0, \forall x$.
- ▶ Let X be a continuous rv and let $Y = g(X)$.
- ▶ Then Y is a continuous rv with pdf

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|, \quad a \leq y \leq b$$

where $a = \min(g(\infty), g(-\infty))$ and
 $b = \max(g(\infty), g(-\infty))$

Recap

- ▶ Let $g : \Re \rightarrow \Re$ be differentiable with $g'(x) > 0, \forall x$ or $g'(x) < 0, \forall x$.
- ▶ Let X be a continuous rv and let $Y = g(X)$.
- ▶ Then Y is a continuous rv with pdf

$$f_Y(y) = f_X(g^{-1}(y)) \left| \frac{d}{dy} g^{-1}(y) \right|, \quad a \leq y \leq b$$

where $a = \min(g(\infty), g(-\infty))$ and
 $b = \max(g(\infty), g(-\infty))$

- ▶ This theorem is useful in some cases to find the densities of functions of continuous random variables

Recap: Expectation

Recap: Expectation

- ▶ Let X be a discrete rv with $X \in \{x_1, x_2, \dots\}$. Then

$$E[X] = \sum_i x_i f_X(x_i)$$

Recap: Expectation

- ▶ Let X be a discrete rv with $X \in \{x_1, x_2, \dots\}$. Then

$$E[X] = \sum_i x_i f_X(x_i)$$

- ▶ If X is a continuous random variable with pdf, f_X ,

$$E[X] = \int_{-\infty}^{\infty} x f_X(x) dx$$

Recap: Expectation

- ▶ Let X be a discrete rv with $X \in \{x_1, x_2, \dots\}$. Then

$$E[X] = \sum_i x_i f_X(x_i)$$

- ▶ If X is a continuous random variable with pdf, f_X ,

$$E[X] = \int_{-\infty}^{\infty} x f_X(x) dx$$

- ▶ Sometimes we use the following notation to denote expectation of both kinds of rv

$$E[X] = \int_{-\infty}^{\infty} x dF_X(x)$$

Recap: Expectation

- ▶ Let X be a discrete rv with $X \in \{x_1, x_2, \dots\}$. Then

$$E[X] = \sum_i x_i f_X(x_i)$$

- ▶ If X is a continuous random variable with pdf, f_X ,

$$E[X] = \int_{-\infty}^{\infty} x f_X(x) dx$$

- ▶ Sometimes we use the following notation to denote expectation of both kinds of rv

$$E[X] = \int_{-\infty}^{\infty} x dF_X(x)$$

- ▶ We take the expectation to exist when the sum or integral above is absolutely convergent

Recap: Expectation

- ▶ Let X be a discrete rv with $X \in \{x_1, x_2, \dots\}$. Then

$$E[X] = \sum_i x_i f_X(x_i)$$

- ▶ If X is a continuous random variable with pdf, f_X ,

$$E[X] = \int_{-\infty}^{\infty} x f_X(x) dx$$

- ▶ Sometimes we use the following notation to denote expectation of both kinds of rv

$$E[X] = \int_{-\infty}^{\infty} x dF_X(x)$$

- ▶ We take the expectation to exist when the sum or integral above is absolutely convergent
- ▶ Note that expectation is defined for all random variables

Recap: Expectation of a function of a random variable

- ▶ Let X be a rv and let $Y = g(X)$. Then,

Recap: Expectation of a function of a random variable

- ▶ Let X be a rv and let $Y = g(X)$. Then,
- ▶ $EY = \int y \, dF_Y(y) = \int g(x) \, dF_X(x)$

Recap: Expectation of a function of a random variable

- ▶ Let X be a rv and let $Y = g(X)$. Then,
- ▶ $EY = \int y \, dF_Y(y) = \int g(x) \, dF_X(x)$
- ▶ That is, if X is discrete, then

$$EY = \sum_j y_j f_Y(y_j) = \sum_i g(x_i) f_X(x_i)$$

Recap: Expectation of a function of a random variable

- ▶ Let X be a rv and let $Y = g(X)$. Then,
- ▶ $EY = \int y \, dF_Y(y) = \int g(x) \, dF_X(x)$
- ▶ That is, if X is discrete, then

$$EY = \sum_j y_j f_Y(y_j) = \sum_i g(x_i) f_X(x_i)$$

- ▶ If X and Y are continuous

$$EY = \int y f_Y(y) \, dy = \int g(x) f_X(x) \, dx$$

Recap: Expectation of a function of a random variable

- ▶ Let X be a rv and let $Y = g(X)$. Then,
- ▶ $EY = \int y dF_Y(y) = \int g(x) dF_X(x)$
- ▶ That is, if X is discrete, then

$$EY = \sum_j y_j f_Y(y_j) = \sum_i g(x_i) f_X(x_i)$$

- ▶ If X and Y are continuous

$$EY = \int y f_Y(y) dy = \int g(x) f_X(x) dx$$

- ▶ This is true for all rv's.

Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

► If $X \geq 0$ then $EX \geq 0$

Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

- ▶ If $X \geq 0$ then $EX \geq 0$
- ▶ $E[b] = b$ where b is a constant

Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

- ▶ If $X \geq 0$ then $EX \geq 0$
- ▶ $E[b] = b$ where b is a constant
- ▶ $E[ag(X)] = aE[g(X)]$ where a is a constant

Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

- ▶ If $X \geq 0$ then $EX \geq 0$
- ▶ $E[b] = b$ where b is a constant
- ▶ $E[ag(X)] = aE[g(X)]$ where a is a constant
- ▶ $E[aX + b] = aE[X] + b$ where a, b are constants.

Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

- ▶ If $X \geq 0$ then $EX \geq 0$
- ▶ $E[b] = b$ where b is a constant
- ▶ $E[ag(X)] = aE[g(X)]$ where a is a constant
- ▶ $E[aX + b] = aE[X] + b$ where a, b are constants.
- ▶ $E[ag_1(X) + bg_2(X)] = aE[g_1(X)] + bE[g_2(X)]$

Recap: Properties of Expectation

$$E[g(X)] = \sum_i g(x_i) f_X(x_i) \quad \text{or} \quad E[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$$

- ▶ If $X \geq 0$ then $EX \geq 0$
- ▶ $E[b] = b$ where b is a constant
- ▶ $E[ag(X)] = aE[g(X)]$ where a is a constant
- ▶ $E[aX + b] = aE[X] + b$ where a, b are constants.
- ▶ $E[ag_1(X) + bg_2(X)] = aE[g_1(X)] + bE[g_2(X)]$
- ▶ $E[(X - c)^2] \geq E[(X - EX)^2], \forall c$

Recap: Variance of random variable



$$\text{Var}(X) = E[(X - EX)^2] = E[X^2] - (EX)^2$$

Recap: Variance of random variable



$$\text{Var}(X) = E[(X - EX)^2] = E[X^2] - (EX)^2$$



Properties of Variance:

Recap: Variance of random variable



$$\text{Var}(X) = E[(X - EX)^2] = E[X^2] - (EX)^2$$

▶ Properties of Variance:

▶ $\text{Var}(X) \geq 0$

Recap: Variance of random variable



$$\text{Var}(X) = E[(X - EX)^2] = E[X^2] - (EX)^2$$

▶ Properties of Variance:

- ▶ $\text{Var}(X) \geq 0$
- ▶ $\text{Var}(X + c) = \text{Var}(X)$

Recap: Variance of random variable



$$\text{Var}(X) = E[(X - EX)^2] = E[X^2] - (EX)^2$$

► Properties of Variance:

- $\text{Var}(X) \geq 0$
- $\text{Var}(X + c) = \text{Var}(X)$
- $\text{Var}(cX) = c^2 \text{Var}(X)$

Recap: Moments of a random variable

- ▶ The k^{th} (order) moment of X is

$$m_k = E[X^k] = \int x^k dF_X(x)$$

Recap: Moments of a random variable

- ▶ The k^{th} (order) moment of X is

$$m_k = E[X^k] = \int x^k dF_X(x)$$

- ▶ The k^{th} central moment of X is

$$s_k = E[(X - EX)^k] = \int (x - EX)^k dF_X(x)$$

Recap: Moments of a random variable

- ▶ The k^{th} (order) moment of X is

$$m_k = E[X^k] = \int x^k dF_X(x)$$

- ▶ The k^{th} central moment of X is

$$s_k = E[(X - EX)^k] = \int (x - EX)^k dF_X(x)$$

- ▶ If moment of order k is finite then so is moment of order s for $s < k$.

Recap: Moment Generating function

- The moment generating function – $M_X : \Re \rightarrow \Re$

$$M_X(t) = Ee^{tX} = \sum_i e^{tx_i} f_X(x_i) \quad \text{or} \quad \int e^{tx} f_X(x) dx, \quad t \in \Re$$

Recap: Moment Generating function

- ▶ The moment generating function – $M_X : \Re \rightarrow \Re$

$$M_X(t) = Ee^{tX} = \sum_i e^{tx_i} f_X(x_i) \quad \text{or} \quad \int e^{tx} f_X(x) dx, \quad t \in \Re$$

- ▶ We say the mgf exists if $E[e^{tX}] < \infty$ for t in some interval around zero

Recap: Moment Generating function

- ▶ The moment generating function – $M_X : \Re \rightarrow \Re$

$$M_X(t) = Ee^{tX} = \sum_i e^{tx_i} f_X(x_i) \quad \text{or} \quad \int e^{tx} f_X(x) dx, \quad t \in \Re$$

- ▶ We say the mgf exists if $E[e^{tX}] < \infty$ for t in some interval around zero
- ▶ If $M_X(t)$ exists (for $t \in [-a, a]$ for some $a > 0$) then all its derivatives also exist and

$$\left. \frac{d^k M_X(t)}{dt^k} \right|_{t=0} = E[X^k]$$

Generating function

- For $X \in \{0, 1, 2, \dots\}$ the (probability) generating function of X is defined by

$$P_X(s) = \sum_{k=0}^{\infty} f_X(k) s^k, \quad s \in \mathfrak{R}$$

Generating function

- ▶ For $X \in \{0, 1, 2, \dots\}$ the (probability) generating function of X is defined by

$$P_X(s) = \sum_{k=0}^{\infty} f_X(k) s^k, \quad s \in \mathfrak{R}$$

- ▶ We get the pmf from it as

$$f_X(0) = P_X(0); \quad f_X(1) = \frac{P'_X(0)}{1!}; \quad f_X(2) = \frac{P''_X(0)}{2!}$$

Generating function

- ▶ For $X \in \{0, 1, 2, \dots\}$ the (probability) generating function of X is defined by

$$P_X(s) = \sum_{k=0}^{\infty} f_X(k)s^k, \quad s \in \mathfrak{R}$$

- ▶ We get the pmf from it as

$$f_X(0) = P_X(0); \quad f_X(1) = \frac{P'_X(0)}{1!}; \quad f_X(2) = \frac{P''_X(0)}{2!}$$

- ▶ We can also get the moments:

$$P'_X(1) = EX, \quad P''_X(1) = E[X(X-1)]$$

quantiles of a distribution

- ▶ Let $p \in (0, 1)$. The number $x \in \Re$ that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad p[X \geq x] \geq 1 - p$$

is called the quantile of order p or the $100p^{th}$ percentile of rv X .

quantiles of a distribution

- ▶ Let $p \in (0, 1)$. The number $x \in \Re$ that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad p[X \geq x] \geq 1 - p$$

is called the quantile of order p or the $100p^{th}$ percentile of rv X .

- ▶ If x is quantile of order p , it satisfies

$$p \leq F_X(x) \leq p + P[X = x]$$

quantiles of a distribution

- ▶ Let $p \in (0, 1)$. The number $x \in \Re$ that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad p[X \geq x] \geq 1 - p$$

is called the quantile of order p or the $100p^{th}$ percentile of rv X .

- ▶ If x is quantile of order p , it satisfies

$$p \leq F_X(x) \leq p + P[X = x]$$

- ▶ For a given p there can be multiple values for x to satisfy the above.

quantiles of a distribution

- ▶ Let $p \in (0, 1)$. The number $x \in \Re$ that satisfies

$$P[X \leq x] \geq p \quad \text{and} \quad p[X \geq x] \geq 1 - p$$

is called the quantile of order p or the $100p^{th}$ percentile of rv X .

- ▶ If x is quantile of order p , it satisfies

$$p \leq F_X(x) \leq p + P[X = x]$$

- ▶ For a given p there can be multiple values for x to satisfy the above.
- ▶ For $p = 0.5$, it is called the median.

Recap: some moment inequalities

- ▶ **Markov inequality:** For a non-negative function, g ,

Recap: some moment inequalities

- **Markov inequality:** For a non-negative function, g ,

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}$$

Recap: some moment inequalities

- ▶ **Markov inequality:** For a non-negative function, g ,

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}$$

- ▶ A specific instance of this is

$$P[|X| > c] \leq \frac{E[|X|^k]}{c^k}$$

Recap: some moment inequalities

- ▶ **Markov inequality:** For a non-negative function, g ,

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}$$

- ▶ A specific instance of this is

$$P[|X| > c] \leq \frac{E[|X|^k]}{c^k}$$

- ▶ **Chebyshev inequality**

$$P[|X - EX| > c] \leq \frac{\text{Var}(X)}{c^2}$$

Recap: some moment inequalities

- ▶ **Markov inequality:** For a non-negative function, g ,

$$P[g(X) > c] \leq \frac{E[g(X)]}{c}$$

- ▶ A specific instance of this is

$$P[|X| > c] \leq \frac{E[|X|^k]}{c^k}$$

- ▶ **Chebyshev inequality**

$$P[|X - EX| > c] \leq \frac{\text{Var}(X)}{c^2}$$

- ▶ With $EX = \mu$ and $\text{Var}(X) = \sigma^2$, we get

$$P[|X - \mu| > k\sigma] \leq \frac{1}{k^2}$$

Recap: A pair of random variables

- ▶ Let X, Y be random variables on the probability space (Ω, \mathcal{F}, P)

Recap: A pair of random variables

- ▶ Let X, Y be random variables on the probability space (Ω, \mathcal{F}, P)
- ▶ We can think of X, Y together as a vector-valued function mapping Ω to \mathbb{R}^2 .

Recap: A pair of random variables

- ▶ Let X, Y be random variables on the probability space (Ω, \mathcal{F}, P)
- ▶ We can think of X, Y together as a vector-valued function mapping Ω to \mathbb{R}^2 .
- ▶ This gives rise to a new probability space $(\mathbb{R}^2, \mathcal{B}^2, P_{XY})$ with P_{XY} given by

Recap: A pair of random variables

- ▶ Let X, Y be random variables on the probability space (Ω, \mathcal{F}, P)
- ▶ We can think of X, Y together as a vector-valued function mapping Ω to \mathbb{R}^2 .
- ▶ This gives rise to a new probability space $(\mathbb{R}^2, \mathcal{B}^2, P_{XY})$ with P_{XY} given by

$$P_{XY}(B) = P[(X, Y) \in B], \forall B \in \mathcal{B}^2$$

Recap: A pair of random variables

- ▶ Let X, Y be random variables on the probability space (Ω, \mathcal{F}, P)
- ▶ We can think of X, Y together as a vector-valued function mapping Ω to \mathbb{R}^2 .
- ▶ This gives rise to a new probability space $(\mathbb{R}^2, \mathcal{B}^2, P_{XY})$ with P_{XY} given by

$$\begin{aligned} P_{XY}(B) &= P[(X, Y) \in B], \forall B \in \mathcal{B}^2 \\ &= P(\{\omega : (X(\omega), Y(\omega)) \in B\}) \end{aligned}$$

Recap: Joint distribution function

Recap: Joint distribution function

- ▶ Let X, Y be random variables on the same probability space (Ω, \mathcal{F}, P)

Recap: Joint distribution function

- ▶ Let X, Y be random variables on the same probability space (Ω, \mathcal{F}, P)
- ▶ The joint distribution function of X, Y is $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$, defined by

Recap: Joint distribution function

- ▶ Let X, Y be random variables on the same probability space (Ω, \mathcal{F}, P)
- ▶ The joint distribution function of X, Y is $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$, defined by

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

Recap: Joint distribution function

- ▶ Let X, Y be random variables on the same probability space (Ω, \mathcal{F}, P)
- ▶ The joint distribution function of X, Y is $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$, defined by

$$\begin{aligned} F_{XY}(x, y) &= P[X \leq x, Y \leq y] \\ &= P(\{\omega : X(\omega) \leq x\} \cap \{\omega : Y(\omega) \leq y\}) \end{aligned}$$

Recap: Joint distribution function

- ▶ Let X, Y be random variables on the same probability space (Ω, \mathcal{F}, P)
- ▶ The joint distribution function of X, Y is $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$, defined by

$$\begin{aligned} F_{XY}(x, y) &= P[X \leq x, Y \leq y] \quad (= P_{XY}((-\infty, x] \times (-\infty, y])) \\ &= P(\{\omega : X(\omega) \leq x\} \cap \{\omega : Y(\omega) \leq y\}) \end{aligned}$$

Recap: Joint distribution function

- ▶ Let X, Y be random variables on the same probability space (Ω, \mathcal{F}, P)
- ▶ The joint distribution function of X, Y is $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$, defined by

$$\begin{aligned} F_{XY}(x, y) &= P[X \leq x, Y \leq y] \quad (= P_{XY}((-\infty, x] \times (-\infty, y])) \\ &= P(\{\omega : X(\omega) \leq x\} \cap \{\omega : Y(\omega) \leq y\}) \end{aligned}$$

- ▶ The joint distribution function is the probability of the intersection of the events $[X \leq x]$ and $[Y \leq y]$.

Recap: Joint distribution function

- ▶ Let X, Y be random variables on the same probability space (Ω, \mathcal{F}, P)
- ▶ The joint distribution function of X, Y is $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$, defined by

$$\begin{aligned} F_{XY}(x, y) &= P[X \leq x, Y \leq y] \quad (= P_{XY}((-\infty, x] \times (-\infty, y])) \\ &= P(\{\omega : X(\omega) \leq x\} \cap \{\omega : Y(\omega) \leq y\}) \end{aligned}$$

- ▶ The joint distribution function is the probability of the intersection of the events $[X \leq x]$ and $[Y \leq y]$.
- ▶ Given the joint distribution function, probability of any event involving the pair of random variables can be (in principle) calculated.

Recap: Properties of Joint Distribution Function

- ▶ Joint distribution function: $F_{XY} : \Re^2 \rightarrow \Re$

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

Recap: Properties of Joint Distribution Function

- ▶ Joint distribution function: $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶ It satisfies

1. $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$
 $F_{XY}(\infty, \infty) = 1$

Recap: Properties of Joint Distribution Function

- ▶ Joint distribution function: $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶ It satisfies

1. $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$
 $F_{XY}(\infty, \infty) = 1$
2. F_{XY} is non-decreasing in each of its arguments

Recap: Properties of Joint Distribution Function

- ▶ Joint distribution function: $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶ It satisfies

1. $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$
 $F_{XY}(\infty, \infty) = 1$
2. F_{XY} is non-decreasing in each of its arguments
3. F_{XY} is right continuous and has left-hand limits in each of its arguments

Recap: Properties of Joint Distribution Function

- ▶ Joint distribution function: $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶ It satisfies

1. $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$
 $F_{XY}(\infty, \infty) = 1$
2. F_{XY} is non-decreasing in each of its arguments
3. F_{XY} is right continuous and has left-hand limits in each of its arguments
4. For all $x_1 < x_2$ and $y_1 < y_2$

$$F_{XY}(x_2, y_2) - F_{XY}(x_2, y_1) - F_{XY}(x_1, y_2) + F_{XY}(x_1, y_1) \geq 0$$

Recap: Properties of Joint Distribution Function

- ▶ Joint distribution function: $F_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$

$$F_{XY}(x, y) = P[X \leq x, Y \leq y]$$

- ▶ It satisfies

1. $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y;$
 $F_{XY}(\infty, \infty) = 1$
2. F_{XY} is non-decreasing in each of its arguments
3. F_{XY} is right continuous and has left-hand limits in each of its arguments
4. For all $x_1 < x_2$ and $y_1 < y_2$

$$F_{XY}(x_2, y_2) - F_{XY}(x_2, y_1) - F_{XY}(x_1, y_2) + F_{XY}(x_1, y_1) \geq 0$$

- ▶ Any $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ satisfying the above would be a joint distribution function.

Recap: Joint Probability mass function

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$ be discrete random variables.

Recap: Joint Probability mass function

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$ be discrete random variables.
- ▶ The joint pmf: $f_{XY}(x, y) = P[X = x, Y = y]$.

Recap: Joint Probability mass function

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$ be discrete random variables.
- ▶ The joint pmf: $f_{XY}(x, y) = P[X = x, Y = y]$.
- ▶ The joint pmf satisfies:

Recap: Joint Probability mass function

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$ be discrete random variables.
- ▶ The joint pmf: $f_{XY}(x, y) = P[X = x, Y = y]$.
- ▶ The joint pmf satisfies:
 - ▶ $f_{XY}(x, y) \geq 0, \forall x, y$ and

Recap: Joint Probability mass function

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$ be discrete random variables.
- ▶ The joint pmf: $f_{XY}(x, y) = P[X = x, Y = y]$.
- ▶ The joint pmf satisfies:
 - ▶ $f_{XY}(x, y) \geq 0, \forall x, y$ and
 - ▶ $\sum_i \sum_j f_{XY}(x_i, y_j) = 1$

Recap: Joint Probability mass function

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$ be discrete random variables.
- ▶ The joint pmf: $f_{XY}(x, y) = P[X = x, Y = y]$.
- ▶ The joint pmf satisfies:
 - ▶ $f_{XY}(x, y) \geq 0, \forall x, y$ and
 - ▶ $\sum_i \sum_j f_{XY}(x_i, y_j) = 1$
- ▶ Given the joint pmf, we can get the joint df as

$$F_{XY}(x, y) = \sum_{\substack{i: \\ x_i \leq x}} \sum_{\substack{j: \\ y_j \leq y}} f_{XY}(x_i, y_j)$$

Recap: Joint pmf

- ▶ Given sets $\{x_1, x_2, \dots\}$ and $\{y_1, y_2, \dots\}$.

Recap: Joint pmf

- ▶ Given sets $\{x_1, x_2, \dots\}$ and $\{y_1, y_2, \dots\}$.
- ▶ Suppose $f_{XY} : \mathbb{R}^2 \rightarrow [0, 1]$ be such that

Recap: Joint pmf

- ▶ Given sets $\{x_1, x_2, \dots\}$ and $\{y_1, y_2, \dots\}$.
- ▶ Suppose $f_{XY} : \mathbb{R}^2 \rightarrow [0, 1]$ be such that
 - ▶ $f_{XY}(x, y) = 0$ unless $x = x_i$ for some i and $y = y_j$ for some j , and

Recap: Joint pmf

- ▶ Given sets $\{x_1, x_2, \dots\}$ and $\{y_1, y_2, \dots\}$.
- ▶ Suppose $f_{XY} : \mathbb{R}^2 \rightarrow [0, 1]$ be such that
 - ▶ $f_{XY}(x, y) = 0$ unless $x = x_i$ for some i and $y = y_j$ for some j , and
 - ▶ $\sum_i \sum_j f_{XY}(x_i, y_j) = 1$

Recap: Joint pmf

- ▶ Given sets $\{x_1, x_2, \dots\}$ and $\{y_1, y_2, \dots\}$.
- ▶ Suppose $f_{XY} : \mathbb{R}^2 \rightarrow [0, 1]$ be such that
 - ▶ $f_{XY}(x, y) = 0$ unless $x = x_i$ for some i and $y = y_j$ for some j , and
 - ▶ $\sum_i \sum_j f_{XY}(x_i, y_j) = 1$
- ▶ Then f_{XY} is a joint pmf.

Recap: Joint pmf

- ▶ Given sets $\{x_1, x_2, \dots\}$ and $\{y_1, y_2, \dots\}$.
- ▶ Suppose $f_{XY} : \mathbb{R}^2 \rightarrow [0, 1]$ be such that
 - ▶ $f_{XY}(x, y) = 0$ unless $x = x_i$ for some i and $y = y_j$ for some j , and
 - ▶ $\sum_i \sum_j f_{XY}(x_i, y_j) = 1$
- ▶ Then f_{XY} is a joint pmf.
- ▶ This is because, if we define

$$F_{XY}(x, y) = \sum_{\substack{i: \\ x_i \leq x}} \sum_{\substack{j: \\ y_j \leq y}} f_{XY}(x_i, y_j)$$

then F_{XY} satisfies all properties of a df.

Recap: Joint pmf

- ▶ Given sets $\{x_1, x_2, \dots\}$ and $\{y_1, y_2, \dots\}$.
- ▶ Suppose $f_{XY} : \mathbb{R}^2 \rightarrow [0, 1]$ be such that
 - ▶ $f_{XY}(x, y) = 0$ unless $x = x_i$ for some i and $y = y_j$ for some j , and
 - ▶ $\sum_i \sum_j f_{XY}(x_i, y_j) = 1$
- ▶ Then f_{XY} is a joint pmf.
- ▶ This is because, if we define

$$F_{XY}(x, y) = \sum_{\substack{i: \\ x_i \leq x}} \sum_{\substack{j: \\ y_j \leq y}} f_{XY}(x_i, y_j)$$

then F_{XY} satisfies all properties of a df.

- ▶ We normally specify a pair of discrete random variables by giving the joint pmf

Recap: Joint pmf

- ▶ Given the joint pmf, we can (in principle) compute the probability of any event involving the two discrete random variables.

Recap: Joint pmf

- ▶ Given the joint pmf, we can (in principle) compute the probability of any event involving the two discrete random variables.

$$P[(X, Y) \in B] = \sum_{\substack{i,j: \\ (x_i, y_j) \in B}} f_{XY}(x_i, y_j)$$

Recap: Joint pmf

- ▶ Given the joint pmf, we can (in principle) compute the probability of any event involving the two discrete random variables.

$$P[(X, Y) \in B] = \sum_{\substack{i,j: \\ (x_i, y_j) \in B}} f_{XY}(x_i, y_j)$$

- ▶ Now, events can be specified in terms of relations between the two rv's too

Recap: Joint pmf

- ▶ Given the joint pmf, we can (in principle) compute the probability of any event involving the two discrete random variables.

$$P[(X, Y) \in B] = \sum_{\substack{i,j: \\ (x_i, y_j) \in B}} f_{XY}(x_i, y_j)$$

- ▶ Now, events can be specified in terms of relations between the two rv's too

$$[X < Y + 2] = \{\omega : X(\omega) < Y(\omega) + 2\}$$

Recap: Joint pmf

- ▶ Given the joint pmf, we can (in principle) compute the probability of any event involving the two discrete random variables.

$$P[(X, Y) \in B] = \sum_{\substack{i,j: \\ (x_i, y_j) \in B}} f_{XY}(x_i, y_j)$$

- ▶ Now, events can be specified in terms of relations between the two rv's too

$$[X < Y + 2] = \{\omega : X(\omega) < Y(\omega) + 2\}$$

- ▶ Thus,

$$P[X < Y + 2] = \sum_{\substack{i,j: \\ x_i < y_j + 2}} f_{XY}(x_i, y_j)$$

Joint density function

- ▶ Let X, Y be two continuous rv's with df F_{XY} .

Joint density function

- ▶ Let X, Y be two continuous rv's with df F_{XY} .
- ▶ If there exists a function f_{XY} that satisfies

Joint density function

- ▶ Let X, Y be two continuous rv's with df F_{XY} .
- ▶ If there exists a function f_{XY} that satisfies

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

then we say that X, Y have a joint probability density function which is f_{XY}

Joint density function

- ▶ Let X, Y be two continuous rv's with df F_{XY} .
- ▶ If there exists a function f_{XY} that satisfies

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

then we say that X, Y have a joint probability density function which is f_{XY}

- ▶ Please note the difference in the definition of joint pmf and joint pdf.

Joint density function

- ▶ Let X, Y be two continuous rv's with df F_{XY} .
- ▶ If there exists a function f_{XY} that satisfies

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

then we say that X, Y have a joint probability density function which is f_{XY}

- ▶ Please note the difference in the definition of joint pmf and joint pdf.
- ▶ When X, Y are discrete we defined a joint pmf

Joint density function

- ▶ Let X, Y be two continuous rv's with df F_{XY} .
- ▶ If there exists a function f_{XY} that satisfies

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

then we say that X, Y have a joint probability density function which is f_{XY}

- ▶ Please note the difference in the definition of joint pmf and joint pdf.
- ▶ When X, Y are discrete we defined a joint pmf
- ▶ We are not saying that if X, Y are continuous rv's then a joint density exists.

properties of joint density

- ▶ The joint density (or joint pdf) of X, Y is f_{XY} that satisfies

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

properties of joint density

- ▶ The joint density (or joint pdf) of X, Y is f_{XY} that satisfies

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

- ▶ Since F_{XY} is non-decreasing in each argument, we must have $f_{XY}(x, y) \geq 0$.

properties of joint density

- ▶ The joint density (or joint pdf) of X, Y is f_{XY} that satisfies

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

- ▶ Since F_{XY} is non-decreasing in each argument, we must have $f_{XY}(x, y) \geq 0$.
- ▶ $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' = 1$ is needed to ensure $F_{XY}(\infty, \infty) = 1$.

properties of joint density

- ▶ The joint density f_{XY} satisfies the following

properties of joint density

- ▶ The joint density f_{XY} satisfies the following
 1. $f_{XY}(x, y) \geq 0, \forall x, y$

properties of joint density

► The joint density f_{XY} satisfies the following

1. $f_{XY}(x, y) \geq 0, \quad \forall x, y$

2. $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') \, dy' \, dx' = 1$

properties of joint density

- ▶ The joint density f_{XY} satisfies the following
 1. $f_{XY}(x, y) \geq 0, \quad \forall x, y$
 2. $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' = 1$
- ▶ These are very similar to the properties of the density of a single rv

Example: Joint Density

- ▶ Consider the function

$$f(x, y) = 2, \quad 0 < x < y < 1$$

Example: Joint Density

- ▶ Consider the function

$$f(x, y) = 2, \quad 0 < x < y < 1 \quad (f(x, y) = 0, \text{ otherwise})$$

Example: Joint Density

- ▶ Consider the function

$$f(x, y) = 2, \quad 0 < x < y < 1 \quad (f(x, y) = 0, \text{ otherwise})$$

- ▶ Let us show this is a density

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy =$$

Example: Joint Density

- ▶ Consider the function

$$f(x, y) = 2, \quad 0 < x < y < 1 \quad (f(x, y) = 0, \text{ otherwise})$$

- ▶ Let us show this is a density

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = \int_0^1 \int_0^y 2 \, dx \, dy$$

Example: Joint Density

- ▶ Consider the function

$$f(x, y) = 2, \quad 0 < x < y < 1 \quad (f(x, y) = 0, \text{ otherwise})$$

- ▶ Let us show this is a density

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = \int_0^1 \int_0^y 2 \, dx \, dy = \int_0^1 2x|_0^y \, dy$$

Example: Joint Density

- ▶ Consider the function

$$f(x, y) = 2, \quad 0 < x < y < 1 \quad (f(x, y) = 0, \text{ otherwise})$$

- ▶ Let us show this is a density

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = \int_0^1 \int_0^y 2 \, dx \, dy = \int_0^1 2x|_0^y \, dy = \int_0^1 2y \, dy$$

Example: Joint Density

- ▶ Consider the function

$$f(x, y) = 2, \quad 0 < x < y < 1 \quad (f(x, y) = 0, \text{ otherwise})$$

- ▶ Let us show this is a density

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = \int_0^1 \int_0^y 2 \, dx \, dy = \int_0^1 2x \Big|_0^y \, dy = \int_0^1 2y \, dy = 1$$

Example: Joint Density


- ▶ Consider the function

$$f(x, y) = 2, \quad 0 < x < y < 1 \quad (f(x, y) = 0, \text{ otherwise})$$

- ▶ Let us show this is a density

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = \int_0^1 \int_0^y 2 \, dx \, dy = \int_0^1 2x \Big|_0^y \, dy = \int_0^1 2y \, dy = 1$$

- ▶ We can say this density is uniform over the region



Example: Joint Density


- ▶ Consider the function

$$f(x, y) = 2, \quad 0 < x < y < 1 \quad (f(x, y) = 0, \text{ otherwise})$$

- ▶ Let us show this is a density

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = \int_0^1 \int_0^y 2 \, dx \, dy = \int_0^1 2x \Big|_0^y \, dy = \int_0^1 2y \, dy = 1$$

- ▶ We can say this density is uniform over the region



The figure is not a plot of the density function!!

properties of joint density

- ▶ The joint density f_{XY} satisfies the following

properties of joint density

- ▶ The joint density f_{XY} satisfies the following
 1. $f_{XY}(x, y) \geq 0, \forall x, y$

properties of joint density

► The joint density f_{XY} satisfies the following

1. $f_{XY}(x, y) \geq 0, \quad \forall x, y$

2. $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' = 1$

properties of joint density

- ▶ The joint density f_{XY} satisfies the following
 1. $f_{XY}(x, y) \geq 0, \quad \forall x, y$
 2. $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' = 1$
- ▶ Any function $f_{XY} : \Re^2 \rightarrow \Re$ satisfying the above two is a joint density function.

properties of joint density

- ▶ The joint density f_{XY} satisfies the following
 1. $f_{XY}(x, y) \geq 0, \quad \forall x, y$
 2. $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' = 1$
- ▶ Any function $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ satisfying the above two is a joint density function.
- ▶ Given f_{XY} satisfying the above, define

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

properties of joint density

- ▶ The joint density f_{XY} satisfies the following
 1. $f_{XY}(x, y) \geq 0, \quad \forall x, y$
 2. $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' = 1$
- ▶ Any function $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ satisfying the above two is a joint density function.
- ▶ Given f_{XY} satisfying the above, define

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

- ▶ Then we can show F_{XY} is a joint distribution.

► $f_{XY}(x, y) \geq 0$ and $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' = 1$

- ▶ $f_{XY}(x, y) \geq 0$ and $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' = 1$
- ▶ Define

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

- ▶ $f_{XY}(x, y) \geq 0$ and $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' = 1$
- ▶ Define

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

- ▶ Then, $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y$ and $F_{XY}(\infty, \infty) = 1$

- ▶ $f_{XY}(x, y) \geq 0$ and $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' = 1$
- ▶ Define

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

- ▶ Then, $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y$ and $F_{XY}(\infty, \infty) = 1$
- ▶ Since $f_{XY}(x, y) \geq 0$, F_{XY} is non-decreasing in each argument.

- ▶ $f_{XY}(x, y) \geq 0$ and $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' = 1$
- ▶ Define

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

- ▶ Then, $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y$ and $F_{XY}(\infty, \infty) = 1$
- ▶ Since $f_{XY}(x, y) \geq 0$, F_{XY} is non-decreasing in each argument.
- ▶ Since it is given as an integral, the above also shows that F_{XY} is continuous in each argument.

- ▶ $f_{XY}(x, y) \geq 0$ and $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' = 1$
- ▶ Define

$$F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx', \quad \forall x, y$$

- ▶ Then, $F_{XY}(-\infty, y) = F_{XY}(x, -\infty) = 0, \forall x, y$ and $F_{XY}(\infty, \infty) = 1$
- ▶ Since $f_{XY}(x, y) \geq 0$, F_{XY} is non-decreasing in each argument.
- ▶ Since it is given as an integral, the above also shows that F_{XY} is continuous in each argument.
- ▶ The only property left is the special property of F_{XY} we mentioned earlier.

$$\Delta \triangleq F_{XY}(x_2, y_2) - F_{XY}(x_1, y_2) - F_{XY}(x_2, y_1) + F_{XY}(x_1, y_1).$$

$$\Delta \triangleq F_{XY}(x_2, y_2) - F_{XY}(x_1, y_2) - F_{XY}(x_2, y_1) + F_{XY}(x_1, y_1).$$

► We need to show $\Delta \geq 0$ if $x_1 < x_2$ and $y_1 < y_2$.

$$\Delta \triangleq F_{XY}(x_2, y_2) - F_{XY}(x_1, y_2) - F_{XY}(x_2, y_1) + F_{XY}(x_1, y_1).$$

- ▶ We need to show $\Delta \geq 0$ if $x_1 < x_2$ and $y_1 < y_2$.
- ▶ We have

$$\begin{aligned} \Delta = & \int_{-\infty}^{x_2} \int_{-\infty}^{y_2} f_{XY} \, dy \, dx - \int_{-\infty}^{x_1} \int_{-\infty}^{y_2} f_{XY} \, dy \, dx \\ & - \int_{-\infty}^{x_2} \int_{-\infty}^{y_1} f_{XY} \, dy \, dx + \int_{-\infty}^{x_1} \int_{-\infty}^{y_1} f_{XY} \, dy \, dx \end{aligned}$$

$$\Delta \triangleq F_{XY}(x_2, y_2) - F_{XY}(x_1, y_2) - F_{XY}(x_2, y_1) + F_{XY}(x_1, y_1).$$

- ▶ We need to show $\Delta \geq 0$ if $x_1 < x_2$ and $y_1 < y_2$.
- ▶ We have

$$\begin{aligned} \Delta &= \int_{-\infty}^{x_2} \int_{-\infty}^{y_2} f_{XY} \, dy \, dx - \int_{-\infty}^{x_1} \int_{-\infty}^{y_2} f_{XY} \, dy \, dx \\ &\quad - \int_{-\infty}^{x_2} \int_{-\infty}^{y_1} f_{XY} \, dy \, dx + \int_{-\infty}^{x_1} \int_{-\infty}^{y_1} f_{XY} \, dy \, dx \\ &= \int_{-\infty}^{x_2} \left(\int_{-\infty}^{y_2} f_{XY} \, dy - \int_{-\infty}^{y_1} f_{XY} \, dy \right) dx \end{aligned}$$

$$\Delta \triangleq F_{XY}(x_2, y_2) - F_{XY}(x_1, y_2) - F_{XY}(x_2, y_1) + F_{XY}(x_1, y_1).$$

- ▶ We need to show $\Delta \geq 0$ if $x_1 < x_2$ and $y_1 < y_2$.
- ▶ We have

$$\begin{aligned} \Delta &= \int_{-\infty}^{x_2} \int_{-\infty}^{y_2} f_{XY} \, dy \, dx - \int_{-\infty}^{x_1} \int_{-\infty}^{y_2} f_{XY} \, dy \, dx \\ &\quad - \int_{-\infty}^{x_2} \int_{-\infty}^{y_1} f_{XY} \, dy \, dx + \int_{-\infty}^{x_1} \int_{-\infty}^{y_1} f_{XY} \, dy \, dx \\ &= \int_{-\infty}^{x_2} \left(\int_{-\infty}^{y_2} f_{XY} \, dy - \int_{-\infty}^{y_1} f_{XY} \, dy \right) dx \\ &\quad - \int_{-\infty}^{x_1} \left(\int_{-\infty}^{y_2} f_{XY} \, dy - \int_{-\infty}^{y_1} f_{XY} \, dy \right) dx \end{aligned}$$

► Thus we have

$$\begin{aligned}\Delta &= \int_{-\infty}^{x_2} \left(\int_{-\infty}^{y_2} f_{XY} dy - \int_{-\infty}^{y_1} f_{XY} dy \right) dx \\ &\quad - \int_{-\infty}^{x_1} \left(\int_{-\infty}^{y_2} f_{XY} dy - \int_{-\infty}^{y_1} f_{XY} dy \right) dx\end{aligned}$$

► Thus we have

$$\begin{aligned}\Delta &= \int_{-\infty}^{x_2} \left(\int_{-\infty}^{y_2} f_{XY} dy - \int_{-\infty}^{y_1} f_{XY} dy \right) dx \\ &\quad - \int_{-\infty}^{x_1} \left(\int_{-\infty}^{y_2} f_{XY} dy - \int_{-\infty}^{y_1} f_{XY} dy \right) dx \\ &= \int_{-\infty}^{x_2} \int_{y_1}^{y_2} f_{XY} dy dx - \int_{-\infty}^{x_1} \int_{y_1}^{y_2} f_{XY} dy dx\end{aligned}$$

► Thus we have

$$\begin{aligned}\Delta &= \int_{-\infty}^{x_2} \left(\int_{-\infty}^{y_2} f_{XY} dy - \int_{-\infty}^{y_1} f_{XY} dy \right) dx \\ &\quad - \int_{-\infty}^{x_1} \left(\int_{-\infty}^{y_2} f_{XY} dy - \int_{-\infty}^{y_1} f_{XY} dy \right) dx \\ &= \int_{-\infty}^{x_2} \int_{y_1}^{y_2} f_{XY} dy dx - \int_{-\infty}^{x_1} \int_{y_1}^{y_2} f_{XY} dy dx \\ &= \int_{x_1}^{x_2} \int_{y_1}^{y_2} f_{XY} dy dx \geq 0\end{aligned}$$

► Thus we have

$$\begin{aligned}\Delta &= \int_{-\infty}^{x_2} \left(\int_{-\infty}^{y_2} f_{XY} dy - \int_{-\infty}^{y_1} f_{XY} dy \right) dx \\ &\quad - \int_{-\infty}^{x_1} \left(\int_{-\infty}^{y_2} f_{XY} dy - \int_{-\infty}^{y_1} f_{XY} dy \right) dx \\ &= \int_{-\infty}^{x_2} \int_{y_1}^{y_2} f_{XY} dy dx - \int_{-\infty}^{x_1} \int_{y_1}^{y_2} f_{XY} dy dx \\ &= \int_{x_1}^{x_2} \int_{y_1}^{y_2} f_{XY} dy dx \geq 0\end{aligned}$$

► This actually shows

$$P[x_1 \leq X \leq x_2, y_1 \leq Y \leq y_2] = \int_{x_1}^{x_2} \int_{y_1}^{y_2} f_{XY} dy dx$$

- ▶ What we showed is the following

- ▶ What we showed is the following
- ▶ Any function $f_{XY} : \Re^2 \rightarrow \Re$ that satisfies

- ▶ What we showed is the following
- ▶ Any function $f_{XY} : \Re^2 \rightarrow \Re$ that satisfies
 - ▶ $f_{XY}(x, y) \geq 0, \forall x, y$

- ▶ What we showed is the following
- ▶ Any function $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ that satisfies
 - ▶ $f_{XY}(x, y) \geq 0, \forall x, y$
 - ▶ $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x, y) dx dy = 1$is a joint density function.

- ▶ What we showed is the following
- ▶ Any function $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ that satisfies
 - ▶ $f_{XY}(x, y) \geq 0, \forall x, y$
 - ▶ $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x, y) dx dy = 1$is a joint density function.
- ▶ This is because now
$$F_{XY}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{XY}(x, y) dx dy$$
would satisfy all conditions for a df.

- ▶ What we showed is the following
- ▶ Any function $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ that satisfies
 - ▶ $f_{XY}(x, y) \geq 0, \forall x, y$
 - ▶ $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x, y) dx dy = 1$is a joint density function.
- ▶ This is because now
$$F_{XY}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{XY}(x, y) dx dy$$
would satisfy all conditions for a df.
- ▶ Convenient to specify joint density (when it exists)

- ▶ What we showed is the following
- ▶ Any function $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ that satisfies

- ▶ $f_{XY}(x, y) \geq 0, \forall x, y$

- ▶ $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x, y) dx dy = 1$

is a joint density function.

- ▶ This is because now

$$F_{XY}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{XY}(x, y) dx dy$$

would satisfy all conditions for a df.

- ▶ Convenient to specify joint density (when it exists)
- ▶ We also showed

$$P[x_1 \leq X \leq x_2, y_1 \leq Y \leq y_2] = \int_{x_1}^{x_2} \int_{y_1}^{y_2} f_{XY} dy dx$$

- ▶ What we showed is the following
- ▶ Any function $f_{XY} : \mathbb{R}^2 \rightarrow \mathbb{R}$ that satisfies
 - ▶ $f_{XY}(x, y) \geq 0, \forall x, y$
 - ▶ $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x, y) dx dy = 1$
 is a joint density function.

- ▶ This is because now
 $F_{XY}(x, y) = \int_{-\infty}^y \int_{-\infty}^x f_{XY}(x, y) dx dy$
 would satisfy all conditions for a df.

- ▶ Convenient to specify joint density (when it exists)
- ▶ We also showed

$$P[x_1 \leq X \leq x_2, y_1 \leq Y \leq y_2] = \int_{x_1}^{x_2} \int_{y_1}^{y_2} f_{XY} dy dx$$

- ▶ In general

$$P[(X, Y) \in B] = \int_B f_{XY}(x, y) dx dy, \quad \forall B \in \mathcal{B}^2$$

- ▶ Let us consider the example

$$f(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ Let us consider the example

$$f(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ Suppose we want probability of $[Y > X + 0.5]$

- ▶ Let us consider the example

$$f(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ Suppose we want probability of $[Y > X + 0.5]$

$$P[Y > X + 0.5]$$

- ▶ Let us consider the example

$$f(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ Suppose we want probability of $[Y > X + 0.5]$

$$P[Y > X + 0.5] = P[(X, Y) \in \{(x, y) : y > x + 0.5\}]$$

- ▶ Let us consider the example

$$f(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ Suppose we want probability of $[Y > X + 0.5]$

$$\begin{aligned} P[Y > X + 0.5] &= P[(X, Y) \in \{(x, y) : y > x + 0.5\}] \\ &= \int_{\{(x, y) : y > x + 0.5\}} f_{XY}(x, y) \, dx \, dy \end{aligned}$$

- Let us consider the example

$$f(x, y) = 2, \quad 0 < x < y < 1$$

- Suppose we want probability of $[Y > X + 0.5]$

$$\begin{aligned} P[Y > X + 0.5] &= P[(X, Y) \in \{(x, y) : y > x + 0.5\}] \\ &= \int_{\{(x, y) : y > x + 0.5\}} f_{XY}(x, y) \, dx \, dy \\ &= \int_{0.5}^1 \int_0^{y-0.5} 2 \, dx \, dy \end{aligned}$$

- ▶ Let us consider the example

$$f(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ Suppose we want probability of $[Y > X + 0.5]$

$$\begin{aligned} P[Y > X + 0.5] &= P[(X, Y) \in \{(x, y) : y > x + 0.5\}] \\ &= \int_{\{(x, y) : y > x + 0.5\}} f_{XY}(x, y) \, dx \, dy \\ &= \int_{0.5}^1 \int_0^{y-0.5} 2 \, dx \, dy \\ &= \int_{0.5}^1 2(y - 0.5) \, dy \end{aligned}$$

- ▶ Let us consider the example

$$f(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ Suppose we want probability of $[Y > X + 0.5]$

$$\begin{aligned} P[Y > X + 0.5] &= P[(X, Y) \in \{(x, y) : y > x + 0.5\}] \\ &= \int_{\{(x, y) : y > x + 0.5\}} f_{XY}(x, y) \, dx \, dy \\ &= \int_{0.5}^1 \int_0^{y-0.5} 2 \, dx \, dy \\ &= \int_{0.5}^1 2(y - 0.5) \, dy \\ &= 2 \left. \frac{y^2}{2} \right|_{0.5}^1 - y \Big|_{0.5}^1 \end{aligned}$$


- ▶ Let us consider the example

$$f(x, y) = 2, \quad 0 < x < y < 1$$


- ▶ Suppose we want probability of $[Y > X + 0.5]$

$$\begin{aligned} P[Y > X + 0.5] &= P[(X, Y) \in \{(x, y) : y > x + 0.5\}] \\ &= \int_{\{(x, y) : y > x + 0.5\}} f_{XY}(x, y) \, dx \, dy \\ &= \int_{0.5}^1 \int_0^{y-0.5} 2 \, dx \, dy \\ &= \int_{0.5}^1 2(y - 0.5) \, dy \\ &= 2 \left. \frac{y^2}{2} \right|_{0.5}^1 - y \Big|_{0.5}^1 = 1 - 0.25 - 1 + 0.5 = 0.25 \end{aligned}$$

- We can look at it geometrically



- We can look at it geometrically



- The probability of the event we want is the area of the small triangle divided by that of the big triangle.

Marginal Distributions

- ▶ Let X, Y be random variables with joint distribution function F_{XY} .

Marginal Distributions

- ▶ Let X, Y be random variables with joint distribution function F_{XY} .
- ▶ We know $F_{XY}(x, y) = P[X \leq x, Y \leq y]$.

Marginal Distributions

- ▶ Let X, Y be random variables with joint distribution function F_{XY} .
- ▶ We know $F_{XY}(x, y) = P[X \leq x, Y \leq y]$.
- ▶ Hence

$$F_{XY}(x, \infty) = P[X \leq x, Y \leq \infty]$$

Marginal Distributions

- ▶ Let X, Y be random variables with joint distribution function F_{XY} .
- ▶ We know $F_{XY}(x, y) = P[X \leq x, Y \leq y]$.
- ▶ Hence

$$F_{XY}(x, \infty) = P[X \leq x, Y \leq \infty] = P[X \leq x]$$

Marginal Distributions

- ▶ Let X, Y be random variables with joint distribution function F_{XY} .
- ▶ We know $F_{XY}(x, y) = P[X \leq x, Y \leq y]$.
- ▶ Hence

$$F_{XY}(x, \infty) = P[X \leq x, Y \leq \infty] = P[X \leq x] = F_X(x)$$

Marginal Distributions

- ▶ Let X, Y be random variables with joint distribution function F_{XY} .
- ▶ We know $F_{XY}(x, y) = P[X \leq x, Y \leq y]$.
- ▶ Hence

$$F_{XY}(x, \infty) = P[X \leq x, Y \leq \infty] = P[X \leq x] = F_X(x)$$

- ▶ We define the marginal distribution functions of X, Y by

$$F_X(x) = F_{XY}(x, \infty); \quad F_Y(y) = F_{XY}(\infty, y)$$

Marginal Distributions

- ▶ Let X, Y be random variables with joint distribution function F_{XY} .
- ▶ We know $F_{XY}(x, y) = P[X \leq x, Y \leq y]$.
- ▶ Hence

$$F_{XY}(x, \infty) = P[X \leq x, Y \leq \infty] = P[X \leq x] = F_X(x)$$

- ▶ We define the marginal distribution functions of X, Y by

$$F_X(x) = F_{XY}(x, \infty); \quad F_Y(y) = F_{XY}(\infty, y)$$

- ▶ These are simply distribution functions of X and Y obtained from the joint distribution function.

Marginal mass functions

Marginal mass functions

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$

Marginal mass functions

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$
- ▶ Let f_{XY} be their joint mass function.

Marginal mass functions

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$
- ▶ Let f_{XY} be their joint mass function.
- ▶ Then

$$P[X = x_i] = \sum_j P[X = x_i, Y = y_j]$$

Marginal mass functions

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$
- ▶ Let f_{XY} be their joint mass function.
- ▶ Then

$$P[X = x_i] = \sum_j P[X = x_i, Y = y_j]$$

(This is because $[Y = y_j]$, $j = 1, \dots$, form a partition and $P(A) = \sum_i P(AB_i)$ when B_i is a partition)

Marginal mass functions

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$
- ▶ Let f_{XY} be their joint mass function.
- ▶ Then

$$P[X = x_i] = \sum_j P[X = x_i, Y = y_j] = \sum_j f_{XY}(x_i, y_j)$$

(This is because $[Y = y_j]$, $j = 1, \dots$, form a partition and $P(A) = \sum_i P(AB_i)$ when B_i is a partition)

Marginal mass functions

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$
- ▶ Let f_{XY} be their joint mass function.
- ▶ Then

$$P[X = x_i] = \sum_j P[X = x_i, Y = y_j] = \sum_j f_{XY}(x_i, y_j)$$

(This is because $[Y = y_j]$, $j = 1, \dots$, form a partition and $P(A) = \sum_i P(AB_i)$ when B_i is a partition)

- ▶ We define the marginal mass functions of X and Y as

$$f_X(x_i) = \sum_j f_{XY}(x_i, y_j); \quad f_Y(y_j) = \sum_i f_{XY}(x_i, y_j)$$

Marginal mass functions

- ▶ Let $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$
- ▶ Let f_{XY} be their joint mass function.
- ▶ Then

$$P[X = x_i] = \sum_j P[X = x_i, Y = y_j] = \sum_j f_{XY}(x_i, y_j)$$

(This is because $[Y = y_j]$, $j = 1, \dots$, form a partition and $P(A) = \sum_i P(AB_i)$ when B_i is a partition)

- ▶ We define the marginal mass functions of X and Y as

$$f_X(x_i) = \sum_j f_{XY}(x_i, y_j); \quad f_Y(y_j) = \sum_i f_{XY}(x_i, y_j)$$

- ▶ These are mass functions of X and Y obtained from the joint mass function

marginal density functions

- ▶ Let X, Y be continuous rv with joint density f_{XY} .

marginal density functions

- ▶ Let X, Y be continuous rv with joint density f_{XY} .
- ▶ Then we know $F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx'$

marginal density functions

- ▶ Let X, Y be continuous rv with joint density f_{XY} .
- ▶ Then we know $F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx'$
- ▶ Hence, we have

$$F_X(x) = F_{XY}(x, \infty)$$

marginal density functions

- ▶ Let X, Y be continuous rv with joint density f_{XY} .
- ▶ Then we know $F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx'$
- ▶ Hence, we have

$$F_X(x) = F_{XY}(x, \infty) = \int_{-\infty}^x \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx'$$

marginal density functions

- ▶ Let X, Y be continuous rv with joint density f_{XY} .
- ▶ Then we know $F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx'$
- ▶ Hence, we have

$$\begin{aligned} F_X(x) = F_{XY}(x, \infty) &= \int_{-\infty}^x \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' \\ &= \int_{-\infty}^x \left(\int_{-\infty}^{\infty} f_{XY}(x', y') dy' \right) dx' \end{aligned}$$

marginal density functions

- ▶ Let X, Y be continuous rv with joint density f_{XY} .
- ▶ Then we know $F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx'$
- ▶ Hence, we have

$$\begin{aligned} F_X(x) = F_{XY}(x, \infty) &= \int_{-\infty}^x \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' \\ &= \int_{-\infty}^x \left(\int_{-\infty}^{\infty} f_{XY}(x', y') dy' \right) dx' \end{aligned}$$

- ▶ Since X is a continuous rv, this means

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy$$

marginal density functions

- ▶ Let X, Y be continuous rv with joint density f_{XY} .
- ▶ Then we know $F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx'$
- ▶ Hence, we have

$$\begin{aligned} F_X(x) = F_{XY}(x, \infty) &= \int_{-\infty}^x \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' \\ &= \int_{-\infty}^x \left(\int_{-\infty}^{\infty} f_{XY}(x', y') dy' \right) dx' \end{aligned}$$

- ▶ Since X is a continuous rv, this means

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy$$

We call this the marginal density of X .

marginal density functions

- ▶ Let X, Y be continuous rv with joint density f_{XY} .
- ▶ Then we know $F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx'$
- ▶ Hence, we have

$$\begin{aligned} F_X(x) = F_{XY}(x, \infty) &= \int_{-\infty}^x \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' \\ &= \int_{-\infty}^x \left(\int_{-\infty}^{\infty} f_{XY}(x', y') dy' \right) dx' \end{aligned}$$

- ▶ Since X is a continuous rv, this means

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy$$

We call this the marginal density of X .

- ▶ Similarly, marginal density of Y is

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx$$

marginal density functions

- ▶ Let X, Y be continuous rv with joint density f_{XY} .
- ▶ Then we know $F_{XY}(x, y) = \int_{-\infty}^x \int_{-\infty}^y f_{XY}(x', y') dy' dx'$
- ▶ Hence, we have

$$\begin{aligned} F_X(x) = F_{XY}(x, \infty) &= \int_{-\infty}^x \int_{-\infty}^{\infty} f_{XY}(x', y') dy' dx' \\ &= \int_{-\infty}^x \left(\int_{-\infty}^{\infty} f_{XY}(x', y') dy' \right) dx' \end{aligned}$$

- ▶ Since X is a continuous rv, this means

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy$$

We call this the marginal density of X .

- ▶ Similarly, marginal density of Y is

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx$$

- ▶ These are pdf's of X and Y obtained from the joint

Example

- ▶ Rolling two dice, X is max, Y is sum

Example

- ▶ Rolling two dice, X is max, Y is sum
- ▶ We had, for $1 \leq m \leq 6$ and $2 \leq n \leq 12$,

Example

- ▶ Rolling two dice, X is max, Y is sum
- ▶ We had, for $1 \leq m \leq 6$ and $2 \leq n \leq 12$,

$$f_{XY}(m, n) = \begin{cases} \frac{2}{36} & \text{if } m < n < 2m \\ \frac{1}{36} & \text{if } n = 2m \end{cases}$$

Example

- ▶ Rolling two dice, X is max, Y is sum
- ▶ We had, for $1 \leq m \leq 6$ and $2 \leq n \leq 12$,

$$f_{XY}(m, n) = \begin{cases} \frac{2}{36} & \text{if } m < n < 2m \\ \frac{1}{36} & \text{if } n = 2m \end{cases}$$

- ▶ We know, $f_X(m) = \sum_n f_{XY}(m, n)$, $m = 1, \dots, 6$.

Example

- ▶ Rolling two dice, X is max, Y is sum
- ▶ We had, for $1 \leq m \leq 6$ and $2 \leq n \leq 12$,

$$f_{XY}(m, n) = \begin{cases} \frac{2}{36} & \text{if } m < n < 2m \\ \frac{1}{36} & \text{if } n = 2m \end{cases}$$

- ▶ We know, $f_X(m) = \sum_n f_{XY}(m, n)$, $m = 1, \dots, 6$.
- ▶ Given m , for what values of n , $f_{XY}(m, n) > 0$?

Example

- ▶ Rolling two dice, X is max, Y is sum
- ▶ We had, for $1 \leq m \leq 6$ and $2 \leq n \leq 12$,

$$f_{XY}(m, n) = \begin{cases} \frac{2}{36} & \text{if } m < n < 2m \\ \frac{1}{36} & \text{if } n = 2m \end{cases}$$

- ▶ We know, $f_X(m) = \sum_n f_{XY}(m, n)$, $m = 1, \dots, 6$.
- ▶ Given m , for what values of n , $f_{XY}(m, n) > 0$?
We can only have $n = m + 1, \dots, 2m$.

Example

- ▶ Rolling two dice, X is max, Y is sum
- ▶ We had, for $1 \leq m \leq 6$ and $2 \leq n \leq 12$,

$$f_{XY}(m, n) = \begin{cases} \frac{2}{36} & \text{if } m < n < 2m \\ \frac{1}{36} & \text{if } n = 2m \end{cases}$$

- ▶ We know, $f_X(m) = \sum_n f_{XY}(m, n)$, $m = 1, \dots, 6$.
- ▶ Given m , for what values of n , $f_{XY}(m, n) > 0$?
We can only have $n = m + 1, \dots, 2m$.
- ▶ Hence we get

$$f_X(m) = \sum_{n=m+1}^{2m} f_{XY}(m, n)$$

Example

- ▶ Rolling two dice, X is max, Y is sum
- ▶ We had, for $1 \leq m \leq 6$ and $2 \leq n \leq 12$,

$$f_{XY}(m, n) = \begin{cases} \frac{2}{36} & \text{if } m < n < 2m \\ \frac{1}{36} & \text{if } n = 2m \end{cases}$$

- ▶ We know, $f_X(m) = \sum_n f_{XY}(m, n)$, $m = 1, \dots, 6$.
- ▶ Given m , for what values of n , $f_{XY}(m, n) > 0$?
We can only have $n = m + 1, \dots, 2m$.
- ▶ Hence we get

$$f_X(m) = \sum_{n=m+1}^{2m} f_{XY}(m, n) = \sum_{n=m+1}^{2m-1} \frac{2}{36} + \frac{1}{36}$$

Example

- ▶ Rolling two dice, X is max, Y is sum
- ▶ We had, for $1 \leq m \leq 6$ and $2 \leq n \leq 12$,

$$f_{XY}(m, n) = \begin{cases} \frac{2}{36} & \text{if } m < n < 2m \\ \frac{1}{36} & \text{if } n = 2m \end{cases}$$

- ▶ We know, $f_X(m) = \sum_n f_{XY}(m, n)$, $m = 1, \dots, 6$.
- ▶ Given m , for what values of n , $f_{XY}(m, n) > 0$?
We can only have $n = m + 1, \dots, 2m$.
- ▶ Hence we get

$$f_X(m) = \sum_{n=m+1}^{2m} f_{XY}(m, n) = \sum_{n=m+1}^{2m-1} \frac{2}{36} + \frac{1}{36} = \frac{2}{36}(m-1) + \frac{1}{36}$$

Example

- ▶ Rolling two dice, X is max, Y is sum
- ▶ We had, for $1 \leq m \leq 6$ and $2 \leq n \leq 12$,

$$f_{XY}(m, n) = \begin{cases} \frac{2}{36} & \text{if } m < n < 2m \\ \frac{1}{36} & \text{if } n = 2m \end{cases}$$

- ▶ We know, $f_X(m) = \sum_n f_{XY}(m, n)$, $m = 1, \dots, 6$.
- ▶ Given m , for what values of n , $f_{XY}(m, n) > 0$?
We can only have $n = m + 1, \dots, 2m$.
- ▶ Hence we get

$$f_X(m) = \sum_{n=m+1}^{2m} f_{XY}(m, n) = \sum_{n=m+1}^{2m-1} \frac{2}{36} + \frac{1}{36} = \frac{2}{36}(m-1) + \frac{1}{36} = \frac{2m-1}{36}$$

Example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

Example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ The marginal density of X is:

Example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ The marginal density of X is:

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) \, dy$$

Example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ The marginal density of X is: for $0 < x < 1$,

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) \, dy = \int_x^1 2 \, dy$$

Example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ The marginal density of X is: for $0 < x < 1$,

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) \, dy = \int_x^1 2 \, dy = 2(1 - x)$$

Example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ The marginal density of X is: for $0 < x < 1$,

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) \, dy = \int_x^1 2 \, dy = 2(1 - x)$$

Thus, $f_X(x) = 2(1 - x)$, $0 < x < 1$

Example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ The marginal density of X is: for $0 < x < 1$,

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) \, dy = \int_x^1 2 \, dy = 2(1 - x)$$

Thus, $f_X(x) = 2(1 - x)$, $0 < x < 1$

- ▶ We can easily verify this is a density

$$\int_{-\infty}^{\infty} f_X(x) \, dx = \int_0^1 2(1 - x) \, dx$$

Example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ The marginal density of X is: for $0 < x < 1$,

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) \, dy = \int_x^1 2 \, dy = 2(1 - x)$$

Thus, $f_X(x) = 2(1 - x)$, $0 < x < 1$

- ▶ We can easily verify this is a density

$$\int_{-\infty}^{\infty} f_X(x) \, dx = \int_0^1 2(1 - x) \, dx = (2x - x^2) \Big|_0^1 = 1$$

We have: $f_{XY}(x, y) = 2, \quad 0 < x < y < 1$

We have: $f_{XY}(x, y) = 2, \quad 0 < x < y < 1$

► We can similarly find density of Y .

We have: $f_{XY}(x, y) = 2, \quad 0 < x < y < 1$

- ▶ We can similarly find density of Y .
- ▶ For $0 < y < 1$,

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx$$

We have: $f_{XY}(x, y) = 2, \quad 0 < x < y < 1$

- ▶ We can similarly find density of Y .
- ▶ For $0 < y < 1$,

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx = \int_0^y 2 dx$$

We have: $f_{XY}(x, y) = 2, \quad 0 < x < y < 1$

- ▶ We can similarly find density of Y .
- ▶ For $0 < y < 1$,

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx = \int_0^y 2 dx = 2y$$

We have: $f_{XY}(x, y) = 2, \quad 0 < x < y < 1$

- ▶ We can similarly find density of Y .
- ▶ For $0 < y < 1$,

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx = \int_0^y 2 dx = 2y$$

- ▶ Thus, $f_Y(y) = 2y, \quad 0 < y < 1$ and

We have: $f_{XY}(x, y) = 2, \quad 0 < x < y < 1$

- ▶ We can similarly find density of Y .
- ▶ For $0 < y < 1$,

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx = \int_0^y 2 dx = 2y$$

- ▶ Thus, $f_Y(y) = 2y, \quad 0 < y < 1$ and

$$\int_0^1 2y dy = 2 \left. \frac{y^2}{2} \right|_0^1 = 1$$

- ▶ If we are given the joint df or joint pmf/joint density of X, Y , then the individual df or pmf/pdf are uniquely determined.

- ▶ If we are given the joint df or joint pmf/joint density of X, Y , then the individual df or pmf/pdf are uniquely determined.
- ▶ However, given individual pdf of X and Y , we cannot determine the joint density. (same is true of pmf or df)

- ▶ If we are given the joint df or joint pmf/joint density of X, Y , then the individual df or pmf/pdf are uniquely determined.
- ▶ However, given individual pdf of X and Y , we cannot determine the joint density. (same is true of pmf or df)
- ▶ There can be many different joint density functions all having the same marginals

Conditional distributions

- ▶ Let X, Y be rv's on the same probability space

Conditional distributions

- ▶ Let X, Y be rv's on the same probability space
- ▶ We define the conditional distribution of X given Y by

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

(For now ignore the case of $P[Y = y] = 0$).

Conditional distributions

- ▶ Let X, Y be rv's on the same probability space
- ▶ We define the conditional distribution of X given Y by

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

(For now ignore the case of $P[Y = y] = 0$).

- ▶ Note that $F_{X|Y} : \mathbb{R}^2 \rightarrow \mathbb{R}$

Conditional distributions

- ▶ Let X, Y be rv's on the same probability space
- ▶ We define the conditional distribution of X given Y by

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

(For now ignore the case of $P[Y = y] = 0$).

- ▶ Note that $F_{X|Y} : \mathbb{R}^2 \rightarrow \mathbb{R}$
- ▶ $F_{X|Y}(x|y)$ is a notation. We could write $F_{X|Y}(x, y)$.

- Conditional distribution of X given Y is

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- Conditional distribution of X given Y is

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

It is the conditional probability of $[X \leq x]$ given (or conditioned on) $[Y = y]$.

- ▶ Conditional distribution of X given Y is

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

It is the conditional probability of $[X \leq x]$ given (or conditioned on) $[Y = y]$.

- ▶ Consider example: rolling 2 dice, X is max, Y is sum

- Conditional distribution of X given Y is

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

It is the conditional probability of $[X \leq x]$ given (or conditioned on) $[Y = y]$.

- Consider example: rolling 2 dice, X is max, Y is sum

$$P[X \leq 4 | Y = 3] = 1;$$

- Conditional distribution of X given Y is

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

It is the conditional probability of $[X \leq x]$ given (or conditioned on) $[Y = y]$.

- Consider example: rolling 2 dice, X is max, Y is sum

$$P[X \leq 4 | Y = 3] = 1; \quad P[X \leq 4 | Y = 9] = 0$$

- ▶ Conditional distribution of X given Y is

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

It is the conditional probability of $[X \leq x]$ given (or conditioned on) $[Y = y]$.

- ▶ Consider example: rolling 2 dice, X is max, Y is sum

$$P[X \leq 4 | Y = 3] = 1; \quad P[X \leq 4 | Y = 9] = 0$$

- ▶ This is what conditional distribution captures.

- ▶ Conditional distribution of X given Y is

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

It is the conditional probability of $[X \leq x]$ given (or conditioned on) $[Y = y]$.

- ▶ Consider example: rolling 2 dice, X is max, Y is sum

$$P[X \leq 4 | Y = 3] = 1; \quad P[X \leq 4 | Y = 9] = 0$$

- ▶ This is what conditional distribution captures.
- ▶ For every value of y , $F_{X|Y}(x|y)$ is a distribution function in the variable x .

- ▶ Conditional distribution of X given Y is

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

It is the conditional probability of $[X \leq x]$ given (or conditioned on) $[Y = y]$.

- ▶ Consider example: rolling 2 dice, X is max, Y is sum

$$P[X \leq 4 | Y = 3] = 1; \quad P[X \leq 4 | Y = 9] = 0$$

- ▶ This is what conditional distribution captures.
- ▶ For every value of y , $F_{X|Y}(x|y)$ is a distribution function in the variable x .
- ▶ It defines a new distribution for X based on knowing the value of Y .

► Let: $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$. Then

► Let: $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$. Then

$$F_{X|Y}(x|y_j) = P[X \leq x | Y = y_j]$$

► Let: $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$. Then

$$F_{X|Y}(x|y_j) = P[X \leq x|Y = y_j] = \frac{P[X \leq x, Y = y_j]}{P[Y = y_j]}$$

► Let: $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$. Then

$$F_{X|Y}(x|y_j) = P[X \leq x | Y = y_j] = \frac{P[X \leq x, Y = y_j]}{P[Y = y_j]}$$

(We define $F_{X|Y}(x|y)$ only when $y = y_j$ for some j).

- Let: $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$. Then

$$F_{X|Y}(x|y_j) = P[X \leq x | Y = y_j] = \frac{P[X \leq x, Y = y_j]}{P[Y = y_j]}$$

(We define $F_{X|Y}(x|y)$ only when $y = y_j$ for some j).

- For each y_j , $F_{X|Y}(x|y_j)$ is a df of a discrete rv in x .

- Let: $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$. Then

$$F_{X|Y}(x|y_j) = P[X \leq x | Y = y_j] = \frac{P[X \leq x, Y = y_j]}{P[Y = y_j]}$$

(We define $F_{X|Y}(x|y)$ only when $y = y_j$ for some j).

- For each y_j , $F_{X|Y}(x|y_j)$ is a df of a discrete rv in x .
- Since X is a discrete rv, we can write the above as

- Let: $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$. Then

$$F_{X|Y}(x|y_j) = P[X \leq x | Y = y_j] = \frac{P[X \leq x, Y = y_j]}{P[Y = y_j]}$$

(We define $F_{X|Y}(x|y)$ only when $y = y_j$ for some j).

- For each y_j , $F_{X|Y}(x|y_j)$ is a df of a discrete rv in x .
- Since X is a discrete rv, we can write the above as

$$F_{X|Y}(x|y_j) = \frac{P[X \leq x, Y = y_j]}{P[Y = y_j]}$$

- Let: $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$. Then

$$F_{X|Y}(x|y_j) = P[X \leq x | Y = y_j] = \frac{P[X \leq x, Y = y_j]}{P[Y = y_j]}$$

(We define $F_{X|Y}(x|y)$ only when $y = y_j$ for some j).

- For each y_j , $F_{X|Y}(x|y_j)$ is a df of a discrete rv in x .
- Since X is a discrete rv, we can write the above as

$$F_{X|Y}(x|y_j) = \frac{P[X \leq x, Y = y_j]}{P[Y = y_j]} = \frac{\sum_{i: x_i \leq x} P[X = x_i, Y = y_j]}{P[Y = y_j]}$$

- Let: $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$. Then

$$F_{X|Y}(x|y_j) = P[X \leq x|Y = y_j] = \frac{P[X \leq x, Y = y_j]}{P[Y = y_j]}$$

(We define $F_{X|Y}(x|y)$ only when $y = y_j$ for some j).

- For each y_j , $F_{X|Y}(x|y_j)$ is a df of a discrete rv in x .
- Since X is a discrete rv, we can write the above as

$$\begin{aligned} F_{X|Y}(x|y_j) &= \frac{P[X \leq x, Y = y_j]}{P[Y = y_j]} = \frac{\sum_{i:x_i \leq x} P[X = x_i, Y = y_j]}{P[Y = y_j]} \\ &= \sum_{i:x_i \leq x} \frac{P[X = x_i, Y = y_j]}{P[Y = y_j]} \end{aligned}$$

- Let: $X \in \{x_1, x_2, \dots\}$ and $Y \in \{y_1, y_2, \dots\}$. Then

$$F_{X|Y}(x|y_j) = P[X \leq x|Y = y_j] = \frac{P[X \leq x, Y = y_j]}{P[Y = y_j]}$$

(We define $F_{X|Y}(x|y)$ only when $y = y_j$ for some j).

- For each y_j , $F_{X|Y}(x|y_j)$ is a df of a discrete rv in x .
► Since X is a discrete rv, we can write the above as

$$\begin{aligned} F_{X|Y}(x|y_j) &= \frac{P[X \leq x, Y = y_j]}{P[Y = y_j]} = \frac{\sum_{i:x_i \leq x} P[X = x_i, Y = y_j]}{P[Y = y_j]} \\ &= \sum_{i:x_i \leq x} \frac{P[X = x_i, Y = y_j]}{P[Y = y_j]} \\ &= \sum_{i:x_i \leq x} \left(\frac{f_{XY}(x_i, y_j)}{f_Y(y_j)} \right) \end{aligned}$$

Conditional mass function

- We got

$$F_{X|Y}(x|y_j) = \sum_{i:x_i \leq x} \left(\frac{f_{XY}(x_i, y_j)}{f_Y(y_j)} \right)$$

Conditional mass function

- We got

$$F_{X|Y}(x|y_j) = \sum_{i: x_i \leq x} \left(\frac{f_{XY}(x_i, y_j)}{f_Y(y_j)} \right)$$

- We define the conditional mass function of X given Y as

$$f_{X|Y}(x_i|y_j) = \frac{f_{XY}(x_i, y_j)}{f_Y(y_j)}$$

Conditional mass function

- We got

$$F_{X|Y}(x|y_j) = \sum_{i: x_i \leq x} \left(\frac{f_{XY}(x_i, y_j)}{f_Y(y_j)} \right)$$

- We define the conditional mass function of X given Y as

$$f_{X|Y}(x_i|y_j) = \frac{f_{XY}(x_i, y_j)}{f_Y(y_j)} = P[X = x_i | Y = y_j]$$

Conditional mass function

- We got

$$F_{X|Y}(x|y_j) = \sum_{i:x_i \leq x} \left(\frac{f_{XY}(x_i, y_j)}{f_Y(y_j)} \right)$$

- We define the conditional mass function of X given Y as

$$f_{X|Y}(x_i|y_j) = \frac{f_{XY}(x_i, y_j)}{f_Y(y_j)} = P[X = x_i | Y = y_j]$$

- Note that

$$\sum_i f_{X|Y}(x_i|y_j) = 1, \quad \forall y_j; \quad \text{and} \quad F_{X|Y}(x|y_j) = \sum_{i:x_i \leq x} f_{X|Y}(x_i|y_j)$$

Example: Conditional pmf

- ▶ Consider the random experiment of tossing a coin n times.

Example: Conditional pmf

- ▶ Consider the random experiment of tossing a coin n times.
- ▶ Let X denote the number of heads and let Y denote the toss number on which the first head comes.

Example: Conditional pmf

- ▶ Consider the random experiment of tossing a coin n times.
- ▶ Let X denote the number of heads and let Y denote the toss number on which the first head comes.
- ▶ For $1 \leq k \leq n$

Example: Conditional pmf

- ▶ Consider the random experiment of tossing a coin n times.
- ▶ Let X denote the number of heads and let Y denote the toss number on which the first head comes.
- ▶ For $1 \leq k \leq n$

$$f_{Y|X}(k|1) = P[Y = k|X = 1] = \frac{P[Y = k, X = 1]}{P[X = 1]}$$

Example: Conditional pmf

- ▶ Consider the random experiment of tossing a coin n times.
- ▶ Let X denote the number of heads and let Y denote the toss number on which the first head comes.
- ▶ For $1 \leq k \leq n$

$$\begin{aligned} f_{Y|X}(k|1) &= P[Y = k|X = 1] = \frac{P[Y = k, X = 1]}{P[X = 1]} \\ &= \frac{p(1-p)^{n-1}}{{}^nC_1 p(1-p)^{n-1}} \end{aligned}$$

Example: Conditional pmf

- ▶ Consider the random experiment of tossing a coin n times.
- ▶ Let X denote the number of heads and let Y denote the toss number on which the first head comes.
- ▶ For $1 \leq k \leq n$

$$\begin{aligned}f_{Y|X}(k|1) &= P[Y = k|X = 1] = \frac{P[Y = k, X = 1]}{P[X = 1]} \\&= \frac{p(1-p)^{n-1}}{{}^nC_1 p(1-p)^{n-1}} \\&= \frac{1}{n}\end{aligned}$$

Example: Conditional pmf

- ▶ Consider the random experiment of tossing a coin n times.
- ▶ Let X denote the number of heads and let Y denote the toss number on which the first head comes.
- ▶ For $1 \leq k \leq n$

$$\begin{aligned}f_{Y|X}(k|1) &= P[Y = k|X = 1] = \frac{P[Y = k, X = 1]}{P[X = 1]} \\&= \frac{p(1-p)^{n-1}}{{}^nC_1 p(1-p)^{n-1}} \\&= \frac{1}{n}\end{aligned}$$

- ▶ Given there is only one head, it is equally likely to occur on any toss.

- The conditional mass function is

$$f_{X|Y}(x_i|y_j) = P[X = x_i|Y = y_j] = \frac{f_{XY}(x_i, y_j)}{f_Y(y_j)}$$

- ▶ The conditional mass function is

$$f_{X|Y}(x_i|y_j) = P[X = x_i|Y = y_j] = \frac{f_{XY}(x_i, y_j)}{f_Y(y_j)}$$

- ▶ This gives us the useful identity

$$f_{XY}(x_i, y_j) = f_{X|Y}(x_i|y_j)f_Y(y_j)$$

- ▶ The conditional mass function is

$$f_{X|Y}(x_i|y_j) = P[X = x_i|Y = y_j] = \frac{f_{XY}(x_i, y_j)}{f_Y(y_j)}$$

- ▶ This gives us the useful identity

$$f_{XY}(x_i, y_j) = f_{X|Y}(x_i|y_j)f_Y(y_j)$$
$$(P[X = x_i, Y = y_j] = P[X = x_i|Y = y_j]P[Y = y_j])$$

- ▶ The conditional mass function is

$$f_{X|Y}(x_i|y_j) = P[X = x_i|Y = y_j] = \frac{f_{XY}(x_i, y_j)}{f_Y(y_j)}$$

- ▶ This gives us the useful identity

$$f_{XY}(x_i, y_j) = f_{X|Y}(x_i|y_j)f_Y(y_j)$$

$$(P[X = x_i, Y = y_j] = P[X = x_i|Y = y_j]P[Y = y_j])$$

- ▶ This gives us the total probability rule for discrete rv's

$$f_X(x_i) = \sum_j f_{XY}(x_i, y_j)$$

- ▶ The conditional mass function is

$$f_{X|Y}(x_i|y_j) = P[X = x_i|Y = y_j] = \frac{f_{XY}(x_i, y_j)}{f_Y(y_j)}$$

- ▶ This gives us the useful identity

$$f_{XY}(x_i, y_j) = f_{X|Y}(x_i|y_j)f_Y(y_j)$$

$$(P[X = x_i, Y = y_j] = P[X = x_i|Y = y_j]P[Y = y_j])$$

- ▶ This gives us the total probability rule for discrete rv's

$$f_X(x_i) = \sum_j f_{XY}(x_i, y_j) = \sum_j f_{X|Y}(x_i|y_j)f_Y(y_j)$$

- ▶ The conditional mass function is

$$f_{X|Y}(x_i|y_j) = P[X = x_i|Y = y_j] = \frac{f_{XY}(x_i, y_j)}{f_Y(y_j)}$$

- ▶ This gives us the useful identity

$$f_{XY}(x_i, y_j) = f_{X|Y}(x_i|y_j)f_Y(y_j)$$

$$(P[X = x_i, Y = y_j] = P[X = x_i|Y = y_j]P[Y = y_j])$$

- ▶ This gives us the total probability rule for discrete rv's

$$f_X(x_i) = \sum_j f_{XY}(x_i, y_j) = \sum_j f_{X|Y}(x_i|y_j)f_Y(y_j)$$

- ▶ This is same as

$$P[X = x_i] = \sum_j P[X = x_i|Y = y_j]P[Y = y_j]$$

- ▶ The conditional mass function is

$$f_{X|Y}(x_i|y_j) = P[X = x_i|Y = y_j] = \frac{f_{XY}(x_i, y_j)}{f_Y(y_j)}$$

- ▶ This gives us the useful identity

$$f_{XY}(x_i, y_j) = f_{X|Y}(x_i|y_j)f_Y(y_j)$$

$$(P[X = x_i, Y = y_j] = P[X = x_i|Y = y_j]P[Y = y_j])$$

- ▶ This gives us the total probability rule for discrete rv's

$$f_X(x_i) = \sum_j f_{XY}(x_i, y_j) = \sum_j f_{X|Y}(x_i|y_j)f_Y(y_j)$$

- ▶ This is same as

$$P[X = x_i] = \sum_j P[X = x_i|Y = y_j]P[Y = y_j]$$

$(P(A) = \sum_j P(A|B_j)P(B_j)$ when B_1, \dots form a partition)

Bayes Rule for discrete Random Variable

► We have

$$f_{XY}(x_i, y_j) = f_{X|Y}(x_i|y_j)f_Y(y_j)$$

Bayes Rule for discrete Random Variable

► We have

$$f_{XY}(x_i, y_j) = f_{X|Y}(x_i|y_j)f_Y(y_j) = f_{Y|X}(y_j|x_i)f_X(x_i)$$

Bayes Rule for discrete Random Variable

- We have

$$f_{XY}(x_i, y_j) = f_{X|Y}(x_i|y_j)f_Y(y_j) = f_{Y|X}(y_j|x_i)f_X(x_i)$$

- This gives us Bayes rule for discrete rv's

Bayes Rule for discrete Random Variable

- We have

$$f_{XY}(x_i, y_j) = f_{X|Y}(x_i|y_j)f_Y(y_j) = f_{Y|X}(y_j|x_i)f_X(x_i)$$

- This gives us Bayes rule for discrete rv's

$$f_{X|Y}(x_i|y_j) = \frac{f_{Y|X}(y_j|x_i)f_X(x_i)}{f_Y(y_j)}$$

Bayes Rule for discrete Random Variable

- We have

$$f_{XY}(x_i, y_j) = f_{X|Y}(x_i|y_j)f_Y(y_j) = f_{Y|X}(y_j|x_i)f_X(x_i)$$

- This gives us Bayes rule for discrete rv's

$$\begin{aligned} f_{X|Y}(x_i|y_j) &= \frac{f_{Y|X}(y_j|x_i)f_X(x_i)}{f_Y(y_j)} \\ &= \frac{f_{Y|X}(y_j|x_i)f_X(x_i)}{\sum_i f_{XY}(x_i, y_j)} \end{aligned}$$

Bayes Rule for discrete Random Variable

- We have

$$f_{XY}(x_i, y_j) = f_{X|Y}(x_i|y_j)f_Y(y_j) = f_{Y|X}(y_j|x_i)f_X(x_i)$$

- This gives us Bayes rule for discrete rv's

$$\begin{aligned} f_{X|Y}(x_i|y_j) &= \frac{f_{Y|X}(y_j|x_i)f_X(x_i)}{f_Y(y_j)} \\ &= \frac{f_{Y|X}(y_j|x_i)f_X(x_i)}{\sum_i f_{XY}(x_i, y_j)} \\ &= \frac{f_{Y|X}(y_j|x_i)f_X(x_i)}{\sum_i f_{Y|X}(y_j|x_i)f_X(x_i)} \end{aligned}$$

- ▶ Let X, Y be continuous rv's with joint density, f_{XY} .

- ▶ Let X, Y be continuous rv's with joint density, f_{XY} .
- ▶ We once again want to define conditional df

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ Let X, Y be continuous rv's with joint density, f_{XY} .
- ▶ We once again want to define conditional df

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ But the conditioning event, $[Y = y]$ has zero probability.

- ▶ Let X, Y be continuous rv's with joint density, f_{XY} .
- ▶ We once again want to define conditional df

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ But the conditioning event, $[Y = y]$ has zero probability.
- ▶ Hence we define conditional df as follows

- ▶ Let X, Y be continuous rv's with joint density, f_{XY} .
- ▶ We once again want to define conditional df

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ But the conditioning event, $[Y = y]$ has zero probability.
- ▶ Hence we define conditional df as follows

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

- ▶ Let X, Y be continuous rv's with joint density, f_{XY} .
- ▶ We once again want to define conditional df

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ But the conditioning event, $[Y = y]$ has zero probability.
- ▶ Hence we define conditional df as follows

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

- ▶ This is well defined if the limit exists.

- ▶ Let X, Y be continuous rv's with joint density, f_{XY} .
- ▶ We once again want to define conditional df

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ But the conditioning event, $[Y = y]$ has zero probability.
- ▶ Hence we define conditional df as follows

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

- ▶ This is well defined if the limit exists.
- ▶ The limit exists for all y where $f_Y(y) > 0$ (and for all x)

- ▶ The conditional df is given by (assuming $f_Y(y) > 0$)

- The conditional df is given by (assuming $f_Y(y) > 0$)

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

- The conditional df is given by (assuming $f_Y(y) > 0$)

$$\begin{aligned} F_{X|Y}(x|y) &= \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]] \\ &= \lim_{\delta \rightarrow 0} \frac{P[X \leq x, Y \in [y, y + \delta]]}{P[Y \in [y, y + \delta]]} \end{aligned}$$

- The conditional df is given by (assuming $f_Y(y) > 0$)

$$\begin{aligned} F_{X|Y}(x|y) &= \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]] \\ &= \lim_{\delta \rightarrow 0} \frac{P[X \leq x, Y \in [y, y + \delta]]}{P[Y \in [y, y + \delta]]} \\ &= \lim_{\delta \rightarrow 0} \frac{\int_{-\infty}^x \int_y^{y+\delta} f_{XY}(x', y') dy' dx'}{\int_y^{y+\delta} f_Y(y') dy'} \end{aligned}$$

- The conditional df is given by (assuming $f_Y(y) > 0$)

$$\begin{aligned} F_{X|Y}(x|y) &= \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]] \\ &= \lim_{\delta \rightarrow 0} \frac{P[X \leq x, Y \in [y, y + \delta]]}{P[Y \in [y, y + \delta]]} \\ &= \lim_{\delta \rightarrow 0} \frac{\int_{-\infty}^x \int_y^{y+\delta} f_{XY}(x', y') dy' dx'}{\int_y^{y+\delta} f_Y(y') dy'} \\ &= \lim_{\delta \rightarrow 0} \frac{\int_{-\infty}^x f_{XY}(x', y) \delta dx'}{f_Y(y) \delta} \end{aligned}$$

- The conditional df is given by (assuming $f_Y(y) > 0$)

$$\begin{aligned} F_{X|Y}(x|y) &= \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]] \\ &= \lim_{\delta \rightarrow 0} \frac{P[X \leq x, Y \in [y, y + \delta]]}{P[Y \in [y, y + \delta]]} \\ &= \lim_{\delta \rightarrow 0} \frac{\int_{-\infty}^x \int_y^{y+\delta} f_{XY}(x', y') dy' dx'}{\int_y^{y+\delta} f_Y(y') dy'} \\ &= \lim_{\delta \rightarrow 0} \frac{\int_{-\infty}^x f_{XY}(x', y) \delta dx'}{f_Y(y) \delta} \\ &= \int_{-\infty}^x \frac{f_{XY}(x', y)}{f_Y(y)} dx' \end{aligned}$$

- The conditional df is given by (assuming $f_Y(y) > 0$)

$$\begin{aligned} F_{X|Y}(x|y) &= \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]] \\ &= \lim_{\delta \rightarrow 0} \frac{P[X \leq x, Y \in [y, y + \delta]]}{P[Y \in [y, y + \delta]]} \\ &= \lim_{\delta \rightarrow 0} \frac{\int_{-\infty}^x \int_y^{y+\delta} f_{XY}(x', y') dy' dx'}{\int_y^{y+\delta} f_Y(y') dy'} \\ &= \lim_{\delta \rightarrow 0} \frac{\int_{-\infty}^x f_{XY}(x', y) \delta dx'}{f_Y(y) \delta} \\ &= \int_{-\infty}^x \frac{f_{XY}(x', y)}{f_Y(y)} dx' \end{aligned}$$

- We define conditional density of X given Y as

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)}$$

- ▶ Let X, Y have joint density f_{XY} .

- ▶ Let X, Y have joint density f_{XY} .
- ▶ The conditional df of X given Y is

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

- ▶ Let X, Y have joint density f_{XY} .
- ▶ The conditional df of X given Y is

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

- ▶ This exists if $f_Y(y) > 0$

- ▶ Let X, Y have joint density f_{XY} .
- ▶ The conditional df of X given Y is

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

- ▶ This exists if $f_Y(y) > 0$

$$F_{X|Y}(x|y) = \int_{-\infty}^x \frac{f_{XY}(x', y)}{f_Y(y)} dx'$$

- ▶ Let X, Y have joint density f_{XY} .
- ▶ The conditional df of X given Y is

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

- ▶ This exists if $f_Y(y) > 0$ and then it has a density:

$$F_{X|Y}(x|y) = \int_{-\infty}^x \frac{f_{XY}(x', y)}{f_Y(y)} dx' = \int_{-\infty}^x f_{X|Y}(x'|y) dx'$$

- ▶ Let X, Y have joint density f_{XY} .
- ▶ The conditional df of X given Y is

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

- ▶ This exists if $f_Y(y) > 0$ and then it has a density:

$$F_{X|Y}(x|y) = \int_{-\infty}^x \frac{f_{XY}(x', y)}{f_Y(y)} dx' = \int_{-\infty}^x f_{X|Y}(x'|y) dx'$$

- ▶ This conditional density is given by

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)}$$

- ▶ Let X, Y have joint density f_{XY} .
- ▶ The conditional df of X given Y is

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

- ▶ This exists if $f_Y(y) > 0$ and then it has a density:

$$F_{X|Y}(x|y) = \int_{-\infty}^x \frac{f_{XY}(x', y)}{f_Y(y)} dx' = \int_{-\infty}^x f_{X|Y}(x'|y) dx'$$

- ▶ This conditional density is given by

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)}$$

- ▶ We (once again) have the useful identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y) = f_{Y|X}(y|x) f_X(x)$$

Example

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

Example

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- We saw that the marginal densities are

$$f_X(x) = 2(1 - x), \quad 0 < x < 1; \quad f_Y(y) = 2y, \quad 0 < y < 1$$

Example

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We saw that the marginal densities are

$$f_X(x) = 2(1 - x), \quad 0 < x < 1; \quad f_Y(y) = 2y, \quad 0 < y < 1$$

- ▶ Hence the conditional densities are given by

Example

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We saw that the marginal densities are

$$f_X(x) = 2(1 - x), \quad 0 < x < 1; \quad f_Y(y) = 2y, \quad 0 < y < 1$$

- ▶ Hence the conditional densities are given by

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)}$$

Example

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We saw that the marginal densities are

$$f_X(x) = 2(1 - x), \quad 0 < x < 1; \quad f_Y(y) = 2y, \quad 0 < y < 1$$

- ▶ Hence the conditional densities are given by

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{1}{y}, \quad 0 < x < y < 1$$

Example

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We saw that the marginal densities are

$$f_X(x) = 2(1 - x), \quad 0 < x < 1; \quad f_Y(y) = 2y, \quad 0 < y < 1$$

- ▶ Hence the conditional densities are given by

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{1}{y}, \quad 0 < x < y < 1$$

$$f_{Y|X}(y|x) = \frac{f_{XY}(x, y)}{f_X(x)}$$

Example

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We saw that the marginal densities are

$$f_X(x) = 2(1 - x), \quad 0 < x < 1; \quad f_Y(y) = 2y, \quad 0 < y < 1$$

- ▶ Hence the conditional densities are given by

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{1}{y}, \quad 0 < x < y < 1$$

$$f_{Y|X}(y|x) = \frac{f_{XY}(x, y)}{f_X(x)} = \frac{1}{1 - x}, \quad 0 < x < y < 1$$

Example

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We saw that the marginal densities are


$$f_X(x) = 2(1 - x), \quad 0 < x < 1; \quad f_Y(y) = 2y, \quad 0 < y < 1$$

- ▶ Hence the conditional densities are given by

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{1}{y}, \quad 0 < x < y < 1$$

$$f_{Y|X}(y|x) = \frac{f_{XY}(x, y)}{f_X(x)} = \frac{1}{1 - x}, \quad 0 < x < y < 1$$

- ▶ We can see this intuitively like this



Example

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We saw that the marginal densities are


$$f_X(x) = 2(1 - x), \quad 0 < x < 1; \quad f_Y(y) = 2y, \quad 0 < y < 1$$

- ▶ Hence the conditional densities are given by

$$f_{X|Y}(x|y) = \frac{f_{XY}(x, y)}{f_Y(y)} = \frac{1}{y}, \quad 0 < x < y < 1$$

$$f_{Y|X}(y|x) = \frac{f_{XY}(x, y)}{f_X(x)} = \frac{1}{1 - x}, \quad 0 < x < y < 1$$

- ▶ We can see this intuitively like this



- ▶ The identity $f_{XY}(x, y) = f_{X|Y}(x|y)f_Y(y)$ can be used to specify the joint density of two continuous rv's

- ▶ The identity $f_{XY}(x, y) = f_{X|Y}(x|y)f_Y(y)$ can be used to specify the joint density of two continuous rv's
- ▶ We can specify the marginal density of one and the conditional density of the other given the first.

- ▶ The identity $f_{XY}(x, y) = f_{X|Y}(x|y)f_Y(y)$ can be used to specify the joint density of two continuous rv's
- ▶ We can specify the marginal density of one and the conditional density of the other given the first.
- ▶ This may actually be the model of how the the rv's are generated.

Example

- ▶ Let X be uniform over $(0, 1)$ and let Y be uniform over 0 to X . Find the density of Y .

Example

- ▶ Let X be uniform over $(0, 1)$ and let Y be uniform over 0 to X . Find the density of Y .
- ▶ What we are given is

$$f_X(x) = 1, 0 < x < 1; \quad f_{Y|X}(y|x) = \frac{1}{x}, 0 < y < x < 1$$

Example

- ▶ Let X be uniform over $(0, 1)$ and let Y be uniform over 0 to X . Find the density of Y .
- ▶ What we are given is

$$f_X(x) = 1, 0 < x < 1; \quad f_{Y|X}(y|x) = \frac{1}{x}, 0 < y < x < 1$$

- ▶ Hence the joint density is:
 $f_{XY}(x, y) = \frac{1}{x}, 0 < y < x < 1.$

Example

- ▶ Let X be uniform over $(0, 1)$ and let Y be uniform over 0 to X . Find the density of Y .
- ▶ What we are given is

$$f_X(x) = 1, 0 < x < 1; \quad f_{Y|X}(y|x) = \frac{1}{x}, 0 < y < x < 1$$

- ▶ Hence the joint density is:
 $f_{XY}(x, y) = \frac{1}{x}, 0 < y < x < 1.$
- ▶ Hence the density of Y is

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx$$

Example

- ▶ Let X be uniform over $(0, 1)$ and let Y be uniform over 0 to X . Find the density of Y .
- ▶ What we are given is

$$f_X(x) = 1, 0 < x < 1; \quad f_{Y|X}(y|x) = \frac{1}{x}, 0 < y < x < 1$$

- ▶ Hence the joint density is:
 $f_{XY}(x, y) = \frac{1}{x}, 0 < y < x < 1.$
- ▶ Hence the density of Y is

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx = \int_y^1 \frac{1}{x} dx$$

Example

- ▶ Let X be uniform over $(0, 1)$ and let Y be uniform over 0 to X . Find the density of Y .
- ▶ What we are given is

$$f_X(x) = 1, 0 < x < 1; \quad f_{Y|X}(y|x) = \frac{1}{x}, 0 < y < x < 1$$

- ▶ Hence the joint density is:
 $f_{XY}(x, y) = \frac{1}{x}, 0 < y < x < 1.$
- ▶ Hence the density of Y is

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx = \int_y^1 \frac{1}{x} dx = -\ln(y), 0 < y < 1$$

Example

- ▶ Let X be uniform over $(0, 1)$ and let Y be uniform over 0 to X . Find the density of Y .
- ▶ What we are given is

$$f_X(x) = 1, 0 < x < 1; \quad f_{Y|X}(y|x) = \frac{1}{x}, 0 < y < x < 1$$

- ▶ Hence the joint density is:
 $f_{XY}(x, y) = \frac{1}{x}, 0 < y < x < 1.$
- ▶ Hence the density of Y is

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x, y) dx = \int_y^1 \frac{1}{x} dx = -\ln(y), 0 < y < 1$$

- ▶ We can verify it to be a density

$$-\int_0^1 \ln(y) dy = -y \ln(y)|_0^1 + \int_0^1 y \frac{1}{y} dy = 1$$

- ▶ We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y)$$

- ▶ We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y)$$

- ▶ By integrating both sides

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_{-\infty}^{\infty} f_{X|Y}(x|y) f_Y(y) dy$$

- ▶ We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y)$$

- ▶ By integrating both sides

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_{-\infty}^{\infty} f_{X|Y}(x|y) f_Y(y) dy$$

- ▶ This is a continuous analogue of total probability rule.

- ▶ We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y)$$

- ▶ By integrating both sides

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_{-\infty}^{\infty} f_{X|Y}(x|y) f_Y(y) dy$$

- ▶ This is a continuous analogue of total probability rule.
- ▶ But note that, since X is continuous rv, $f_X(x)$ is **NOT** $P[X = x]$

- ▶ We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y)$$

- ▶ By integrating both sides

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_{-\infty}^{\infty} f_{X|Y}(x|y) f_Y(y) dy$$

- ▶ This is a continuous analogue of total probability rule.
- ▶ But note that, since X is continuous rv, $f_X(x)$ is **NOT** $P[X = x]$
- ▶ In case of discrete rv, the mass function value $f_X(x)$ is equal to $P[X = x]$ and we had

$$f_X(x) = \sum_y f_{X|Y}(x|y) f_Y(y)$$

- ▶ We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y)$$

- ▶ By integrating both sides

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_{-\infty}^{\infty} f_{X|Y}(x|y) f_Y(y) dy$$

- ▶ This is a continuous analogue of total probability rule.
- ▶ But note that, since X is continuous rv, $f_X(x)$ is **NOT** $P[X = x]$
- ▶ In case of discrete rv, the mass function value $f_X(x)$ is equal to $P[X = x]$ and we had

$$f_X(x) = \sum_y f_{X|Y}(x|y) f_Y(y)$$

- ▶ It is as if one can simply replace pmf by pdf and summation by integration!!

- ▶ We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y)$$

- ▶ By integrating both sides

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy = \int_{-\infty}^{\infty} f_{X|Y}(x|y) f_Y(y) dy$$

- ▶ This is a continuous analogue of total probability rule.
- ▶ But note that, since X is continuous rv, $f_X(x)$ is **NOT** $P[X = x]$
- ▶ In case of discrete rv, the mass function value $f_X(x)$ is equal to $P[X = x]$ and we had

$$f_X(x) = \sum_y f_{X|Y}(x|y) f_Y(y)$$

- ▶ It is as if one can simply replace pmf by pdf and summation by integration!!
- ▶ While often that gives the right result, one needs to be very careful

- We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y)$$

- We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y) = f_{Y|X}(y|x) f_X(x)$$

- ▶ We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y) = f_{Y|X}(y|x) f_X(x)$$

- ▶ This gives rise to Bayes rule for continuous rv

- We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y) = f_{Y|X}(y|x) f_X(x)$$

- This gives rise to Bayes rule for continuous rv

$$f_{X|Y}(x|y) = \frac{f_{Y|X}(y|x) f_X(x)}{f_Y(y)}$$

- We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y) = f_{Y|X}(y|x) f_X(x)$$

- This gives rise to Bayes rule for continuous rv

$$\begin{aligned} f_{X|Y}(x|y) &= \frac{f_{Y|X}(y|x) f_X(x)}{f_Y(y)} \\ &= \frac{f_{Y|X}(y|x) f_X(x)}{\int_{-\infty}^{\infty} f_{Y|X}(y|x) f_X(x) dx} \end{aligned}$$

- ▶ We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y) = f_{Y|X}(y|x) f_X(x)$$

- ▶ This gives rise to Bayes rule for continuous rv

$$\begin{aligned} f_{X|Y}(x|y) &= \frac{f_{Y|X}(y|x) f_X(x)}{f_Y(y)} \\ &= \frac{f_{Y|X}(y|x) f_X(x)}{\int_{-\infty}^{\infty} f_{Y|X}(y|x) f_X(x) dx} \end{aligned}$$

- ▶ This is essentially identical to Bayes rule for discrete rv's.

- ▶ We have the identity

$$f_{XY}(x, y) = f_{X|Y}(x|y) f_Y(y) = f_{Y|X}(y|x) f_X(x)$$

- ▶ This gives rise to Bayes rule for continuous rv

$$\begin{aligned} f_{X|Y}(x|y) &= \frac{f_{Y|X}(y|x) f_X(x)}{f_Y(y)} \\ &= \frac{f_{Y|X}(y|x) f_X(x)}{\int_{-\infty}^{\infty} f_{Y|X}(y|x) f_X(x) dx} \end{aligned}$$

- ▶ This is essentially identical to Bayes rule for discrete rv's. We have essentially put the pdf wherever there was pmf

- ▶ To recap, we started by defining conditional distribution function.

- ▶ To recap, we started by defining conditional distribution function.

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ To recap, we started by defining conditional distribution function.

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ When X, Y are discrete, we define this only for $y = y_j$.

- ▶ To recap, we started by defining conditional distribution function.

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ When X, Y are discrete, we define this only for $y = y_j$. That is, we define it only for all values that Y can take.

- ▶ To recap, we started by defining conditional distribution function.

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ When X, Y are discrete, we define this only for $y = y_j$. That is, we define it only for all values that Y can take.
- ▶ When X, Y have joint density, we defined it by

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

- ▶ To recap, we started by defining conditional distribution function.

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ When X, Y are discrete, we define this only for $y = y_j$. That is, we define it only for all values that Y can take.
- ▶ When X, Y have joint density, we defined it by

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

This limit exists and $F_{X|Y}$ is well defined if $f_Y(y) > 0$.

- ▶ To recap, we started by defining conditional distribution function.

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ When X, Y are discrete, we define this only for $y = y_j$. That is, we define it only for all values that Y can take.
- ▶ When X, Y have joint density, we defined it by

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

This limit exists and $F_{X|Y}$ is well defined if $f_Y(y) > 0$. That is, essentially again for all values that Y can take.

- ▶ To recap, we started by defining conditional distribution function.

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ When X, Y are discrete, we define this only for $y = y_j$. That is, we define it only for all values that Y can take.
- ▶ When X, Y have joint density, we defined it by

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

This limit exists and $F_{X|Y}$ is well defined if $f_Y(y) > 0$. That is, essentially again for all values that Y can take.

- ▶ In the discrete case, we define $f_{X|Y}$ as the pmf corresponding to $F_{X|Y}$. This conditional pmf can also be defined as a conditional probability

- ▶ To recap, we started by defining conditional distribution function.

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ When X, Y are discrete, we define this only for $y = y_j$. That is, we define it only for all values that Y can take.
- ▶ When X, Y have joint density, we defined it by

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

This limit exists and $F_{X|Y}$ is well defined if $f_Y(y) > 0$. That is, essentially again for all values that Y can take.

- ▶ In the discrete case, we define $f_{X|Y}$ as the pmf corresponding to $F_{X|Y}$. This conditional pmf can also be defined as a conditional probability
- ▶ In the continuous case $f_{X|Y}$ is the density corresponding to $F_{X|Y}$.

- ▶ To recap, we started by defining conditional distribution function.

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ When X, Y are discrete, we define this only for $y = y_j$. That is, we define it only for all values that Y can take.
- ▶ When X, Y have joint density, we defined it by

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

This limit exists and $F_{X|Y}$ is well defined if $f_Y(y) > 0$. That is, essentially again for all values that Y can take.

- ▶ In the discrete case, we define $f_{X|Y}$ as the pmf corresponding to $F_{X|Y}$. This conditional pmf can also be defined as a conditional probability
- ▶ In the continuous case $f_{X|Y}$ is the density corresponding to $F_{X|Y}$.
- ▶ In both cases we have: $f_{XY}(x, y) = f_{X|Y}(x|y)f_Y(y)$

- ▶ To recap, we started by defining conditional distribution function.

$$F_{X|Y}(x|y) = P[X \leq x | Y = y]$$

- ▶ When X, Y are discrete, we define this only for $y = y_j$. That is, we define it only for all values that Y can take.
- ▶ When X, Y have joint density, we defined it by

$$F_{X|Y}(x|y) = \lim_{\delta \rightarrow 0} P[X \leq x | Y \in [y, y + \delta]]$$

This limit exists and $F_{X|Y}$ is well defined if $f_Y(y) > 0$. That is, essentially again for all values that Y can take.

- ▶ In the discrete case, we define $f_{X|Y}$ as the pmf corresponding to $F_{X|Y}$. This conditional pmf can also be defined as a conditional probability
- ▶ In the continuous case $f_{X|Y}$ is the density corresponding to $F_{X|Y}$.
- ▶ In both cases we have: $f_{XY}(x, y) = f_{X|Y}(x|y)f_Y(y)$
- ▶ This gives total probability rule and Bayes rule for random variables