

## Recap: Functions of multiple rv

- ▶ Given  $X_1, \dots, X_n$ , random variables on the same probability space,  $Z = g(X_1, \dots, X_n)$  is a rv

## Recap: Functions of multiple rv

- ▶ Given  $X_1, \dots, X_n$ , random variables on the same probability space,  $Z = g(X_1, \dots, X_n)$  is a rv (if  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  is borel measurable).

## Recap: Functions of multiple rv

- ▶ Given  $X_1, \dots, X_n$ , random variables on the same probability space,  $Z = g(X_1, \dots, X_n)$  is a rv (if  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  is borel measurable).
- ▶ We can determine distribution of  $Z$  from the joint distribution of all  $X_i$

## Recap: Functions of multiple rv

- ▶ Given  $X_1, \dots, X_n$ , random variables on the same probability space,  $Z = g(X_1, \dots, X_n)$  is a rv (if  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  is borel measurable).
- ▶ We can determine distribution of  $Z$  from the joint distribution of all  $X_i$

$$F_Z(z) = P[Z \leq z] = P[g(X_1, \dots, X_n) \leq z]$$

## Recap: iid random variables

- ▶  $X_1, \dots, X_n$  are said to be independent if events  $[X_1 \in B_1], \dots, [X_n \in B_n]$  are independent.

## Recap: iid random variables

- ▶  $X_1, \dots, X_n$  are said to be independent if events  $[X_1 \in B_1], \dots, [X_n \in B_n]$  are independent.
- ▶ If  $X_1, \dots, X_n$  are independent and all of them have the same distribution function then they are said to be iid – independent and identically distributed

# Recap: Independence of functions of random variables

## Recap: Independence of functions of random variables

- ▶ If  $\mathbf{X}, \mathbf{Y}$  are vector random variables (or random vectors), independence implies  $[\mathbf{X} \in B_1]$  is independent of  $[\mathbf{Y} \in B_2]$  for all borel sets  $B_1, B_2$  (in appropriate spaces).



## Recap: Independence of functions of random variables

- ▶ If  $\mathbf{X}, \mathbf{Y}$  are vector random variables (or random vectors), independence implies  $[\mathbf{X} \in B_1]$  is independent of  $[\mathbf{Y} \in B_2]$  for all borel sets  $B_1, B_2$  (in appropriate spaces).
- ▶ Then  $g(\mathbf{X})$  would be independent of  $h(\mathbf{Y})$ .

# Recap: Independence of functions of random variables

- ▶ If  $\mathbf{X}, \mathbf{Y}$  are vector random variables (or random vectors), independence implies  $[\mathbf{X} \in B_1]$  is independent of  $[\mathbf{Y} \in B_2]$  for all borel sets  $B_1, B_2$  (in appropriate spaces).
- ▶ Then  $g(\mathbf{X})$  would be independent of  $h(\mathbf{Y})$ .
- ▶ That is, suppose  $X_1, \dots, X_m, X_{m+1}, \dots, X_{m+n}$  are independent.

# Recap: Independence of functions of random variables

- ▶ If  $\mathbf{X}, \mathbf{Y}$  are vector random variables (or random vectors), independence implies  $[\mathbf{X} \in B_1]$  is independent of  $[\mathbf{Y} \in B_2]$  for all borel sets  $B_1, B_2$  (in appropriate spaces).
- ▶ Then  $g(\mathbf{X})$  would be independent of  $h(\mathbf{Y})$ .
- ▶ That is, suppose  $X_1, \dots, X_m, X_{m+1}, \dots, X_{m+n}$  are independent.
- ▶ Then,  $g(X_1, \dots, X_m)$  is independent of  $h(X_{m+1}, \dots, X_{m+n})$ .

## Recap: Max of a set of random variables

- ▶ Let  $X_1, \dots, X_n$  be independent and  $Z = \max(X_1, \dots, X_n)$

# Recap: Max of a set of random variables

- ▶ Let  $X_1, \dots, X_n$  be independent and  $Z = \max(X_1, \dots, X_n)$

$$F_Z(z) = \prod_{i=1}^n F_{X_i}(z)$$

# Recap: Max of a set of random variables

- ▶ Let  $X_1, \dots, X_n$  be independent and  $Z = \max(X_1, \dots, X_n)$

$$\begin{aligned} F_Z(z) &= \prod_{i=1}^n F_{X_i}(z) \\ &= (F(z))^n, \quad \text{if they are iid} \end{aligned}$$

# Recap: Min of a set of random variables

- ▶ Let  $X_1, \dots, X_n$  be independent and  $Z = \min(X_1, \dots, X_n)$

# Recap: Min of a set of random variables

- ▶ Let  $X_1, \dots, X_n$  be independent and  $Z = \min(X_1, \dots, X_n)$

$$F_Z(z) = 1 - \prod_{i=1}^n (1 - F_{X_i}(z))$$



# Recap: Min of a set of random variables

- ▶ Let  $X_1, \dots, X_n$  be independent and  $Z = \min(X_1, \dots, X_n)$

$$\begin{aligned} F_Z(z) &= 1 - \prod_{i=1}^n (1 - F_{X_i}(z)) \\ &= 1 - (1 - F(z))^n, \quad \text{if they are iid} \end{aligned}$$

# Recap: Order Statistics

- ▶ Let  $X_1, \dots, X_n$  be iid with density  $f$ .

# Recap: Order Statistics

- ▶ Let  $X_1, \dots, X_n$  be iid with density  $f$ .
- ▶ Let  $X_{(k)}$  denote the  $k^{th}$  smallest of these.

# Recap: Order Statistics

- ▶ Let  $X_1, \dots, X_n$  be iid with density  $f$ .
- ▶ Let  $X_{(k)}$  denote the  $k^{th}$  smallest of these.
- ▶  $X_{(1)} = \min(X_1, \dots, X_n)$ ,  $X_{(n)} = \max(X_1, \dots, X_n)$

# Recap: Order Statistics

- ▶ Let  $X_1, \dots, X_n$  be iid with density  $f$ .
- ▶ Let  $X_{(k)}$  denote the  $k^{th}$  smallest of these.
- ▶  $X_{(1)} = \min(X_1, \dots, X_n)$ ,  $X_{(n)} = \max(X_1, \dots, X_n)$
- ▶ We have:

$$F_{X_{(k)}}(y) = \sum_{j=k}^n {}^nC_j (F(y))^j (1 - F(y))^{n-j}$$

# Recap: Order Statistics

- ▶ Let  $X_1, \dots, X_n$  be iid with density  $f$ .
- ▶ Let  $X_{(k)}$  denote the  $k^{th}$  smallest of these.
- ▶  $X_{(1)} = \min(X_1, \dots, X_n)$ ,  $X_{(n)} = \max(X_1, \dots, X_n)$
- ▶ We have:

$$F_{X_{(k)}}(y) = \sum_{j=k}^n {}^nC_j (F(y))^j (1 - F(y))^{n-j}$$

- ▶ Joint distribution of  $X_{(1)}, \dots, X_{(n)}$  is called the order statistics.

$$f_{X_{(1)} \dots X_{(n)}}(x_1, \dots, x_n) = n! \prod_{i=1}^n f(x_i), \quad x_1 < x_2 < \dots < x_n$$

## Recap: Sum of two independent rv

- ▶ Let  $X, Y$  be random variables and  $Z = X + Y$ .

## Recap: Sum of two independent rv

- ▶ Let  $X, Y$  be random variables and  $Z = X + Y$ .
- ▶ If  $X, Y$  are discrete

$$f_Z(z) = \sum_k f_{XY}(k, z - k)$$



## Recap: Sum of two independent rv

- ▶ Let  $X, Y$  be random variables and  $Z = X + Y$ .
- ▶ If  $X, Y$  are discrete

$$\begin{aligned}f_Z(z) &= \sum_k f_{XY}(k, z - k) \\&= \sum_k f_X(k) f_Y(z - k) \quad \text{when } X, Y \text{ are independent.}\end{aligned}$$

## Recap: Sum of two independent rv

- ▶ Let  $X, Y$  be random variables and  $Z = X + Y$ .
- ▶ If  $X, Y$  are discrete

$$\begin{aligned}f_Z(z) &= \sum_k f_{XY}(k, z - k) \\&= \sum_k f_X(k) f_Y(z - k) \quad \text{when } X, Y \text{ are independent.}\end{aligned}$$

- ▶ If  $X, Y$  have a joint density

$$f_Z(z) = \int_{-\infty}^{\infty} f_{XY}(t, z - t) dt$$

## Recap: Sum of two independent rv

- ▶ Let  $X, Y$  be random variables and  $Z = X + Y$ .
- ▶ If  $X, Y$  are discrete

$$\begin{aligned}f_Z(z) &= \sum_k f_{XY}(k, z - k) \\&= \sum_k f_X(k) f_Y(z - k) \quad \text{when } X, Y \text{ are independent.}\end{aligned}$$

- ▶ If  $X, Y$  have a joint density

$$\begin{aligned}f_Z(z) &= \int_{-\infty}^{\infty} f_{XY}(t, z - t) dt \\&= \int_{-\infty}^{\infty} f_X(t) f_Y(z - t) dt, \quad \text{when } X, Y \text{ are independent}\end{aligned}$$

## Recap: Sum of two independent rv

- ▶ Let  $X, Y$  be random variables and  $Z = X + Y$ .
- ▶ If  $X, Y$  are discrete

$$\begin{aligned}f_Z(z) &= \sum_k f_{XY}(k, z - k) \\&= \sum_k f_X(k) f_Y(z - k) \quad \text{when } X, Y \text{ are independent.}\end{aligned}$$

- ▶ If  $X, Y$  have a joint density

$$\begin{aligned}f_Z(z) &= \int_{-\infty}^{\infty} f_{XY}(t, z - t) dt \\&= \int_{-\infty}^{\infty} f_X(t) f_Y(z - t) dt, \quad \text{when } X, Y \text{ are independent}\end{aligned}$$

Density of sum of independent random variables is the convolution of their densities.

# Recap: sum of two independent random variables

- ▶ Given independent random variables  $X, Y$

## Recap: sum of two independent random variables

- ▶ Given independent random variables  $X, Y$
- ▶ If  $X \sim \text{Poisson}(\lambda_1)$  and  $Y \sim \text{Poisson}(\lambda_2)$   
 $X + Y \sim \text{Poisson}(\lambda_1 + \lambda_2)$

# Recap: sum of two independent random variables

- ▶ Given independent random variables  $X, Y$
- ▶ If  $X \sim \text{Poisson}(\lambda_1)$  and  $Y \sim \text{Poisson}(\lambda_2)$   
 $X + Y \sim \text{Poisson}(\lambda_1 + \lambda_2)$
- ▶ If  $X \sim \text{Gamma}(\alpha_1, \lambda)$  and  $Y \sim \text{Gamma}(\alpha_2, \lambda)$   
 $X + Y \sim \text{Gamma}(\alpha_1 + \alpha_2, \lambda)$

# Recap: sum of two independent random variables

- ▶ Given independent random variables  $X, Y$
- ▶ If  $X \sim \text{Poisson}(\lambda_1)$  and  $Y \sim \text{Poisson}(\lambda_2)$   
 $X + Y \sim \text{Poisson}(\lambda_1 + \lambda_2)$
- ▶ If  $X \sim \text{Gamma}(\alpha_1, \lambda)$  and  $Y \sim \text{Gamma}(\alpha_2, \lambda)$   
 $X + Y \sim \text{Gamma}(\alpha_1 + \alpha_2, \lambda)$
- ▶ If  $X \sim \mathcal{N}(\mu_1, \sigma_1^2)$  and  $Y \sim \mathcal{N}(\mu_2, \sigma_2^2)$   
 $X + Y \sim \mathcal{N}(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$



## Recap: Theorem for functions of cont rv

- ▶  $X_1, \dots, X_n$  are continuous rv with joint density

## Recap: Theorem for functions of cont rv

- ▶  $X_1, \dots, X_n$  are continuous rv with joint density

$$Y_1 = g_1(X_1, \dots, X_n) \quad \dots \quad Y_n = g_n(X_1, \dots, X_n)$$

# Recap: Theorem for functions of cont rv

- ▶  $X_1, \dots, X_n$  are continuous rv with joint density

$$Y_1 = g_1(X_1, \dots, X_n) \quad \dots \quad Y_n = g_n(X_1, \dots, X_n)$$

- ▶ The transformation is continuous with continuous first partials and is invertible and

$$X_1 = h_1(Y_1, \dots, Y_n) \quad \dots \quad X_n = h_n(Y_1, \dots, Y_n)$$

## Recap: Theorem for functions of cont rv

- ▶  $X_1, \dots, X_n$  are continuous rv with joint density

$$Y_1 = g_1(X_1, \dots, X_n) \quad \dots \quad Y_n = g_n(X_1, \dots, X_n)$$

- ▶ The transformation is continuous with continuous first partials and is invertible and

$$X_1 = h_1(Y_1, \dots, Y_n) \quad \dots \quad X_n = h_n(Y_1, \dots, Y_n)$$

- ▶ Jacobian of the inverse transform,  $J$ , is non-zero

## Recap: Theorem for functions of cont rv

- ▶  $X_1, \dots, X_n$  are continuous rv with joint density

$$Y_1 = g_1(X_1, \dots, X_n) \quad \dots \quad Y_n = g_n(X_1, \dots, X_n)$$

- ▶ The transformation is continuous with continuous first partials and is invertible and

$$X_1 = h_1(Y_1, \dots, Y_n) \quad \dots \quad X_n = h_n(Y_1, \dots, Y_n)$$

- ▶ Jacobian of the inverse transform,  $J$ , is non-zero
- ▶ Then the density of  $\mathbf{Y}$  is

$$f_{Y_1 \dots Y_n}(y_1, \dots, y_n) = |J| f_{X_1 \dots X_n}(h_1(y_1, \dots, y_n), \dots, h_n(y_1, \dots, y_n))$$

## Recap: Theorem for functions of cont rv

- ▶  $X_1, \dots, X_n$  are continuous rv with joint density

$$Y_1 = g_1(X_1, \dots, X_n) \quad \dots \quad Y_n = g_n(X_1, \dots, X_n)$$

- ▶ The transformation is continuous with continuous first partials and is invertible and

$$X_1 = h_1(Y_1, \dots, Y_n) \quad \dots \quad X_n = h_n(Y_1, \dots, Y_n)$$

- ▶ Jacobian of the inverse transform,  $J$ , is non-zero
- ▶ Then the density of  $\mathbf{Y}$  is

$$f_{Y_1 \dots Y_n}(y_1, \dots, y_n) = |J| f_{X_1 \dots X_n}(h_1(y_1, \dots, y_n), \dots, h_n(y_1, \dots, y_n))$$

- ▶ Called multidimensional change of variable formula

- Let  $X, Y$  have joint density  $f_{XY}$ . Let  $Z = X + Y$ .

- ▶ Let  $X, Y$  have joint density  $f_{XY}$ . Let  $Z = X + Y$ .
- ▶ We want to find  $f_Z$  using the theorem.



- ▶ Let  $X, Y$  have joint density  $f_{XY}$ . Let  $Z = X + Y$ .
- ▶ We want to find  $f_Z$  using the theorem.
- ▶ To use the theorem, we need an invertible transformation of  $\mathbb{R}^2$  onto  $\mathbb{R}^2$  of which one component is  $x + y$ .

- ▶ Let  $X, Y$  have joint density  $f_{XY}$ . Let  $Z = X + Y$ .
- ▶ We want to find  $f_Z$  using the theorem.
- ▶ To use the theorem, we need an invertible transformation of  $\mathbb{R}^2$  onto  $\mathbb{R}^2$  of which one component is  $x + y$ .
- ▶ Take  $Z = X + Y$  and  $W = X - Y$ . This is invertible.

- ▶ Let  $X, Y$  have joint density  $f_{XY}$ . Let  $Z = X + Y$ .
- ▶ We want to find  $f_Z$  using the theorem.
- ▶ To use the theorem, we need an invertible transformation of  $\mathbb{R}^2$  onto  $\mathbb{R}^2$  of which one component is  $x + y$ .
- ▶ Take  $Z = X + Y$  and  $W = X - Y$ . This is invertible.
- ▶  $X = (Z + W)/2$  and  $Y = (Z - W)/2$ .

- ▶ Let  $X, Y$  have joint density  $f_{XY}$ . Let  $Z = X + Y$ .
- ▶ We want to find  $f_Z$  using the theorem.
- ▶ To use the theorem, we need an invertible transformation of  $\mathbb{R}^2$  onto  $\mathbb{R}^2$  of which one component is  $x + y$ .
- ▶ Take  $Z = X + Y$  and  $W = X - Y$ . This is invertible.
- ▶  $X = (Z + W)/2$  and  $Y = (Z - W)/2$ . The Jacobian is

$$J = \begin{vmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{vmatrix} = -\frac{1}{2}$$

- ▶ Let  $X, Y$  have joint density  $f_{XY}$ . Let  $Z = X + Y$ .
- ▶ We want to find  $f_Z$  using the theorem.
- ▶ To use the theorem, we need an invertible transformation of  $\mathbb{R}^2$  onto  $\mathbb{R}^2$  of which one component is  $x + y$ .
- ▶ Take  $Z = X + Y$  and  $W = X - Y$ . This is invertible.
- ▶  $X = (Z + W)/2$  and  $Y = (Z - W)/2$ . The Jacobian is

$$J = \begin{vmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{vmatrix} = -\frac{1}{2}$$

- ▶ Hence we get

$$f_{ZW}(z, w) = \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right)$$

- ▶ Let  $X, Y$  have joint density  $f_{XY}$ . Let  $Z = X + Y$ .
- ▶ We want to find  $f_Z$  using the theorem.
- ▶ To use the theorem, we need an invertible transformation of  $\mathbb{R}^2$  onto  $\mathbb{R}^2$  of which one component is  $x + y$ .
- ▶ Take  $Z = X + Y$  and  $W = X - Y$ . This is invertible.
- ▶  $X = (Z + W)/2$  and  $Y = (Z - W)/2$ . The Jacobian is

$$J = \begin{vmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{vmatrix} = -\frac{1}{2}$$

- ▶ Hence we get

$$f_{ZW}(z, w) = \frac{1}{2} f_{XY} \left( \frac{z + w}{2}, \frac{z - w}{2} \right)$$

- ▶ Now we get density of  $Z$  as

$$f_Z(z) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z + w}{2}, \frac{z - w}{2} \right) dw$$

► let  $Z = X + Y$  and  $W = X - Y$ . Then

► let  $Z = X + Y$  and  $W = X - Y$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dw$$



► let  $Z = X + Y$  and  $W = X - Y$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dw$$

change the variable:  $t = \frac{z+w}{2}$

► let  $Z = X + Y$  and  $W = X - Y$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dw$$

$$\text{change the variable: } t = \frac{z+w}{2} \Rightarrow dt = \frac{1}{2} dw$$

► let  $Z = X + Y$  and  $W = X - Y$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dw$$

$$\begin{aligned} \text{change the variable: } t &= \frac{z+w}{2} \Rightarrow dt = \frac{1}{2} dw \\ \Rightarrow w &= 2t - z \Rightarrow z - w = 2z - 2t \end{aligned}$$

► let  $Z = X + Y$  and  $W = X - Y$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dw$$

$$\begin{aligned} \text{change the variable: } t &= \frac{z+w}{2} \Rightarrow dt = \frac{1}{2} dw \\ &\Rightarrow w = 2t - z \Rightarrow z - w = 2z - 2t \end{aligned}$$

$$f_Z(z) = \int_{-\infty}^{\infty} f_{XY}(t, z - t) dt$$

► let  $Z = X + Y$  and  $W = X - Y$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dw$$

$$\begin{aligned} \text{change the variable: } t = \frac{z+w}{2} &\Rightarrow dt = \frac{1}{2} dw \\ &\Rightarrow w = 2t - z \Rightarrow z - w = 2z - 2t \end{aligned}$$

$$\begin{aligned} f_Z(z) &= \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt \\ &= \int_{-\infty}^{\infty} f_{XY}(z-s, s) ds, \end{aligned}$$

- let  $Z = X + Y$  and  $W = X - Y$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dw$$

$$\begin{aligned} \text{change the variable: } t = \frac{z+w}{2} &\Rightarrow dt = \frac{1}{2} dw \\ &\Rightarrow w = 2t - z \Rightarrow z - w = 2z - 2t \end{aligned}$$

$$\begin{aligned} f_Z(z) &= \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt \\ &= \int_{-\infty}^{\infty} f_{XY}(z-s, s) ds, \end{aligned}$$

- We get same result as earlier.

- let  $Z = X + Y$  and  $W = X - Y$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dw$$

$$\begin{aligned} \text{change the variable: } t &= \frac{z+w}{2} \Rightarrow dt = \frac{1}{2} dw \\ \Rightarrow w &= 2t - z \Rightarrow z - w = 2z - 2t \end{aligned}$$

$$\begin{aligned} f_Z(z) &= \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt \\ &= \int_{-\infty}^{\infty} f_{XY}(z-s, s) ds, \end{aligned}$$

- We get same result as earlier. If,  $X, Y$  are independent

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(t) f_Y(z-t) dt$$

► let  $Z = X + Y$  and  $W = X - Y$ . We got



► let  $Z = X + Y$  and  $W = X - Y$ . We got

$$f_{ZW}(z, w) = \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right)$$

- ▶ let  $Z = X + Y$  and  $W = X - Y$ . We got

$$f_{ZW}(z, w) = \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right)$$

- ▶ Now we can calculate  $f_W$  also.

- let  $Z = X + Y$  and  $W = X - Y$ . We got

$$f_{ZW}(z, w) = \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right)$$

- Now we can calculate  $f_W$  also.

$$f_W(w) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dz$$

- let  $Z = X + Y$  and  $W = X - Y$ . We got

$$f_{ZW}(z, w) = \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right)$$

- Now we can calculate  $f_W$  also.

$$f_W(w) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dz$$

change the variable:  $t = \frac{z+w}{2}$

- let  $Z = X + Y$  and  $W = X - Y$ . We got

$$f_{ZW}(z, w) = \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right)$$

- Now we can calculate  $f_W$  also.

$$f_W(w) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dz$$

$$\text{change the variable: } t = \frac{z+w}{2} \Rightarrow dt = \frac{1}{2} dz$$

- let  $Z = X + Y$  and  $W = X - Y$ . We got

$$f_{ZW}(z, w) = \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right)$$

- Now we can calculate  $f_W$  also.

$$f_W(w) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dz$$

$$\begin{aligned} \text{change the variable: } t = \frac{z+w}{2} &\Rightarrow dt = \frac{1}{2} dz \\ \Rightarrow z = 2t - w &\Rightarrow z - w = 2t - 2w \end{aligned}$$

- let  $Z = X + Y$  and  $W = X - Y$ . We got

$$f_{ZW}(z, w) = \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right)$$

- Now we can calculate  $f_W$  also.

$$f_W(w) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dz$$

$$\begin{aligned} \text{change the variable: } t = \frac{z+w}{2} &\Rightarrow dt = \frac{1}{2} dz \\ \Rightarrow z = 2t - w &\Rightarrow z - w = 2t - 2w \end{aligned}$$

$$f_W(w) = \int_{-\infty}^{\infty} f_{XY}(t, t - w) dt$$

- let  $Z = X + Y$  and  $W = X - Y$ . We got

$$f_{ZW}(z, w) = \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right)$$

- Now we can calculate  $f_W$  also.

$$f_W(w) = \int_{-\infty}^{\infty} \frac{1}{2} f_{XY} \left( \frac{z+w}{2}, \frac{z-w}{2} \right) dz$$

$$\begin{aligned} \text{change the variable: } t = \frac{z+w}{2} &\Rightarrow dt = \frac{1}{2} dz \\ \Rightarrow z = 2t - w &\Rightarrow z - w = 2t - 2w \end{aligned}$$

$$\begin{aligned} f_W(w) &= \int_{-\infty}^{\infty} f_{XY}(t, t-w) dt \\ &= \int_{-\infty}^{\infty} f_{XY}(s+w, s) ds, \end{aligned}$$



## Example

- ▶ Let  $X, Y$  be iid  $U[0, 1]$ . Let  $Z = X - Y$ .

## Example

- Let  $X, Y$  be iid  $U[0, 1]$ . Let  $Z = X - Y$ .

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(t) f_Y(t - z) dt$$

## Example

- ▶ Let  $X, Y$  be iid  $U[0, 1]$ . Let  $Z = X - Y$ .

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(t) f_Y(t - z) dt$$

- ▶ For the integrand to be non-zero

## Example

- ▶ Let  $X, Y$  be iid  $U[0, 1]$ . Let  $Z = X - Y$ .

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(t) f_Y(t - z) dt$$

- ▶ For the integrand to be non-zero
  - ▶  $0 \leq t \leq 1 \Rightarrow t \geq 0, t \leq 1$

## Example

- ▶ Let  $X, Y$  be iid  $U[0, 1]$ . Let  $Z = X - Y$ .

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(t) f_Y(t - z) dt$$

- ▶ For the integrand to be non-zero
  - ▶  $0 \leq t \leq 1 \Rightarrow t \geq 0, t \leq 1$
  - ▶  $0 \leq t - z \leq 1 \Rightarrow t \geq z, t \leq 1 + z$

# Example

- ▶ Let  $X, Y$  be iid  $U[0, 1]$ . Let  $Z = X - Y$ .

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(t) f_Y(t - z) dt$$

- ▶ For the integrand to be non-zero
  - ▶  $0 \leq t \leq 1 \Rightarrow t \geq 0, t \leq 1$
  - ▶  $0 \leq t - z \leq 1 \Rightarrow t \geq z, t \leq 1 + z$
  - ▶  $\Rightarrow \max(0, z) \leq t \leq \min(1, 1 + z)$

## Example

- ▶ Let  $X, Y$  be iid  $U[0, 1]$ . Let  $Z = X - Y$ .

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(t) f_Y(t - z) dt$$

- ▶ For the integrand to be non-zero
  - ▶  $0 \leq t \leq 1 \Rightarrow t \geq 0, t \leq 1$
  - ▶  $0 \leq t - z \leq 1 \Rightarrow t \geq z, t \leq 1 + z$
  - ▶  $\Rightarrow \max(0, z) \leq t \leq \min(1, 1 + z)$
- ▶ Thus, we get density as (note  $Z \in (-1, 1)$ )

$$f_Z(z) = \begin{cases} \int_0^{1+z} 1 dt = 1 + z, & \text{if } -1 \leq z \leq 0 \\ \int_z^1 1 dt = 1 - z, & \text{if } 0 \leq z \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

## Example

- ▶ Let  $X, Y$  be iid  $U[0, 1]$ . Let  $Z = X - Y$ .

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(t) f_Y(t - z) dt$$

- ▶ For the integrand to be non-zero
  - ▶  $0 \leq t \leq 1 \Rightarrow t \geq 0, t \leq 1$
  - ▶  $0 \leq t - z \leq 1 \Rightarrow t \geq z, t \leq 1 + z$
  - ▶  $\Rightarrow \max(0, z) \leq t \leq \min(1, 1 + z)$
- ▶ Thus, we get density as (note  $Z \in (-1, 1)$ )

$$f_Z(z) = \begin{cases} \int_0^{1+z} 1 dt = 1 + z, & \text{if } -1 \leq z \leq 0 \\ \int_z^1 1 dt = 1 - z, & 0 \leq z \leq 1 \end{cases}$$



## Example

- ▶ Let  $X, Y$  be iid  $U[0, 1]$ . Let  $Z = X - Y$ .

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(t) f_Y(t - z) dt$$

- ▶ For the integrand to be non-zero
  - ▶  $0 \leq t \leq 1 \Rightarrow t \geq 0, t \leq 1$
  - ▶  $0 \leq t - z \leq 1 \Rightarrow t \geq z, t \leq 1 + z$
  - ▶  $\Rightarrow \max(0, z) \leq t \leq \min(1, 1 + z)$
- ▶ Thus, we get density as (note  $Z \in (-1, 1)$ )

$$f_Z(z) = \begin{cases} \int_0^{1+z} 1 dt = 1 + z, & \text{if } -1 \leq z \leq 0 \\ \int_z^1 1 dt = 1 - z, & 0 \leq z \leq 1 \end{cases}$$

- ▶ Thus, when  $X, Y \sim U(0, 1)$  iid

$$f_{X-Y}(z) = 1 - |z|, \quad -1 < z < 1$$

► We showed that

► We showed that

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt = \int_{-\infty}^{\infty} f_{XY}(z-t, t) dt$$

► We showed that

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt = \int_{-\infty}^{\infty} f_{XY}(z-t, t) dt$$

$$f_{X-Y}(w) = \int_{-\infty}^{\infty} f_{XY}(t, t-w) dt = \int_{-\infty}^{\infty} f_{XY}(t+w, t) dt$$

- We showed that

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt = \int_{-\infty}^{\infty} f_{XY}(z-t, t) dt$$

$$f_{X-Y}(w) = \int_{-\infty}^{\infty} f_{XY}(t, t-w) dt = \int_{-\infty}^{\infty} f_{XY}(t+w, t) dt$$

- Suppose  $X, Y$  are discrete. Then we have

- We showed that

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt = \int_{-\infty}^{\infty} f_{XY}(z-t, t) dt$$

$$f_{X-Y}(w) = \int_{-\infty}^{\infty} f_{XY}(t, t-w) dt = \int_{-\infty}^{\infty} f_{XY}(t+w, t) dt$$

- Suppose  $X, Y$  are discrete. Then we have

$$f_{X+Y}(z) = P[X+Y=z] = \sum_k P[X=k, Y=z-k]$$

- We showed that

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt = \int_{-\infty}^{\infty} f_{XY}(z-t, t) dt$$

$$f_{X-Y}(w) = \int_{-\infty}^{\infty} f_{XY}(t, t-w) dt = \int_{-\infty}^{\infty} f_{XY}(t+w, t) dt$$

- Suppose  $X, Y$  are discrete. Then we have

$$\begin{aligned} f_{X+Y}(z) &= P[X+Y=z] = \sum_k P[X=k, Y=z-k] \\ &= \sum_k f_{XY}(k, z-k) \end{aligned}$$

- We showed that

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt = \int_{-\infty}^{\infty} f_{XY}(z-t, t) dt$$

$$f_{X-Y}(w) = \int_{-\infty}^{\infty} f_{XY}(t, t-w) dt = \int_{-\infty}^{\infty} f_{XY}(t+w, t) dt$$

- Suppose  $X, Y$  are discrete. Then we have

$$f_{X+Y}(z) = P[X+Y=z] = \sum_k P[X=k, Y=z-k]$$

$$= \sum_k f_{XY}(k, z-k)$$

$$f_{X-Y}(w) = P[X-Y=w] = \sum_k P[X=k, Y=k-w]$$



- We showed that

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_{XY}(t, z-t) dt = \int_{-\infty}^{\infty} f_{XY}(z-t, t) dt$$

$$f_{X-Y}(w) = \int_{-\infty}^{\infty} f_{XY}(t, t-w) dt = \int_{-\infty}^{\infty} f_{XY}(t+w, t) dt$$

- Suppose  $X, Y$  are discrete. Then we have

$$f_{X+Y}(z) = P[X+Y=z] = \sum_k P[X=k, Y=z-k]$$

$$= \sum_k f_{XY}(k, z-k)$$

$$f_{X-Y}(w) = P[X-Y=w] = \sum_k P[X=k, Y=k-w]$$

$$= \sum_k f_{XY}(k, k-w)$$

# Distribution of product of random variables

- ▶ We want density of  $Z = XY$ .

# Distribution of product of random variables

- ▶ We want density of  $Z = XY$ .
- ▶ We need one more function to make an invertible transformation

# Distribution of product of random variables

- ▶ We want density of  $Z = XY$ .
- ▶ We need one more function to make an invertible transformation
- ▶ A possible choice:  $Z = XY$      $W = Y$

# Distribution of product of random variables

- ▶ We want density of  $Z = XY$ .
- ▶ We need one more function to make an invertible transformation
- ▶ A possible choice:  $Z = XY \quad W = Y$
- ▶ This is invertible:  $X = Z/W \quad Y = W$

# Distribution of product of random variables

- ▶ We want density of  $Z = XY$ .
- ▶ We need one more function to make an invertible transformation
- ▶ A possible choice:  $Z = XY \quad W = Y$
- ▶ This is invertible:  $X = Z/W \quad Y = W$

$$J = \begin{vmatrix} \frac{1}{w} & \frac{-z}{w^2} \\ 0 & 1 \end{vmatrix} = \frac{1}{w}$$

# Distribution of product of random variables

- ▶ We want density of  $Z = XY$ .
- ▶ We need one more function to make an invertible transformation
- ▶ A possible choice:  $Z = XY$     $W = Y$
- ▶ This is invertible:  $X = Z/W$     $Y = W$

$$J = \begin{vmatrix} \frac{1}{w} & \frac{-z}{w^2} \\ 0 & 1 \end{vmatrix} = \frac{1}{w}$$

- ▶ Hence we get

$$f_{ZW}(z, w) = \left| \frac{1}{w} \right| f_{XY}\left(\frac{z}{w}, w\right)$$

# Distribution of product of random variables

- ▶ We want density of  $Z = XY$ .
- ▶ We need one more function to make an invertible transformation
- ▶ A possible choice:  $Z = XY \quad W = Y$
- ▶ This is invertible:  $X = Z/W \quad Y = W$

$$J = \begin{vmatrix} \frac{1}{w} & \frac{-z}{w^2} \\ 0 & 1 \end{vmatrix} = \frac{1}{w}$$

- ▶ Hence we get

$$f_{ZW}(z, w) = \left| \frac{1}{w} \right| f_{XY} \left( \frac{z}{w}, w \right)$$

- ▶ Thus we get the density of product as

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{1}{w} \right| f_{XY} \left( \frac{z}{w}, w \right) dw$$



## example

- ▶ Let  $X, Y$  be iid  $U(0, 1)$ . Let  $Z = XY$ .

## example

- Let  $X, Y$  be iid  $U(0, 1)$ . Let  $Z = XY$ .

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{1}{w} \right| f_X\left(\frac{z}{w}\right) f_Y(w) dw$$

## example

- ▶ Let  $X, Y$  be iid  $U(0, 1)$ . Let  $Z = XY$ .

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{1}{w} \right| f_X\left(\frac{z}{w}\right) f_Y(w) dw$$

- ▶ We need:  $0 < w < 1$  and  $0 < \frac{z}{w} < 1$ . Hence

## example

- ▶ Let  $X, Y$  be iid  $U(0, 1)$ . Let  $Z = XY$ .

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{1}{w} \right| f_X\left(\frac{z}{w}\right) f_Y(w) dw$$

- ▶ We need:  $0 < w < 1$  and  $0 < \frac{z}{w} < 1$ . Hence

$$f_Z(z) = \int_z^1 \left| \frac{1}{w} \right| dw$$

## example

- ▶ Let  $X, Y$  be iid  $U(0, 1)$ . Let  $Z = XY$ .

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{1}{w} \right| f_X\left(\frac{z}{w}\right) f_Y(w) dw$$

- ▶ We need:  $0 < w < 1$  and  $0 < \frac{z}{w} < 1$ . Hence

$$f_Z(z) = \int_z^1 \left| \frac{1}{w} \right| dw = \int_z^1 \frac{1}{w} dw = -\ln(z), \quad 0 < z < 1$$

- ▶  $X, Y$  have joint density and  $Z = XY$ . Then

- $X, Y$  have joint density and  $Z = XY$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{1}{w} \right| f_{XY} \left( \frac{z}{w}, w \right) dw$$

- $X, Y$  have joint density and  $Z = XY$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{1}{w} \right| f_{XY} \left( \frac{z}{w}, w \right) dw$$

Suppose  $X, Y$  are discrete and  $Z = XY$



- $X, Y$  have joint density and  $Z = XY$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{1}{w} \right| f_{XY} \left( \frac{z}{w}, w \right) dw$$

Suppose  $X, Y$  are discrete and  $Z = XY$

$$f_Z(0) = P[X = 0 \text{ or } Y = 0] = \sum_x f_{XY}(x, 0) + \sum_y f_{XY}(0, y)$$

- $X, Y$  have joint density and  $Z = XY$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{1}{w} \right| f_{XY} \left( \frac{z}{w}, w \right) dw$$

Suppose  $X, Y$  are discrete and  $Z = XY$

$$f_Z(0) = P[X = 0 \text{ or } Y = 0] = \sum_x f_{XY}(x, 0) + \sum_y f_{XY}(0, y)$$

$$f_Z(k) = \sum_{y \neq 0} P \left[ X = \frac{k}{y}, Y = y \right] = \sum_{y \neq 0} f_{XY} \left( \frac{k}{y}, y \right), \quad k \neq 0$$

- $X, Y$  have joint density and  $Z = XY$ . Then

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{1}{w} \right| f_{XY} \left( \frac{z}{w}, w \right) dw$$

Suppose  $X, Y$  are discrete and  $Z = XY$

$$f_Z(0) = P[X = 0 \text{ or } Y = 0] = \sum_x f_{XY}(x, 0) + \sum_y f_{XY}(0, y)$$

$$f_Z(k) = \sum_{y \neq 0} P \left[ X = \frac{k}{y}, Y = y \right] = \sum_{y \neq 0} f_{XY} \left( \frac{k}{y}, y \right), \quad k \neq 0$$

- We cannot always interchange density and mass functions!!

- ▶ We wanted density of  $Z = XY$ .

- ▶ We wanted density of  $Z = XY$ .
- ▶ We used:  $Z = XY$  and  $W = Y$ .

- ▶ We wanted density of  $Z = XY$ .
- ▶ We used:  $Z = XY$  and  $W = Y$ .
- ▶ We could have used:  $Z = XY$  and  $W = X$ .

- ▶ We wanted density of  $Z = XY$ .
- ▶ We used:  $Z = XY$  and  $W = Y$ .
- ▶ We could have used:  $Z = XY$  and  $W = X$ .
- ▶ This is invertible:  $X = W$  and  $Y = Z/W$ .

- ▶ We wanted density of  $Z = XY$ .
- ▶ We used:  $Z = XY$  and  $W = Y$ .
- ▶ We could have used:  $Z = XY$  and  $W = X$ .
- ▶ This is invertible:  $X = W$  and  $Y = Z/W$ .

$$J = \begin{vmatrix} 0 & 1 \\ \frac{1}{w} & \frac{-z}{w^2} \end{vmatrix} = -\frac{1}{w}$$



- ▶ We wanted density of  $Z = XY$ .
- ▶ We used:  $Z = XY$  and  $W = Y$ .
- ▶ We could have used:  $Z = XY$  and  $W = X$ .
- ▶ This is invertible:  $X = W$  and  $Y = Z/W$ .

$$J = \begin{vmatrix} 0 & 1 \\ \frac{1}{w} & \frac{-z}{w^2} \end{vmatrix} = -\frac{1}{w}$$

- ▶ This gives

- ▶ We wanted density of  $Z = XY$ .
- ▶ We used:  $Z = XY$  and  $W = Y$ .
- ▶ We could have used:  $Z = XY$  and  $W = X$ .
- ▶ This is invertible:  $X = W$  and  $Y = Z/W$ .

$$J = \begin{vmatrix} 0 & 1 \\ \frac{1}{w} & \frac{-z}{w^2} \end{vmatrix} = -\frac{1}{w}$$

- ▶ This gives

$$f_{ZW}(z, w) = \left| \frac{1}{w} \right| f_{XY} \left( w, \frac{z}{w} \right)$$

- ▶ We wanted density of  $Z = XY$ .
- ▶ We used:  $Z = XY$  and  $W = Y$ .
- ▶ We could have used:  $Z = XY$  and  $W = X$ .
- ▶ This is invertible:  $X = W$  and  $Y = Z/W$ .

$$J = \begin{vmatrix} 0 & 1 \\ \frac{1}{w} & \frac{-z}{w^2} \end{vmatrix} = -\frac{1}{w}$$

- ▶ This gives

$$\begin{aligned} f_{ZW}(z, w) &= \left| \frac{1}{w} \right| f_{XY} \left( w, \frac{z}{w} \right) \\ f_Z(z) &= \int_{-\infty}^{\infty} \left| \frac{1}{w} \right| f_{XY} \left( w, \frac{z}{w} \right) dw \end{aligned}$$

- ▶ We wanted density of  $Z = XY$ .
- ▶ We used:  $Z = XY$  and  $W = Y$ .
- ▶ We could have used:  $Z = XY$  and  $W = X$ .
- ▶ This is invertible:  $X = W$  and  $Y = Z/W$ .

$$J = \begin{vmatrix} 0 & 1 \\ \frac{1}{w} & \frac{-z}{w^2} \end{vmatrix} = -\frac{1}{w}$$

- ▶ This gives

$$f_{ZW}(z, w) = \left| \frac{1}{w} \right| f_{XY} \left( w, \frac{z}{w} \right)$$

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{1}{w} \right| f_{XY} \left( w, \frac{z}{w} \right) dw$$

- ▶ The  $f_Z$  should be same in both cases.

# Distributions of quotients

- ▶  $X, Y$  have joint density and  $Z = X/Y$ .

# Distributions of quotients

- ▶  $X, Y$  have joint density and  $Z = X/Y$ .
- ▶ We can take:  $Z = X/Y$      $W = Y$

# Distributions of quotients

- ▶  $X, Y$  have joint density and  $Z = X/Y$ .
- ▶ We can take:  $Z = X/Y$      $W = Y$
- ▶ This is invertible:  $X = ZW$      $Y = W$

# Distributions of quotients

- ▶  $X, Y$  have joint density and  $Z = X/Y$ .
- ▶ We can take:  $Z = X/Y$     $W = Y$
- ▶ This is invertible:  $X = ZW$     $Y = W$

$$J = \begin{vmatrix} w & z \\ 0 & 1 \end{vmatrix} = w$$



# Distributions of quotients

- ▶  $X, Y$  have joint density and  $Z = X/Y$ .
- ▶ We can take:  $Z = X/Y$     $W = Y$
- ▶ This is invertible:  $X = ZW$     $Y = W$

$$J = \begin{vmatrix} w & z \\ 0 & 1 \end{vmatrix} = w$$

- ▶ Hence we get

$$f_{ZW}(z, w) = |w| f_{XY}(zw, w)$$

# Distributions of quotients

- ▶  $X, Y$  have joint density and  $Z = X/Y$ .
- ▶ We can take:  $Z = X/Y$     $W = Y$
- ▶ This is invertible:  $X = ZW$     $Y = W$

$$J = \begin{vmatrix} w & z \\ 0 & 1 \end{vmatrix} = w$$

- ▶ Hence we get

$$f_{ZW}(z, w) = |w| f_{XY}(zw, w)$$

- ▶ Thus we get the density of quotient as

$$f_Z(z) = \int_{-\infty}^{\infty} |w| f_{XY}(zw, w) dw$$

## example

- ▶ Let  $X, Y$  be iid  $U(0, 1)$ . Let  $Z = X/Y$ .

## example

- ▶ Let  $X, Y$  be iid  $U(0, 1)$ . Let  $Z = X/Y$ .  
Note  $Z \in (0, \infty)$

## example

- ▶ Let  $X, Y$  be iid  $U(0, 1)$ . Let  $Z = X/Y$ .  
Note  $Z \in (0, \infty)$

$$f_Z(z) = \int_{-\infty}^{\infty} |w| f_X(zw) f_Y(w) dw$$

## example

- ▶ Let  $X, Y$  be iid  $U(0, 1)$ . Let  $Z = X/Y$ .  
Note  $Z \in (0, \infty)$

$$f_Z(z) = \int_{-\infty}^{\infty} |w| f_X(zw) f_Y(w) dw$$

- ▶ We need  $0 < w < 1$  and  $0 < zw < 1 \Rightarrow w < 1/z$ .

## example

- ▶ Let  $X, Y$  be iid  $U(0, 1)$ . Let  $Z = X/Y$ .  
Note  $Z \in (0, \infty)$

$$f_Z(z) = \int_{-\infty}^{\infty} |w| f_X(zw) f_Y(w) dw$$

- ▶ We need  $0 < w < 1$  and  $0 < zw < 1 \Rightarrow w < 1/z$ .
- ▶ So, when  $z \leq 1$ ,  $w$  goes from 0 to 1; when  $z > 1$ ,  $w$  goes from 0 to  $1/z$ .

## example

- ▶ Let  $X, Y$  be iid  $U(0, 1)$ . Let  $Z = X/Y$ .  
Note  $Z \in (0, \infty)$

$$f_Z(z) = \int_{-\infty}^{\infty} |w| f_X(zw) f_Y(w) dw$$

- ▶ We need  $0 < w < 1$  and  $0 < zw < 1 \Rightarrow w < 1/z$ .
- ▶ So, when  $z \leq 1$ ,  $w$  goes from 0 to 1; when  $z > 1$ ,  $w$  goes from 0 to  $1/z$ .
- ▶ Hence we get density as

$$f_Z(z) = \begin{cases} \int_0^1 w dw = \frac{1}{2}, & \text{if } 0 < z \leq 1 \\ \int_0^{1/z} w dw = \frac{1}{2z^2}, & \text{if } z > 1 \end{cases}$$



## example

- ▶ Let  $X, Y$  be iid  $U(0, 1)$ . Let  $Z = X/Y$ .  
Note  $Z \in (0, \infty)$

$$f_Z(z) = \int_{-\infty}^{\infty} |w| f_X(zw) f_Y(w) dw$$

- ▶ We need  $0 < w < 1$  and  $0 < zw < 1 \Rightarrow w < 1/z$ .
- ▶ So, when  $z \leq 1$ ,  $w$  goes from 0 to 1; when  $z > 1$ ,  $w$  goes from 0 to  $1/z$ .
- ▶ Hence we get density as

$$f_Z(z) = \begin{cases} \int_0^1 w dw = \frac{1}{2}, & \text{if } 0 < z \leq 1 \\ \int_0^{1/z} w dw = \frac{1}{2z^2}, & 1 < z < \infty \end{cases}$$

- $X, Y$  have joint density and  $Z = X/Y$

$$f_Z(z) = \int_{-\infty}^{\infty} |w| f_{XY}(zw, w) dw$$

- ▶  $X, Y$  have joint density and  $Z = X/Y$

$$f_Z(z) = \int_{-\infty}^{\infty} |w| f_{XY}(zw, w) dw$$

- ▶ Suppose  $X, Y$  are discrete and  $Z = X/Y$

- ▶  $X, Y$  have joint density and  $Z = X/Y$

$$f_Z(z) = \int_{-\infty}^{\infty} |w| f_{XY}(zw, w) dw$$

- ▶ Suppose  $X, Y$  are discrete and  $Z = X/Y$

$$f_Z(z) = P[Z = z] = P[X/Y = z]$$

- ▶  $X, Y$  have joint density and  $Z = X/Y$

$$f_Z(z) = \int_{-\infty}^{\infty} |w| f_{XY}(zw, w) dw$$

- ▶ Suppose  $X, Y$  are discrete and  $Z = X/Y$

$$\begin{aligned} f_Z(z) &= P[Z = z] = P[X/Y = z] \\ &= \sum_y P[X = yz, Y = y] \end{aligned}$$

- ▶  $X, Y$  have joint density and  $Z = X/Y$

$$f_Z(z) = \int_{-\infty}^{\infty} |w| f_{XY}(zw, w) dw$$

- ▶ Suppose  $X, Y$  are discrete and  $Z = X/Y$

$$\begin{aligned} f_Z(z) &= P[Z = z] = P[X/Y = z] \\ &= \sum_y P[X = yz, Y = y] \\ &= \sum_y f_{XY}(yz, y) \end{aligned}$$

- ▶ We chose:  $Z = X/Y$  and  $W = Y$ .

- ▶ We chose:  $Z = X/Y$  and  $W = Y$ .
- ▶ We could have taken:  $Z = X/Y$  and  $W = X$



- ▶ We chose:  $Z = X/Y$  and  $W = Y$ .
- ▶ We could have taken:  $Z = X/Y$  and  $W = X$
- ▶ The inverse is:  $X = W$  and  $Y = W/Z$

- ▶ We chose:  $Z = X/Y$  and  $W = Y$ .
- ▶ We could have taken:  $Z = X/Y$  and  $W = X$
- ▶ The inverse is:  $X = W$  and  $Y = W/Z$

$$J = \begin{vmatrix} 0 & 1 \\ -\frac{w}{z^2} & \frac{1}{z} \end{vmatrix} = -\frac{w}{z^2}$$

- ▶ We chose:  $Z = X/Y$  and  $W = Y$ .
- ▶ We could have taken:  $Z = X/Y$  and  $W = X$
- ▶ The inverse is:  $X = W$  and  $Y = W/Z$

$$J = \begin{vmatrix} 0 & 1 \\ -\frac{w}{z^2} & \frac{1}{z} \end{vmatrix} = -\frac{w}{z^2}$$

- ▶ Thus we get the density of quotient as

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{w}{z^2} \right| f_{XY} \left( w, \frac{w}{z} \right) dw$$

- ▶ We chose:  $Z = X/Y$  and  $W = Y$ .
- ▶ We could have taken:  $Z = X/Y$  and  $W = X$
- ▶ The inverse is:  $X = W$  and  $Y = W/Z$

$$J = \begin{vmatrix} 0 & 1 \\ -\frac{w}{z^2} & \frac{1}{z} \end{vmatrix} = -\frac{w}{z^2}$$

- ▶ Thus we get the density of quotient as

$$f_Z(z) = \int_{-\infty}^{\infty} \left| \frac{w}{z^2} \right| f_{XY} \left( w, \frac{w}{z} \right) dw$$

put  $t = \frac{w}{z} \Rightarrow dt = \frac{dw}{z}, \quad w = tz$

- ▶ We chose:  $Z = X/Y$  and  $W = Y$ .
- ▶ We could have taken:  $Z = X/Y$  and  $W = X$
- ▶ The inverse is:  $X = W$  and  $Y = W/Z$

$$J = \begin{vmatrix} 0 & 1 \\ -\frac{w}{z^2} & \frac{1}{z} \end{vmatrix} = -\frac{w}{z^2}$$

- ▶ Thus we get the density of quotient as

$$\begin{aligned} f_Z(z) &= \int_{-\infty}^{\infty} \left| \frac{w}{z^2} \right| f_{XY} \left( w, \frac{w}{z} \right) dw \\ &\text{put } t = \frac{w}{z} \Rightarrow dt = \frac{dw}{z}, \quad w = tz \\ &= \int_{-\infty}^{\infty} |t| f_{XY}(tz, t) dt \end{aligned}$$

- ▶ We chose:  $Z = X/Y$  and  $W = Y$ .
- ▶ We could have taken:  $Z = X/Y$  and  $W = X$
- ▶ The inverse is:  $X = W$  and  $Y = W/Z$

$$J = \begin{vmatrix} 0 & 1 \\ -\frac{w}{z^2} & \frac{1}{z} \end{vmatrix} = -\frac{w}{z^2}$$

- ▶ Thus we get the density of quotient as

$$\begin{aligned} f_Z(z) &= \int_{-\infty}^{\infty} \left| \frac{w}{z^2} \right| f_{XY} \left( w, \frac{w}{z} \right) dw \\ &\text{put } t = \frac{w}{z} \Rightarrow dt = \frac{dw}{z}, \quad w = tz \\ &= \int_{-\infty}^{\infty} |t| f_{XY}(tz, t) dt \end{aligned}$$

- ▶ We can show that the density of quotient is same in both these approaches.

# Exchangeable Random Variables

- ▶  $X_1, X_2, \dots, X_n$  are said to be exchangeable if their joint distribution is same as that of any permutation of them.

# Exchangeable Random Variables

- ▶  $X_1, X_2, \dots, X_n$  are said to be exchangeable if their joint distribution is same as that of any permutation of them.
- ▶ let  $(i_1, \dots, i_n)$  be a permutation of  $(1, 2, \dots, n)$ . Then joint df of  $(X_{i_1}, \dots, X_{i_n})$  should be same as that  $(X_1, \dots, X_n)$



# Exchangeable Random Variables

- ▶  $X_1, X_2, \dots, X_n$  are said to be exchangeable if their joint distribution is same as that of any permutation of them.
- ▶ let  $(i_1, \dots, i_n)$  be a permutation of  $(1, 2, \dots, n)$ . Then joint df of  $(X_{i_1}, \dots, X_{i_n})$  should be same as that  $(X_1, \dots, X_n)$
- ▶ Take  $n = 3$ . Suppose  $F_{X_1 X_2 X_3}(a, b, c) = g(a, b, c)$ . If they are exchangeable, then

# Exchangeable Random Variables

- ▶  $X_1, X_2, \dots, X_n$  are said to be exchangeable if their joint distribution is same as that of any permutation of them.
- ▶ let  $(i_1, \dots, i_n)$  be a permutation of  $(1, 2, \dots, n)$ . Then joint df of  $(X_{i_1}, \dots, X_{i_n})$  should be same as that  $(X_1, \dots, X_n)$
- ▶ Take  $n = 3$ . Suppose  $F_{X_1 X_2 X_3}(a, b, c) = g(a, b, c)$ . If they are exchangeable, then

$$F_{X_2 X_3 X_1}(a, b, c) = P[X_2 \leq a, X_3 \leq b, X_1 \leq c]$$

# Exchangeable Random Variables

- ▶  $X_1, X_2, \dots, X_n$  are said to be exchangeable if their joint distribution is same as that of any permutation of them.
- ▶ let  $(i_1, \dots, i_n)$  be a permutation of  $(1, 2, \dots, n)$ . Then joint df of  $(X_{i_1}, \dots, X_{i_n})$  should be same as that  $(X_1, \dots, X_n)$
- ▶ Take  $n = 3$ . Suppose  $F_{X_1 X_2 X_3}(a, b, c) = g(a, b, c)$ . If they are exchangeable, then

$$\begin{aligned} F_{X_2 X_3 X_1}(a, b, c) &= P[X_2 \leq a, X_3 \leq b, X_1 \leq c] \\ &= P[X_1 \leq c, X_2 \leq a, X_3 \leq b] \end{aligned}$$

# Exchangeable Random Variables

- ▶  $X_1, X_2, \dots, X_n$  are said to be exchangeable if their joint distribution is same as that of any permutation of them.
- ▶ let  $(i_1, \dots, i_n)$  be a permutation of  $(1, 2, \dots, n)$ . Then joint df of  $(X_{i_1}, \dots, X_{i_n})$  should be same as that  $(X_1, \dots, X_n)$
- ▶ Take  $n = 3$ . Suppose  $F_{X_1 X_2 X_3}(a, b, c) = g(a, b, c)$ . If they are exchangeable, then

$$\begin{aligned} F_{X_2 X_3 X_1}(a, b, c) &= P[X_2 \leq a, X_3 \leq b, X_1 \leq c] \\ &= P[X_1 \leq c, X_2 \leq a, X_3 \leq b] \\ &= g(c, a, b) \end{aligned}$$

# Exchangeable Random Variables

- ▶  $X_1, X_2, \dots, X_n$  are said to be exchangeable if their joint distribution is same as that of any permutation of them.
- ▶ let  $(i_1, \dots, i_n)$  be a permutation of  $(1, 2, \dots, n)$ . Then joint df of  $(X_{i_1}, \dots, X_{i_n})$  should be same as that  $(X_1, \dots, X_n)$
- ▶ Take  $n = 3$ . Suppose  $F_{X_1 X_2 X_3}(a, b, c) = g(a, b, c)$ . If they are exchangeable, then

$$\begin{aligned} F_{X_2 X_3 X_1}(a, b, c) &= P[X_2 \leq a, X_3 \leq b, X_1 \leq c] \\ &= P[X_1 \leq c, X_2 \leq a, X_3 \leq b] \\ &= g(c, a, b) = g(a, b, c) \end{aligned}$$

# Exchangeable Random Variables

- ▶  $X_1, X_2, \dots, X_n$  are said to be exchangeable if their joint distribution is same as that of any permutation of them.
- ▶ let  $(i_1, \dots, i_n)$  be a permutation of  $(1, 2, \dots, n)$ . Then joint df of  $(X_{i_1}, \dots, X_{i_n})$  should be same as that  $(X_1, \dots, X_n)$
- ▶ Take  $n = 3$ . Suppose  $F_{X_1 X_2 X_3}(a, b, c) = g(a, b, c)$ . If they are exchangeable, then

$$\begin{aligned} F_{X_2 X_3 X_1}(a, b, c) &= P[X_2 \leq a, X_3 \leq b, X_1 \leq c] \\ &= P[X_1 \leq c, X_2 \leq a, X_3 \leq b] \\ &= g(c, a, b) = g(a, b, c) \end{aligned}$$

- ▶ The df or density should be “symmetric” in its variables if the random variables are exchangeable.

- Consider the density of three random variables

$$f(x, y, z) = \frac{2}{3}(x + y + z), \quad 0 < x, y, z < 1$$

- ▶ Consider the density of three random variables

$$f(x, y, z) = \frac{2}{3}(x + y + z), \quad 0 < x, y, z < 1$$

- ▶ They are exchangeable (because  $f(x, y, z) = f(y, x, z)$ )



- ▶ Consider the density of three random variables

$$f(x, y, z) = \frac{2}{3}(x + y + z), \quad 0 < x, y, z < 1$$

- ▶ They are exchangeable (because  $f(x, y, z) = f(y, x, z)$ )
- ▶ If random variables are exchangeable then they are identically distributed.

- ▶ Consider the density of three random variables

$$f(x, y, z) = \frac{2}{3}(x + y + z), \quad 0 < x, y, z < 1$$

- ▶ They are exchangeable (because  $f(x, y, z) = f(y, x, z)$ )
- ▶ If random variables are exchangeable then they are identically distributed.

$$F_{XYZ}(a, \infty, \infty) = F_{XYZ}(\infty, \infty, a) \Rightarrow F_X(a) = F_Z(a)$$

- ▶ Consider the density of three random variables

$$f(x, y, z) = \frac{2}{3}(x + y + z), \quad 0 < x, y, z < 1$$

- ▶ They are exchangeable (because  $f(x, y, z) = f(y, x, z)$ )
- ▶ If random variables are exchangeable then they are identically distributed.

$$F_{XYZ}(a, \infty, \infty) = F_{XYZ}(\infty, \infty, a) \Rightarrow F_X(a) = F_Z(a)$$

- ▶ The above example shows that exchangeable random variables need not be independent. The joint density is not factorizable.

- ▶ Consider the density of three random variables

$$f(x, y, z) = \frac{2}{3}(x + y + z), \quad 0 < x, y, z < 1$$

- ▶ They are exchangeable (because  $f(x, y, z) = f(y, x, z)$ )
- ▶ If random variables are exchangeable then they are identically distributed.

$$F_{XYZ}(a, \infty, \infty) = F_{XYZ}(\infty, \infty, a) \Rightarrow F_X(a) = F_Z(a)$$

- ▶ The above example shows that exchangeable random variables need not be independent. The joint density is not factorizable.

$$\int_0^1 \int_0^1 \frac{2}{3}(x + y + z) \, dy \, dz = \frac{2(x+1)}{3}$$

- Consider the density of three random variables

$$f(x, y, z) = \frac{2}{3}(x + y + z), \quad 0 < x, y, z < 1$$

- They are exchangeable (because  $f(x, y, z) = f(y, x, z)$ )
- If random variables are exchangeable then they are identically distributed.

$$F_{XYZ}(a, \infty, \infty) = F_{XYZ}(\infty, \infty, a) \Rightarrow F_X(a) = F_Z(a)$$

- The above example shows that exchangeable random variables need not be independent. The joint density is not factorizable.

$$\int_0^1 \int_0^1 \frac{2}{3}(x + y + z) dy dz = \frac{2(x+1)}{3}$$

- So, the joint density is not the product of marginals

# Expectation of functions of multiple rv

# Expectation of functions of multiple rv

► **Theorem:** Let  $Z = g(X_1, \dots, X_n) = g(\mathbf{X})$ . Then

$$E[Z] = \int_{\mathbb{R}^n} g(\mathbf{x}) dF_{\mathbf{X}}(\mathbf{x})$$

# Expectation of functions of multiple rv

- **Theorem:** Let  $Z = g(X_1, \dots, X_n) = g(\mathbf{X})$ . Then

$$E[Z] = \int_{\mathbb{R}^n} g(\mathbf{x}) dF_{\mathbf{X}}(\mathbf{x})$$

- That is, if they have a joint density, then

$$E[Z] = \int_{\mathbb{R}^n} g(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\mathbf{x}$$



# Expectation of functions of multiple rv

- **Theorem:** Let  $Z = g(X_1, \dots, X_n) = g(\mathbf{X})$ . Then

$$E[Z] = \int_{\mathbb{R}^n} g(\mathbf{x}) dF_{\mathbf{X}}(\mathbf{x})$$

- That is, if they have a joint density, then

$$E[Z] = \int_{\mathbb{R}^n} g(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x}) d\mathbf{x}$$

- Similarly, if all  $X_i$  are discrete

$$E[Z] = \sum_{\mathbf{x}} g(\mathbf{x}) f_{\mathbf{X}}(\mathbf{x})$$

- ▶ Let  $Z = X + Y$ . Let  $X, Y$  have joint density  $f_{XY}$

- Let  $Z = X + Y$ . Let  $X, Y$  have joint density  $f_{XY}$

$$E[X + Y] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x + y) f_{XY}(x, y) dx dy$$

- Let  $Z = X + Y$ . Let  $X, Y$  have joint density  $f_{XY}$

$$\begin{aligned} E[X + Y] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x + y) f_{XY}(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x \int_{-\infty}^{\infty} f_{XY}(x, y) dy dx \\ &\quad + \int_{-\infty}^{\infty} y \int_{-\infty}^{\infty} f_{XY}(x, y) dx dy \end{aligned}$$

- Let  $Z = X + Y$ . Let  $X, Y$  have joint density  $f_{XY}$

$$\begin{aligned} E[X + Y] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x + y) f_{XY}(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x \int_{-\infty}^{\infty} f_{XY}(x, y) dy dx \\ &\quad + \int_{-\infty}^{\infty} y \int_{-\infty}^{\infty} f_{XY}(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x f_X(x) dx + \int_{-\infty}^{\infty} y f_Y(y) dy \end{aligned}$$

► Let  $Z = X + Y$ . Let  $X, Y$  have joint density  $f_{XY}$

$$\begin{aligned} E[X + Y] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x + y) f_{XY}(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x \int_{-\infty}^{\infty} f_{XY}(x, y) dy dx \\ &\quad + \int_{-\infty}^{\infty} y \int_{-\infty}^{\infty} f_{XY}(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x f_X(x) dx + \int_{-\infty}^{\infty} y f_Y(y) dy \\ &= E[X] + E[Y] \end{aligned}$$

- Let  $Z = X + Y$ . Let  $X, Y$  have joint density  $f_{XY}$

$$\begin{aligned} E[X + Y] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x + y) f_{XY}(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x \int_{-\infty}^{\infty} f_{XY}(x, y) dy dx \\ &\quad + \int_{-\infty}^{\infty} y \int_{-\infty}^{\infty} f_{XY}(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x f_X(x) dx + \int_{-\infty}^{\infty} y f_Y(y) dy \\ &= E[X] + E[Y] \end{aligned}$$

- Expectation is a linear operator.

- ▶ Let  $Z = X + Y$ . Let  $X, Y$  have joint density  $f_{XY}$

$$\begin{aligned} E[X + Y] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x + y) f_{XY}(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x \int_{-\infty}^{\infty} f_{XY}(x, y) dy dx \\ &\quad + \int_{-\infty}^{\infty} y \int_{-\infty}^{\infty} f_{XY}(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x f_X(x) dx + \int_{-\infty}^{\infty} y f_Y(y) dy \\ &= E[X] + E[Y] \end{aligned}$$

- ▶ Expectation is a linear operator.
- ▶ This is true for all random variables.



► We saw  $E[X + Y] = E[X] + E[Y]$ .

- ▶ We saw  $E[X + Y] = E[X] + E[Y]$ .
- ▶ Let us calculate  $\text{Var}(X + Y)$ .

- ▶ We saw  $E[X + Y] = E[X] + E[Y]$ .
- ▶ Let us calculate  $\text{Var}(X + Y)$ .

$$\text{Var}(X + Y) = E \left[ ((X + Y) - E[X + Y])^2 \right]$$

- ▶ We saw  $E[X + Y] = E[X] + E[Y]$ .
- ▶ Let us calculate  $\text{Var}(X + Y)$ .

$$\begin{aligned}\text{Var}(X + Y) &= E \left[ ((X + Y) - E[X + Y])^2 \right] \\ &= E \left[ ((X - EX) + (Y - EY))^2 \right]\end{aligned}$$

- ▶ We saw  $E[X + Y] = E[X] + E[Y]$ .
- ▶ Let us calculate  $\text{Var}(X + Y)$ .

$$\begin{aligned}\text{Var}(X + Y) &= E \left[ ((X + Y) - E[X + Y])^2 \right] \\ &= E \left[ ((X - EX) + (Y - EY))^2 \right] \\ &= E \left[ (X - EX)^2 \right] + E \left[ (Y - EY)^2 \right] \\ &\quad + 2E \left[ (X - EX)(Y - EY) \right]\end{aligned}$$

- ▶ We saw  $E[X + Y] = E[X] + E[Y]$ .
- ▶ Let us calculate  $\text{Var}(X + Y)$ .

$$\begin{aligned}\text{Var}(X + Y) &= E \left[ ((X + Y) - E[X + Y])^2 \right] \\ &= E \left[ ((X - EX) + (Y - EY))^2 \right] \\ &= E \left[ (X - EX)^2 \right] + E \left[ (Y - EY)^2 \right] \\ &\quad + 2E \left[ (X - EX)(Y - EY) \right] \\ &= \text{Var}(X) + \text{Var}(Y) + 2\text{Cov}(X, Y)\end{aligned}$$

- ▶ We saw  $E[X + Y] = E[X] + E[Y]$ .
- ▶ Let us calculate  $\text{Var}(X + Y)$ .

$$\begin{aligned}\text{Var}(X + Y) &= E [((X + Y) - E[X + Y])^2] \\&= E [((X - EX) + (Y - EY))^2] \\&= E [(X - EX)^2] + E [(Y - EY)^2] \\&\quad + 2E [(X - EX)(Y - EY)] \\&= \text{Var}(X) + \text{Var}(Y) + 2\text{Cov}(X, Y)\end{aligned}$$

where we define **covariance** between  $X, Y$  as

$$\text{Cov}(X, Y) = E [(X - EX)(Y - EY)]$$

- We define **covariance** between  $X$  and  $Y$  by

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)]$$



- We define **covariance** between  $X$  and  $Y$  by

$$\begin{aligned}\text{Cov}(X, Y) &= E[(X - EX)(Y - EY)] \\ &= E[XY - X(EY) - Y(EX) + EX EY]\end{aligned}$$

- We define **covariance** between  $X$  and  $Y$  by

$$\begin{aligned}\text{Cov}(X, Y) &= E[(X - EX)(Y - EY)] \\ &= E[XY - X(EY) - Y(EX) + EX EY] \\ &= E[XY] - EX EY\end{aligned}$$

- ▶ We define **covariance** between  $X$  and  $Y$  by

$$\begin{aligned}\text{Cov}(X, Y) &= E[(X - EX)(Y - EY)] \\ &= E[XY - X(EY) - Y(EX) + EX EY] \\ &= E[XY] - EX EY\end{aligned}$$

- ▶ Note that  $\text{Cov}(X, Y)$  can be positive or negative

- ▶ We define **covariance** between  $X$  and  $Y$  by

$$\begin{aligned}\text{Cov}(X, Y) &= E[(X - EX)(Y - EY)] \\ &= E[XY - X(EY) - Y(EX) + EX EY] \\ &= E[XY] - EX EY\end{aligned}$$

- ▶ Note that  $\text{Cov}(X, Y)$  can be positive or negative
- ▶  $X$  and  $Y$  are said to be uncorrelated if  $\text{Cov}(X, Y) = 0$

- We define **covariance** between  $X$  and  $Y$  by

$$\begin{aligned}\text{Cov}(X, Y) &= E[(X - EX)(Y - EY)] \\ &= E[XY - X(EY) - Y(EX) + EX EY] \\ &= E[XY] - EX EY\end{aligned}$$

- Note that  $\text{Cov}(X, Y)$  can be positive or negative
- $X$  and  $Y$  are said to be uncorrelated if  $\text{Cov}(X, Y) = 0$
- If  $X$  and  $Y$  are uncorrelated then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y)$$

- ▶ We define **covariance** between  $X$  and  $Y$  by

$$\begin{aligned}\text{Cov}(X, Y) &= E[(X - EX)(Y - EY)] \\ &= E[XY - X(EY) - Y(EX) + EX EY] \\ &= E[XY] - EX EY\end{aligned}$$

- ▶ Note that  $\text{Cov}(X, Y)$  can be positive or negative
- ▶  $X$  and  $Y$  are said to be uncorrelated if  $\text{Cov}(X, Y) = 0$
- ▶ If  $X$  and  $Y$  are uncorrelated then

$$\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y)$$

- ▶ Note that  $E[X + Y] = E[X] + E[Y]$  for all random variables.

## Example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

# Example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We want to calculate  $\text{Cov}(X, Y)$



## Example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We want to calculate  $\text{Cov}(X, Y)$

$$EX = \int_0^1 \int_x^1 x \cdot 2 \, dy \, dx$$

## Example

- ▶ Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- ▶ We want to calculate  $\text{Cov}(X, Y)$

$$EX = \int_0^1 \int_x^1 x \cdot 2 \, dy \, dx = 2 \int_0^1 x (1 - x) \, dx = \frac{1}{3}$$

## Example

- Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- We want to calculate  $\text{Cov}(X, Y)$

$$EX = \int_0^1 \int_x^1 x \cdot 2 \, dy \, dx = 2 \int_0^1 x (1 - x) \, dx = \frac{1}{3}$$

$$EY = \int_0^1 \int_0^y y \cdot 2 \, dx \, dy$$

## Example

- Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- We want to calculate  $\text{Cov}(X, Y)$

$$EX = \int_0^1 \int_x^1 x \cdot 2 \, dy \, dx = 2 \int_0^1 x (1 - x) \, dx = \frac{1}{3}$$

$$EY = \int_0^1 \int_0^y y \cdot 2 \, dx \, dy = 2 \int_0^1 y^2 \, dy = \frac{2}{3}$$

## Example

- Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- We want to calculate  $\text{Cov}(X, Y)$

$$EX = \int_0^1 \int_x^1 x \cdot 2 \, dy \, dx = 2 \int_0^1 x (1 - x) \, dx = \frac{1}{3}$$

$$EY = \int_0^1 \int_0^y y \cdot 2 \, dx \, dy = 2 \int_0^1 y^2 \, dy = \frac{2}{3}$$

$$E[XY] = \int_0^1 \int_0^y xy \cdot 2 \, dx \, dy$$

## Example

- Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- We want to calculate  $\text{Cov}(X, Y)$

$$EX = \int_0^1 \int_x^1 x \cdot 2 \, dy \, dx = 2 \int_0^1 x (1 - x) \, dx = \frac{1}{3}$$

$$EY = \int_0^1 \int_0^y y \cdot 2 \, dx \, dy = 2 \int_0^1 y^2 \, dy = \frac{2}{3}$$

$$E[XY] = \int_0^1 \int_0^y xy \cdot 2 \, dx \, dy = 2 \int_0^1 y \frac{y^2}{2} \, dy = \frac{1}{4}$$

## Example

- Consider the joint density

$$f_{XY}(x, y) = 2, \quad 0 < x < y < 1$$

- We want to calculate  $\text{Cov}(X, Y)$

$$EX = \int_0^1 \int_x^1 x \cdot 2 \, dy \, dx = 2 \int_0^1 x (1 - x) \, dx = \frac{1}{3}$$

$$EY = \int_0^1 \int_0^y y \cdot 2 \, dx \, dy = 2 \int_0^1 y^2 \, dy = \frac{2}{3}$$

$$E[XY] = \int_0^1 \int_0^y xy \cdot 2 \, dx \, dy = 2 \int_0^1 y \frac{y^2}{2} \, dy = \frac{1}{4}$$

- Hence,  $\text{Cov}(X, Y) = E[XY] - EX \cdot EY = \frac{1}{4} - \frac{2}{9} = \frac{1}{36}$

# Independent random variables are uncorrelated

- ▶ Suppose  $X, Y$  are independent. Then



# Independent random variables are uncorrelated

- Suppose  $X, Y$  are independent. Then

$$E[XY] = \int \int x y f_{XY}(x, y) dx dy$$

# Independent random variables are uncorrelated

- Suppose  $X, Y$  are independent. Then

$$\begin{aligned} E[XY] &= \int \int x y f_{XY}(x, y) dx dy \\ &= \int \int x y f_X(x) f_Y(y) dx dy \end{aligned}$$

# Independent random variables are uncorrelated

- Suppose  $X, Y$  are independent. Then

$$\begin{aligned} E[XY] &= \int \int x y f_{XY}(x, y) dx dy \\ &= \int \int x y f_X(x) f_Y(y) dx dy \\ &= \int x f_X(x) dx \int y f_Y(y) dy \end{aligned}$$

# Independent random variables are uncorrelated

- Suppose  $X, Y$  are independent. Then

$$\begin{aligned} E[XY] &= \int \int x y f_{XY}(x, y) dx dy \\ &= \int \int x y f_X(x) f_Y(y) dx dy \\ &= \int x f_X(x) dx \int y f_Y(y) dy = EX EY \end{aligned}$$

# Independent random variables are uncorrelated

- Suppose  $X, Y$  are independent. Then

$$\begin{aligned} E[XY] &= \int \int x y f_{XY}(x, y) dx dy \\ &= \int \int x y f_X(x) f_Y(y) dx dy \\ &= \int x f_X(x) dx \int y f_Y(y) dy = EX EY \end{aligned}$$

- Then,  $\text{Cov}(X, Y) = E[XY] - EX EY = 0$ .

# Independent random variables are uncorrelated

- Suppose  $X, Y$  are independent. Then

$$\begin{aligned} E[XY] &= \int \int x y f_{XY}(x, y) dx dy \\ &= \int \int x y f_X(x) f_Y(y) dx dy \\ &= \int x f_X(x) dx \int y f_Y(y) dy = EX EY \end{aligned}$$

- Then,  $\text{Cov}(X, Y) = E[XY] - EX EY = 0$ .
- $X, Y$  independent  $\Rightarrow X, Y$  uncorrelated

# Uncorrelated random variables may not be independent

- ▶ Suppose  $X \sim \mathcal{N}(0, 1)$  Then,  $EX = EX^3 = 0$

# Uncorrelated random variables may not be independent

- ▶ Suppose  $X \sim \mathcal{N}(0, 1)$  Then,  $EX = EX^3 = 0$
- ▶ Let  $Y = X^2$  Then,

$$E[XY] = EX^3 = 0$$



# Uncorrelated random variables may not be independent

- ▶ Suppose  $X \sim \mathcal{N}(0, 1)$  Then,  $EX = EX^3 = 0$
- ▶ Let  $Y = X^2$  Then,

$$E[XY] = EX^3 = 0 = EX EY$$

# Uncorrelated random variables may not be independent

- ▶ Suppose  $X \sim \mathcal{N}(0, 1)$  Then,  $EX = EX^3 = 0$
- ▶ Let  $Y = X^2$  Then,

$$E[XY] = EX^3 = 0 = EX EY$$

- ▶ Thus  $X, Y$  are uncorrelated.

# Uncorrelated random variables may not be independent

- ▶ Suppose  $X \sim \mathcal{N}(0, 1)$  Then,  $EX = EX^3 = 0$
- ▶ Let  $Y = X^2$  Then,

$$E[XY] = EX^3 = 0 = EX EY$$

- ▶ Thus  $X, Y$  are uncorrelated.
- ▶ Are they independent?

# Uncorrelated random variables may not be independent

- ▶ Suppose  $X \sim \mathcal{N}(0, 1)$  Then,  $EX = EX^3 = 0$
- ▶ Let  $Y = X^2$  Then,

$$E[XY] = EX^3 = 0 = EX EY$$

- ▶ Thus  $X, Y$  are uncorrelated.
- ▶ Are they independent? No

# Uncorrelated random variables may not be independent

- ▶ Suppose  $X \sim \mathcal{N}(0, 1)$  Then,  $EX = EX^3 = 0$
- ▶ Let  $Y = X^2$  Then,

$$E[XY] = EX^3 = 0 = EX EY$$

- ▶ Thus  $X, Y$  are uncorrelated.
- ▶ Are they independent? No  
e.g.,

$$P[X > 2 | Y < 1] = 0$$

# Uncorrelated random variables may not be independent

- ▶ Suppose  $X \sim \mathcal{N}(0, 1)$  Then,  $EX = EX^3 = 0$
- ▶ Let  $Y = X^2$  Then,

$$E[XY] = EX^3 = 0 = EX EY$$

- ▶ Thus  $X, Y$  are uncorrelated.
- ▶ Are they independent? No  
e.g.,

$$P[X > 2 | Y < 1] = 0 \neq P[X > 2]$$

# Uncorrelated random variables may not be independent

- ▶ Suppose  $X \sim \mathcal{N}(0, 1)$  Then,  $EX = EX^3 = 0$
- ▶ Let  $Y = X^2$  Then,

$$E[XY] = EX^3 = 0 = EX EY$$

- ▶ Thus  $X, Y$  are uncorrelated.
- ▶ Are they independent? No  
e.g.,

$$P[X > 2 | Y < 1] = 0 \neq P[X > 2]$$

- ▶  $X, Y$  are uncorrelated does not imply they are independent.

- We define the **correlation coefficient** of  $X, Y$  by

$$\rho_{XY} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$



- ▶ We define the **correlation coefficient** of  $X, Y$  by

$$\rho_{XY} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

- ▶ If  $X, Y$  are uncorrelated then  $\rho_{XY} = 0$ .

- ▶ We define the **correlation coefficient** of  $X, Y$  by

$$\rho_{XY} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

- ▶ If  $X, Y$  are uncorrelated then  $\rho_{XY} = 0$ .
- ▶ We will show that  $|\rho_{XY}| \leq 1$

- ▶ We define the **correlation coefficient** of  $X, Y$  by

$$\rho_{XY} = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X) \text{Var}(Y)}}$$

- ▶ If  $X, Y$  are uncorrelated then  $\rho_{XY} = 0$ .
- ▶ We will show that  $|\rho_{XY}| \leq 1$
- ▶ Hence  $-1 \leq \rho_{XY} \leq 1, \forall X, Y$

► We have  $E[(\alpha X + \beta Y)^2] \geq 0, \forall \alpha, \beta \in \mathbb{R}$

► We have  $E[(\alpha X + \beta Y)^2] \geq 0, \forall \alpha, \beta \in \mathfrak{R}$

$$\alpha^2 E[X^2] + \beta^2 E[Y^2] + 2\alpha\beta E[XY] \geq 0, \quad \forall \alpha, \beta \in \mathfrak{R}$$

► We have  $E[(\alpha X + \beta Y)^2] \geq 0, \forall \alpha, \beta \in \mathfrak{R}$

$$\alpha^2 E[X^2] + \beta^2 E[Y^2] + 2\alpha\beta E[XY] \geq 0, \quad \forall \alpha, \beta \in \mathfrak{R}$$

Take  $\alpha = -\frac{E[XY]}{E[X^2]}$

► We have  $E[(\alpha X + \beta Y)^2] \geq 0, \forall \alpha, \beta \in \mathbb{R}$

$$\alpha^2 E[X^2] + \beta^2 E[Y^2] + 2\alpha\beta E[XY] \geq 0, \quad \forall \alpha, \beta \in \mathbb{R}$$

Take  $\alpha = -\frac{E[XY]}{E[X^2]}$

$$\frac{(E[XY])^2}{E[X^2]} +$$

► We have  $E[(\alpha X + \beta Y)^2] \geq 0, \forall \alpha, \beta \in \mathbb{R}$

$$\alpha^2 E[X^2] + \beta^2 E[Y^2] + 2\alpha\beta E[XY] \geq 0, \quad \forall \alpha, \beta \in \mathbb{R}$$

$$\text{Take } \alpha = -\frac{E[XY]}{E[X^2]}$$

$$\frac{(E[XY])^2}{E[X^2]} + \beta^2 E[Y^2]$$



► We have  $E[(\alpha X + \beta Y)^2] \geq 0, \forall \alpha, \beta \in \mathfrak{R}$

$$\alpha^2 E[X^2] + \beta^2 E[Y^2] + 2\alpha\beta E[XY] \geq 0, \quad \forall \alpha, \beta \in \mathfrak{R}$$

$$\text{Take } \alpha = -\frac{E[XY]}{E[X^2]}$$

$$\frac{(E[XY])^2}{E[X^2]} + \beta^2 E[Y^2] - 2\beta \frac{(E[XY])^2}{E[X^2]} \geq 0, \quad \forall \beta \in \mathfrak{R}$$

► We have  $E[(\alpha X + \beta Y)^2] \geq 0, \forall \alpha, \beta \in \mathbb{R}$

$$\alpha^2 E[X^2] + \beta^2 E[Y^2] + 2\alpha\beta E[XY] \geq 0, \quad \forall \alpha, \beta \in \mathbb{R}$$

$$\text{Take } \alpha = -\frac{E[XY]}{E[X^2]}$$

$$\frac{(E[XY])^2}{E[X^2]} + \beta^2 E[Y^2] - 2\beta \frac{(E[XY])^2}{E[X^2]} \geq 0, \quad \forall \beta \in \mathbb{R}$$

$$a\beta^2 + b\beta + c \geq 0, \quad \forall \beta \Rightarrow b^2 - 4ac \leq 0$$

► We have  $E[(\alpha X + \beta Y)^2] \geq 0, \forall \alpha, \beta \in \mathfrak{R}$

$$\alpha^2 E[X^2] + \beta^2 E[Y^2] + 2\alpha\beta E[XY] \geq 0, \quad \forall \alpha, \beta \in \mathfrak{R}$$

$$\text{Take } \alpha = -\frac{E[XY]}{E[X^2]}$$

$$\frac{(E[XY])^2}{E[X^2]} + \beta^2 E[Y^2] - 2\beta \frac{(E[XY])^2}{E[X^2]} \geq 0, \quad \forall \beta \in \mathfrak{R}$$

$$\Rightarrow 4 \left( \frac{(E[XY])^2}{E[X^2]} \right)^2 - 4E[Y^2] \frac{(E[XY])^2}{E[X^2]} \leq 0$$

► We have  $E[(\alpha X + \beta Y)^2] \geq 0, \forall \alpha, \beta \in \mathbb{R}$

$$\alpha^2 E[X^2] + \beta^2 E[Y^2] + 2\alpha\beta E[XY] \geq 0, \quad \forall \alpha, \beta \in \mathbb{R}$$

$$\text{Take } \alpha = -\frac{E[XY]}{E[X^2]}$$

$$\frac{(E[XY])^2}{E[X^2]} + \beta^2 E[Y^2] - 2\beta \frac{(E[XY])^2}{E[X^2]} \geq 0, \quad \forall \beta \in \mathbb{R}$$

$$\Rightarrow 4 \left( \frac{(E[XY])^2}{E[X^2]} \right)^2 - 4E[Y^2] \frac{(E[XY])^2}{E[X^2]} \leq 0$$

$$\Rightarrow \left( \frac{(E[XY])^2}{E[X^2]} \right)^2 \leq \frac{E[Y^2](E[XY])^2}{E[X^2]}$$

► We have  $E[(\alpha X + \beta Y)^2] \geq 0, \forall \alpha, \beta \in \mathbb{R}$

$$\alpha^2 E[X^2] + \beta^2 E[Y^2] + 2\alpha\beta E[XY] \geq 0, \quad \forall \alpha, \beta \in \mathbb{R}$$

$$\text{Take } \alpha = -\frac{E[XY]}{E[X^2]}$$

$$\frac{(E[XY])^2}{E[X^2]} + \beta^2 E[Y^2] - 2\beta \frac{(E[XY])^2}{E[X^2]} \geq 0, \quad \forall \beta \in \mathbb{R}$$

$$\Rightarrow 4 \left( \frac{(E[XY])^2}{E[X^2]} \right)^2 - 4E[Y^2] \frac{(E[XY])^2}{E[X^2]} \leq 0$$

$$\Rightarrow \left( \frac{(E[XY])^2}{E[X^2]} \right)^2 \leq \frac{E[Y^2](E[XY])^2}{E[X^2]}$$

$$\Rightarrow \frac{(E[XY])^4}{(E[XY])^2} \leq \frac{E[Y^2](E[X^2])^2}{E[X^2]}$$

► We have  $E[(\alpha X + \beta Y)^2] \geq 0, \forall \alpha, \beta \in \mathbb{R}$

$$\alpha^2 E[X^2] + \beta^2 E[Y^2] + 2\alpha\beta E[XY] \geq 0, \quad \forall \alpha, \beta \in \mathbb{R}$$

$$\text{Take } \alpha = -\frac{E[XY]}{E[X^2]}$$

$$\frac{(E[XY])^2}{E[X^2]} + \beta^2 E[Y^2] - 2\beta \frac{(E[XY])^2}{E[X^2]} \geq 0, \quad \forall \beta \in \mathbb{R}$$

$$\Rightarrow 4 \left( \frac{(E[XY])^2}{E[X^2]} \right)^2 - 4E[Y^2] \frac{(E[XY])^2}{E[X^2]} \leq 0$$

$$\Rightarrow \left( \frac{(E[XY])^2}{E[X^2]} \right)^2 \leq \frac{E[Y^2](E[XY])^2}{E[X^2]}$$

$$\Rightarrow \frac{(E[XY])^4}{(E[XY])^2} \leq \frac{E[Y^2](E[X^2])^2}{E[X^2]}$$

$$\Rightarrow (E[XY])^2 \leq E[X^2]E[Y^2]$$

► We showed that

$$(E[XY])^2 \leq E[X^2]E[Y^2]$$

- ▶ We showed that

$$(E[XY])^2 \leq E[X^2]E[Y^2]$$

- ▶ Take  $X - EX$  in place of  $X$  and  $Y - EY$  in place of  $Y$  in the above algebra.



- ▶ We showed that

$$(E[XY])^2 \leq E[X^2]E[Y^2]$$

- ▶ Take  $X - EX$  in place of  $X$  and  $Y - EY$  in place of  $Y$  in the above algebra.
- ▶ This gives us

$$(E[(X - EX)(Y - EY)])^2 \leq E[(X - EX)^2]E[(Y - EY)^2]$$

- ▶ We showed that

$$(E[XY])^2 \leq E[X^2]E[Y^2]$$

- ▶ Take  $X - EX$  in place of  $X$  and  $Y - EY$  in place of  $Y$  in the above algebra.
- ▶ This gives us

$$(E[(X - EX)(Y - EY)])^2 \leq E[(X - EX)^2]E[(Y - EY)^2]$$

$$\Rightarrow (\text{Cov}(X, Y))^2 \leq \text{Var}(X)\text{Var}(Y)$$

- ▶ We showed that

$$(E[XY])^2 \leq E[X^2]E[Y^2]$$

- ▶ Take  $X - EX$  in place of  $X$  and  $Y - EY$  in place of  $Y$  in the above algebra.
- ▶ This gives us

$$(E[(X - EX)(Y - EY)])^2 \leq E[(X - EX)^2]E[(Y - EY)^2]$$

$$\Rightarrow (\text{Cov}(X, Y))^2 \leq \text{Var}(X)\text{Var}(Y)$$

- ▶ Hence we get

$$\rho_{XY}^2 = \left( \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \right)^2 \leq 1$$

- ▶ We showed that

$$(E[XY])^2 \leq E[X^2]E[Y^2]$$

- ▶ Take  $X - EX$  in place of  $X$  and  $Y - EY$  in place of  $Y$  in the above algebra.
- ▶ This gives us

$$(E[(X - EX)(Y - EY)])^2 \leq E[(X - EX)^2]E[(Y - EY)^2]$$

$$\Rightarrow (\text{Cov}(X, Y))^2 \leq \text{Var}(X)\text{Var}(Y)$$

- ▶ Hence we get

$$\rho_{XY}^2 = \left( \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \right)^2 \leq 1$$

- ▶ The equality holds here only if  $E[(\alpha X + \beta Y)^2] = 0$

- ▶ We showed that

$$(E[XY])^2 \leq E[X^2]E[Y^2]$$

- ▶ Take  $X - EX$  in place of  $X$  and  $Y - EY$  in place of  $Y$  in the above algebra.
- ▶ This gives us

$$(E[(X - EX)(Y - EY)])^2 \leq E[(X - EX)^2]E[(Y - EY)^2]$$

$$\Rightarrow (\text{Cov}(X, Y))^2 \leq \text{Var}(X)\text{Var}(Y)$$

- ▶ Hence we get

$$\rho_{XY}^2 = \left( \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \right)^2 \leq 1$$

- ▶ The equality holds here only if  $E[(\alpha X + \beta Y)^2] = 0$

$$\text{Thus, } |\rho_{XY}| = 1 \text{ only if } \alpha X + \beta Y = 0$$

- ▶ We showed that

$$(E[XY])^2 \leq E[X^2]E[Y^2]$$

- ▶ Take  $X - EX$  in place of  $X$  and  $Y - EY$  in place of  $Y$  in the above algebra.
- ▶ This gives us

$$(E[(X - EX)(Y - EY)])^2 \leq E[(X - EX)^2]E[(Y - EY)^2]$$

$$\Rightarrow (\text{Cov}(X, Y))^2 \leq \text{Var}(X)\text{Var}(Y)$$

- ▶ Hence we get

$$\rho_{XY}^2 = \left( \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}} \right)^2 \leq 1$$

- ▶ The equality holds here only if  $E[(\alpha X + \beta Y)^2] = 0$

$$\text{Thus, } |\rho_{XY}| = 1 \text{ only if } \alpha X + \beta Y = 0$$

- ▶ Correlation coefficient of  $X, Y$  is  $\pm 1$  only when  $Y$  is a linear function of  $X$

# Linear Least Squares Estimation

- Suppose we want to approximate  $Y$  as an affine function of  $X$ .

# Linear Least Squares Estimation

- ▶ Suppose we want to approximate  $Y$  as an affine function of  $X$ .
- ▶ We want  $a, b$  to minimize  $E[(Y - (aX + b))^2]$



# Linear Least Squares Estimation

- ▶ Suppose we want to approximate  $Y$  as an affine function of  $X$ .
- ▶ We want  $a, b$  to minimize  $E[(Y - (aX + b))^2]$
- ▶ For a fixed  $a$ , what is the  $b$  that minimizes  $E[((Y - aX) - b)^2]$  ?

# Linear Least Squares Estimation

- ▶ Suppose we want to approximate  $Y$  as an affine function of  $X$ .
- ▶ We want  $a, b$  to minimize  $E[(Y - (aX + b))^2]$
- ▶ For a fixed  $a$ , what is the  $b$  that minimizes  $E[((Y - aX) - b)^2]$  ?
- ▶ We know the best  $b$  here is:  
$$b = E[Y - aX] = EY - aEX.$$

# Linear Least Squares Estimation

- ▶ Suppose we want to approximate  $Y$  as an affine function of  $X$ .
- ▶ We want  $a, b$  to minimize  $E[(Y - (aX + b))^2]$
- ▶ For a fixed  $a$ , what is the  $b$  that minimizes  $E[((Y - aX) - b)^2]$  ?
- ▶ We know the best  $b$  here is:  
$$b = E[Y - aX] = EY - aEX.$$
- ▶ So, we want to find the best  $a$  to minimize  $J(a) = E[(Y - aX - (EY - aEX))^2]$

► We want to find  $a$  to minimize

$$J(a) = E[(Y - aX - (EY - aEX))^2]$$

► We want to find  $a$  to minimize

$$\begin{aligned} J(a) &= E[(Y - aX - (EY - aEX))^2] \\ &= E[((Y - EY) - a(X - EX))^2] \end{aligned}$$

► We want to find  $a$  to minimize

$$\begin{aligned} J(a) &= E[(Y - aX - (EY - aEX))^2] \\ &= E[((Y - EY) - a(X - EX))^2] \\ &= E[(Y - EY)^2 + a^2(X - EX)^2 - 2a(Y - EY)(X - EX)] \end{aligned}$$

► We want to find  $a$  to minimize

$$\begin{aligned} J(a) &= E[(Y - aX - (EY - aEX))^2] \\ &= E[((Y - EY) - a(X - EX))^2] \\ &= E[(Y - EY)^2 + a^2(X - EX)^2 - 2a(Y - EY)(X - EX)] \\ &= \text{Var}(Y) + a^2\text{Var}(X) - 2a\text{Cov}(X, Y) \end{aligned}$$

- We want to find  $a$  to minimize

$$\begin{aligned} J(a) &= E[(Y - aX - (EY - aEX))^2] \\ &= E[((Y - EY) - a(X - EX))^2] \\ &= E[(Y - EY)^2 + a^2(X - EX)^2 - 2a(Y - EY)(X - EX)] \\ &= \text{Var}(Y) + a^2\text{Var}(X) - 2a\text{Cov}(X, Y) \end{aligned}$$

- So, the optimal  $a$  satisfies

$$2a\text{Var}(X) - 2\text{Cov}(X, Y) = 0$$



- We want to find  $a$  to minimize

$$\begin{aligned} J(a) &= E[(Y - aX - (EY - aEX))^2] \\ &= E[((Y - EY) - a(X - EX))^2] \\ &= E[(Y - EY)^2 + a^2(X - EX)^2 - 2a(Y - EY)(X - EX)] \\ &= \text{Var}(Y) + a^2\text{Var}(X) - 2a\text{Cov}(X, Y) \end{aligned}$$

- So, the optimal  $a$  satisfies

$$2a\text{Var}(X) - 2\text{Cov}(X, Y) = 0 \Rightarrow a = \frac{\text{Cov}(X, Y)}{\text{Var}(X)}$$

- The final mean square error, say,  $J^*$  is

$$J^* = \text{Var}(Y) + a^2\text{Var}(X) - 2a\text{Cov}(X, Y)$$

- The final mean square error, say,  $J^*$  is

$$\begin{aligned} J^* &= \text{Var}(Y) + a^2 \text{Var}(X) - 2a \text{Cov}(X, Y) \\ &= \text{Var}(Y) + \left( \frac{\text{Cov}(X, Y)}{\text{Var}(X)} \right)^2 \text{Var}(X) - 2 \frac{\text{Cov}(X, Y)}{\text{Var}(X)} \text{Cov}(X, Y) \end{aligned}$$

- The final mean square error, say,  $J^*$  is

$$\begin{aligned} J^* &= \text{Var}(Y) + a^2 \text{Var}(X) - 2a \text{Cov}(X, Y) \\ &= \text{Var}(Y) + \left( \frac{\text{Cov}(X, Y)}{\text{Var}(X)} \right)^2 \text{Var}(X) - 2 \frac{\text{Cov}(X, Y)}{\text{Var}(X)} \text{Cov}(X, Y) \\ &= \text{Var}(Y) - \frac{(\text{Cov}(X, Y))^2}{\text{Var}(X)} \end{aligned}$$

- The final mean square error, say,  $J^*$  is

$$\begin{aligned} J^* &= \text{Var}(Y) + a^2 \text{Var}(X) - 2a \text{Cov}(X, Y) \\ &= \text{Var}(Y) + \left( \frac{\text{Cov}(X, Y)}{\text{Var}(X)} \right)^2 \text{Var}(X) - 2 \frac{\text{Cov}(X, Y)}{\text{Var}(X)} \text{Cov}(X, Y) \\ &= \text{Var}(Y) - \frac{(\text{Cov}(X, Y))^2}{\text{Var}(X)} \\ &= \text{Var}(Y) \left( 1 - \frac{(\text{Cov}(X, Y))^2}{\text{Var}(Y) \text{Var}(X)} \right) \\ &= \text{Var}(Y) (1 - \rho_{XY}^2) \end{aligned}$$

- The best mean-square approximation of  $Y$  as a 'linear' function of  $X$  is

$$Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)} X + \left( EY - \frac{\text{Cov}(X, Y)}{\text{Var}(X)} EX \right)$$

- ▶ The best mean-square approximation of  $Y$  as a 'linear' function of  $X$  is

$$Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)} X + \left( EY - \frac{\text{Cov}(X, Y)}{\text{Var}(X)} EX \right)$$

- ▶ Called the line of regression of  $Y$  on  $X$ .

- ▶ The best mean-square approximation of  $Y$  as a 'linear' function of  $X$  is

$$Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)} X + \left( EY - \frac{\text{Cov}(X, Y)}{\text{Var}(X)} EX \right)$$

- ▶ Called the line of regression of  $Y$  on  $X$ .
- ▶ If  $\text{cov}(X, Y) = 0$  then this reduces to approximating  $Y$  by a constant,  $EY$ .



- ▶ The best mean-square approximation of  $Y$  as a 'linear' function of  $X$  is

$$Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)} X + \left( EY - \frac{\text{Cov}(X, Y)}{\text{Var}(X)} EX \right)$$

- ▶ Called the line of regression of  $Y$  on  $X$ .
- ▶ If  $\text{cov}(X, Y) = 0$  then this reduces to approximating  $Y$  by a constant,  $EY$ .
- ▶ The final mean square error is

$$\text{Var}(Y) (1 - \rho_{XY}^2)$$

- ▶ The best mean-square approximation of  $Y$  as a 'linear' function of  $X$  is

$$Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)} X + \left( EY - \frac{\text{Cov}(X, Y)}{\text{Var}(X)} EX \right)$$

- ▶ Called the line of regression of  $Y$  on  $X$ .
- ▶ If  $\text{cov}(X, Y) = 0$  then this reduces to approximating  $Y$  by a constant,  $EY$ .
- ▶ The final mean square error is

$$\text{Var}(Y) (1 - \rho_{XY}^2)$$

- ▶ If  $\rho_{XY} = \pm 1$  then the error is zero

- ▶ The best mean-square approximation of  $Y$  as a 'linear' function of  $X$  is

$$Y = \frac{\text{Cov}(X, Y)}{\text{Var}(X)} X + \left( EY - \frac{\text{Cov}(X, Y)}{\text{Var}(X)} EX \right)$$

- ▶ Called the line of regression of  $Y$  on  $X$ .
- ▶ If  $\text{cov}(X, Y) = 0$  then this reduces to approximating  $Y$  by a constant,  $EY$ .
- ▶ The final mean square error is

$$\text{Var}(Y) (1 - \rho_{XY}^2)$$

- ▶ If  $\rho_{XY} = \pm 1$  then the error is zero
- ▶ If  $\rho_{XY} = 0$  the final error is  $\text{Var}(Y)$

- ▶ The covariance of  $X, Y$  is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)]$$

- ▶ The covariance of  $X, Y$  is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

- The covariance of  $X, Y$  is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Note that  $\text{Cov}(X, X) = \text{Var}(X)$

- ▶ The covariance of  $X, Y$  is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Note that  $\text{Cov}(X, X) = \text{Var}(X)$

- ▶  $X, Y$  are called uncorrelated if  $\text{Cov}(X, Y) = 0$ .

- ▶ The covariance of  $X, Y$  is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Note that  $\text{Cov}(X, X) = \text{Var}(X)$

- ▶  $X, Y$  are called uncorrelated if  $\text{Cov}(X, Y) = 0$ .
- ▶  $X, Y$  independent  $\Rightarrow X, Y$  uncorrelated.



- ▶ The covariance of  $X, Y$  is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Note that  $\text{Cov}(X, X) = \text{Var}(X)$

- ▶  $X, Y$  are called uncorrelated if  $\text{Cov}(X, Y) = 0$ .
- ▶  $X, Y$  independent  $\Rightarrow X, Y$  uncorrelated.
- ▶ Uncorrelated random variables need not necessarily be independent

- ▶ The covariance of  $X, Y$  is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Note that  $\text{Cov}(X, X) = \text{Var}(X)$

- ▶  $X, Y$  are called uncorrelated if  $\text{Cov}(X, Y) = 0$ .
- ▶  $X, Y$  independent  $\Rightarrow X, Y$  uncorrelated.
- ▶ Uncorrelated random variables need not necessarily be independent
- ▶ Covariance plays an important role in linear least squares estimation.

- ▶ The covariance of  $X, Y$  is

$$\text{Cov}(X, Y) = E[(X - EX)(Y - EY)] = E[XY] - EX EY$$

Note that  $\text{Cov}(X, X) = \text{Var}(X)$

- ▶  $X, Y$  are called uncorrelated if  $\text{Cov}(X, Y) = 0$ .
- ▶  $X, Y$  independent  $\Rightarrow X, Y$  uncorrelated.
- ▶ Uncorrelated random variables need not necessarily be independent
- ▶ Covariance plays an important role in linear least squares estimation.
- ▶ Informally, covariance captures the 'linear dependence' between the two random variables.

# Covariance Matrix

- ▶ Let  $X_1, \dots, X_n$  be random variables (on the same probability space)

# Covariance Matrix

- ▶ Let  $X_1, \dots, X_n$  be random variables (on the same probability space)
- ▶ We represent them as a vector  $\mathbf{X}$ .

# Covariance Matrix

- ▶ Let  $X_1, \dots, X_n$  be random variables (on the same probability space)
- ▶ We represent them as a vector  $\mathbf{X}$ .
- ▶ As a notation, all vectors are column vectors:

# Covariance Matrix

- ▶ Let  $X_1, \dots, X_n$  be random variables (on the same probability space)
- ▶ We represent them as a vector  $\mathbf{X}$ .
- ▶ As a notation, all vectors are column vectors:  
 $\mathbf{X} = (X_1, \dots, X_n)^T$

# Covariance Matrix

- ▶ Let  $X_1, \dots, X_n$  be random variables (on the same probability space)
- ▶ We represent them as a vector  $\mathbf{X}$ .
- ▶ As a notation, all vectors are column vectors:  
 $\mathbf{X} = (X_1, \dots, X_n)^T$
- ▶ We denote  $E[\mathbf{X}] = (EX_1, \dots, EX_n)^T$



# Covariance Matrix

- ▶ Let  $X_1, \dots, X_n$  be random variables (on the same probability space)
- ▶ We represent them as a vector  $\mathbf{X}$ .
- ▶ As a notation, all vectors are column vectors:  
 $\mathbf{X} = (X_1, \dots, X_n)^T$
- ▶ We denote  $E[\mathbf{X}] = (EX_1, \dots, EX_n)^T$
- ▶ The  $n \times n$  matrix whose  $(i, j)^{th}$  element is  $\text{Cov}(X_i, X_j)$  is called the covariance matrix (or variance-covariance matrix) of  $\mathbf{X}$ . Denoted as  $\Sigma_{\mathbf{X}}$  or  $\Sigma_X$

# Covariance Matrix

- ▶ Let  $X_1, \dots, X_n$  be random variables (on the same probability space)
- ▶ We represent them as a vector  $\mathbf{X}$ .
- ▶ As a notation, all vectors are column vectors:  
 $\mathbf{X} = (X_1, \dots, X_n)^T$
- ▶ We denote  $E[\mathbf{X}] = (EX_1, \dots, EX_n)^T$
- ▶ The  $n \times n$  matrix whose  $(i, j)^{th}$  element is  $\text{Cov}(X_i, X_j)$  is called the covariance matrix (or variance-covariance matrix) of  $\mathbf{X}$ . Denoted as  $\Sigma_{\mathbf{X}}$  or  $\Sigma_X$

$$\Sigma_{\mathbf{X}} = \begin{bmatrix} \text{Cov}(X_1, X_1) & \text{Cov}(X_1, X_2) & \cdots & \text{Cov}(X_1, X_n) \\ \text{Cov}(X_2, X_1) & \text{Cov}(X_2, X_2) & \cdots & \text{Cov}(X_2, X_n) \\ \vdots & \vdots & \vdots & \vdots \\ \text{Cov}(X_n, X_1) & \text{Cov}(X_n, X_2) & \cdots & \text{Cov}(X_n, X_n) \end{bmatrix}$$

# Covariance matrix

- ▶ If  $\mathbf{a} = (a_1, \dots, a_n)^T$  then  $\mathbf{a} \mathbf{a}^T$  is a  $n \times n$  matrix whose  $(i, j)^{th}$  element is  $a_i a_j$ .

# Covariance matrix

- ▶ If  $\mathbf{a} = (a_1, \dots, a_n)^T$  then  $\mathbf{a} \mathbf{a}^T$  is a  $n \times n$  matrix whose  $(i, j)^{th}$  element is  $a_i a_j$ .
- ▶ Hence we get

$$\Sigma_{\mathbf{X}} = E [(\mathbf{X} - E\mathbf{X})(\mathbf{X} - E\mathbf{X})^T]$$

# Covariance matrix

- ▶ If  $\mathbf{a} = (a_1, \dots, a_n)^T$  then  $\mathbf{a} \mathbf{a}^T$  is a  $n \times n$  matrix whose  $(i, j)^{th}$  element is  $a_i a_j$ .
- ▶ Hence we get

$$\Sigma_{\mathbf{X}} = E [(\mathbf{X} - E\mathbf{X})(\mathbf{X} - E\mathbf{X})^T]$$

- ▶ This is because  $((\mathbf{X} - E\mathbf{X})(\mathbf{X} - E\mathbf{X})^T)_{ij} = (X_i - EX_i)(X_j - EX_j)$

# Covariance matrix

- ▶ If  $\mathbf{a} = (a_1, \dots, a_n)^T$  then  $\mathbf{a} \mathbf{a}^T$  is a  $n \times n$  matrix whose  $(i, j)^{th}$  element is  $a_i a_j$ .
- ▶ Hence we get

$$\Sigma_{\mathbf{X}} = E [(\mathbf{X} - E\mathbf{X})(\mathbf{X} - E\mathbf{X})^T]$$

- ▶ This is because  
 $((\mathbf{X} - E\mathbf{X})(\mathbf{X} - E\mathbf{X})^T)_{ij} = (X_i - EX_i)(X_j - EX_j)$   
and  $(\Sigma_{\mathbf{X}})_{ij} = E[(X_i - EX_i)(X_j - EX_j)]$

- ▶ Recall the following about vectors and matrices

- ▶ Recall the following about vectors and matrices
- ▶ let  $\mathbf{a}, \mathbf{b} \in \Re^n$  be column vectors. Then

$$(\mathbf{a}^T \mathbf{b})^2 = (\mathbf{a}^T \mathbf{b})^T (\mathbf{a}^T \mathbf{b})$$



- ▶ Recall the following about vectors and matrices
- ▶ let  $\mathbf{a}, \mathbf{b} \in \Re^n$  be column vectors. Then

$$(\mathbf{a}^T \mathbf{b})^2 = (\mathbf{a}^T \mathbf{b})^T (\mathbf{a}^T \mathbf{b}) = \mathbf{b}^T \mathbf{a} \mathbf{a}^T \mathbf{b}$$

- ▶ Recall the following about vectors and matrices
- ▶ let  $\mathbf{a}, \mathbf{b} \in \Re^n$  be column vectors. Then

$$(\mathbf{a}^T \mathbf{b})^2 = (\mathbf{a}^T \mathbf{b})^T (\mathbf{a}^T \mathbf{b}) = \mathbf{b}^T \mathbf{a} \mathbf{a}^T \mathbf{b} = \mathbf{b}^T (\mathbf{a} \mathbf{a}^T) \mathbf{b}$$

- ▶ Recall the following about vectors and matrices
- ▶ let  $\mathbf{a}, \mathbf{b} \in \Re^n$  be column vectors. Then

$$(\mathbf{a}^T \mathbf{b})^2 = (\mathbf{a}^T \mathbf{b})^T (\mathbf{a}^T \mathbf{b}) = \mathbf{b}^T \mathbf{a} \mathbf{a}^T \mathbf{b} = \mathbf{b}^T (\mathbf{a} \mathbf{a}^T) \mathbf{b}$$

- ▶ Let  $A$  be an  $n \times n$  matrix with elements  $a_{ij}$ . Then

$$\mathbf{b}^T A \mathbf{b} = \sum_{i,j=1}^n b_i b_j a_{ij}$$

where  $\mathbf{b} = (b_1, \dots, b_n)^T$

- ▶ Recall the following about vectors and matrices
- ▶ let  $\mathbf{a}, \mathbf{b} \in \Re^n$  be column vectors. Then

$$(\mathbf{a}^T \mathbf{b})^2 = (\mathbf{a}^T \mathbf{b})^T (\mathbf{a}^T \mathbf{b}) = \mathbf{b}^T \mathbf{a} \mathbf{a}^T \mathbf{b} = \mathbf{b}^T (\mathbf{a} \mathbf{a}^T) \mathbf{b}$$

- ▶ Let  $A$  be an  $n \times n$  matrix with elements  $a_{ij}$ . Then

$$\mathbf{b}^T A \mathbf{b} = \sum_{i,j=1}^n b_i b_j a_{ij}$$

where  $\mathbf{b} = (b_1, \dots, b_n)^T$

- ▶  $A$  is said to be positive semidefinite if  $\mathbf{b}^T A \mathbf{b} \geq 0, \forall \mathbf{b}$

- ▶  $\Sigma_X$  is a real symmetric matrix

- ▶  $\Sigma_X$  is a real symmetric matrix
- ▶ It is positive semidefinite.

- ▶  $\Sigma_X$  is a real symmetric matrix
- ▶ It is positive semidefinite.
- ▶ Let  $\mathbf{a} \in \Re^n$  and let  $Y = \mathbf{a}^T \mathbf{X}$ .

- ▶  $\Sigma_X$  is a real symmetric matrix
- ▶ It is positive semidefinite.
- ▶ Let  $\mathbf{a} \in \Re^n$  and let  $Y = \mathbf{a}^T \mathbf{X}$ .
- ▶ Then,  $EY = \mathbf{a}^T E\mathbf{X}$ . We get variance of  $Y$  as

$$\text{Var}(Y) = E[(Y - EY)^2] = E \left[ (\mathbf{a}^T \mathbf{X} - \mathbf{a}^T E\mathbf{X})^2 \right]$$



- ▶  $\Sigma_X$  is a real symmetric matrix
- ▶ It is positive semidefinite.
- ▶ Let  $\mathbf{a} \in \Re^n$  and let  $Y = \mathbf{a}^T \mathbf{X}$ .
- ▶ Then,  $EY = \mathbf{a}^T E\mathbf{X}$ . We get variance of  $Y$  as

$$\begin{aligned}\text{Var}(Y) &= E[(Y - EY)^2] = E \left[ (\mathbf{a}^T \mathbf{X} - \mathbf{a}^T E\mathbf{X})^2 \right] \\ &= E \left[ (\mathbf{a}^T (\mathbf{X} - E\mathbf{X}))^2 \right]\end{aligned}$$

- ▶  $\Sigma_X$  is a real symmetric matrix
- ▶ It is positive semidefinite.
- ▶ Let  $\mathbf{a} \in \Re^n$  and let  $Y = \mathbf{a}^T \mathbf{X}$ .
- ▶ Then,  $EY = \mathbf{a}^T E\mathbf{X}$ . We get variance of  $Y$  as

$$\begin{aligned}\text{Var}(Y) &= E[(Y - EY)^2] = E \left[ (\mathbf{a}^T \mathbf{X} - \mathbf{a}^T E\mathbf{X})^2 \right] \\ &= E \left[ (\mathbf{a}^T (\mathbf{X} - E\mathbf{X}))^2 \right] \\ &= E \left[ \mathbf{a}^T (\mathbf{X} - E\mathbf{X}) (\mathbf{X} - E\mathbf{X})^T \mathbf{a} \right]\end{aligned}$$

- ▶  $\Sigma_X$  is a real symmetric matrix
- ▶ It is positive semidefinite.
- ▶ Let  $\mathbf{a} \in \Re^n$  and let  $Y = \mathbf{a}^T \mathbf{X}$ .
- ▶ Then,  $EY = \mathbf{a}^T E\mathbf{X}$ . We get variance of  $Y$  as

$$\begin{aligned}
 \text{Var}(Y) &= E[(Y - EY)^2] = E \left[ (\mathbf{a}^T \mathbf{X} - \mathbf{a}^T E\mathbf{X})^2 \right] \\
 &= E \left[ (\mathbf{a}^T (\mathbf{X} - E\mathbf{X}))^2 \right] \\
 &= E \left[ \mathbf{a}^T (\mathbf{X} - E\mathbf{X}) (\mathbf{X} - E\mathbf{X})^T \mathbf{a} \right] \\
 &= \mathbf{a}^T E \left[ (\mathbf{X} - E\mathbf{X}) (\mathbf{X} - E\mathbf{X})^T \right] \mathbf{a}
 \end{aligned}$$

- ▶  $\Sigma_X$  is a real symmetric matrix
- ▶ It is positive semidefinite.
- ▶ Let  $\mathbf{a} \in \Re^n$  and let  $Y = \mathbf{a}^T \mathbf{X}$ .
- ▶ Then,  $EY = \mathbf{a}^T E\mathbf{X}$ . We get variance of  $Y$  as

$$\begin{aligned}
 \text{Var}(Y) &= E[(Y - EY)^2] = E \left[ (\mathbf{a}^T \mathbf{X} - \mathbf{a}^T E\mathbf{X})^2 \right] \\
 &= E \left[ (\mathbf{a}^T (\mathbf{X} - E\mathbf{X}))^2 \right] \\
 &= E \left[ \mathbf{a}^T (\mathbf{X} - E\mathbf{X}) (\mathbf{X} - E\mathbf{X})^T \mathbf{a} \right] \\
 &= \mathbf{a}^T E \left[ (\mathbf{X} - E\mathbf{X}) (\mathbf{X} - E\mathbf{X})^T \right] \mathbf{a} \\
 &= \mathbf{a}^T \Sigma_X \mathbf{a}
 \end{aligned}$$

- ▶  $\Sigma_X$  is a real symmetric matrix
- ▶ It is positive semidefinite.
- ▶ Let  $\mathbf{a} \in \Re^n$  and let  $Y = \mathbf{a}^T \mathbf{X}$ .
- ▶ Then,  $EY = \mathbf{a}^T E\mathbf{X}$ . We get variance of  $Y$  as

$$\begin{aligned}
 \text{Var}(Y) &= E[(Y - EY)^2] = E \left[ (\mathbf{a}^T \mathbf{X} - \mathbf{a}^T E\mathbf{X})^2 \right] \\
 &= E \left[ (\mathbf{a}^T (\mathbf{X} - E\mathbf{X}))^2 \right] \\
 &= E \left[ \mathbf{a}^T (\mathbf{X} - E\mathbf{X}) (\mathbf{X} - E\mathbf{X})^T \mathbf{a} \right] \\
 &= \mathbf{a}^T E \left[ (\mathbf{X} - E\mathbf{X}) (\mathbf{X} - E\mathbf{X})^T \right] \mathbf{a} \\
 &= \mathbf{a}^T \Sigma_X \mathbf{a}
 \end{aligned}$$

- ▶ This gives  $\mathbf{a}^T \Sigma_X \mathbf{a} \geq 0, \forall \mathbf{a}$

- ▶  $\Sigma_X$  is a real symmetric matrix
- ▶ It is positive semidefinite.
- ▶ Let  $\mathbf{a} \in \Re^n$  and let  $Y = \mathbf{a}^T \mathbf{X}$ .
- ▶ Then,  $EY = \mathbf{a}^T E\mathbf{X}$ . We get variance of  $Y$  as

$$\begin{aligned}
 \text{Var}(Y) &= E[(Y - EY)^2] = E \left[ (\mathbf{a}^T \mathbf{X} - \mathbf{a}^T E\mathbf{X})^2 \right] \\
 &= E \left[ (\mathbf{a}^T (\mathbf{X} - E\mathbf{X}))^2 \right] \\
 &= E \left[ \mathbf{a}^T (\mathbf{X} - E\mathbf{X}) (\mathbf{X} - E\mathbf{X})^T \mathbf{a} \right] \\
 &= \mathbf{a}^T E \left[ (\mathbf{X} - E\mathbf{X}) (\mathbf{X} - E\mathbf{X})^T \right] \mathbf{a} \\
 &= \mathbf{a}^T \Sigma_X \mathbf{a}
 \end{aligned}$$

- ▶ This gives  $\mathbf{a}^T \Sigma_X \mathbf{a} \geq 0, \forall \mathbf{a}$
- ▶ This shows  $\Sigma_X$  is positive semidefinite

►  $Y = \mathbf{a}^T \mathbf{X} = \sum_i a_i X_i$

►  $Y = \mathbf{a}^T \mathbf{X} = \sum_i a_i X_i$  – linear combination of  $X_i$ 's.



- ▶  $Y = \mathbf{a}^T \mathbf{X} = \sum_i a_i X_i$  – linear combination of  $X_i$ 's.
- ▶ We know how to find its mean and variance

- ▶  $Y = \mathbf{a}^T \mathbf{X} = \sum_i a_i X_i$  – linear combination of  $X_i$ 's.
- ▶ We know how to find its mean and variance

$$EY = \mathbf{a}^T EX = \sum_i a_i EX_i;$$

- ▶  $Y = \mathbf{a}^T \mathbf{X} = \sum_i a_i X_i$  – linear combination of  $X_i$ 's.
- ▶ We know how to find its mean and variance

$$EY = \mathbf{a}^T E\mathbf{X} = \sum_i a_i EX_i;$$

$$\text{Var}(Y) = \mathbf{a}^T \Sigma_X \mathbf{a} = \sum_{i,j} a_i a_j \text{Cov}(X_i, X_j)$$

- ▶  $Y = \mathbf{a}^T \mathbf{X} = \sum_i a_i X_i$  – linear combination of  $X_i$ 's.
- ▶ We know how to find its mean and variance

$$EY = \mathbf{a}^T E\mathbf{X} = \sum_i a_i EX_i;$$

$$\text{Var}(Y) = \mathbf{a}^T \Sigma_X \mathbf{a} = \sum_{i,j} a_i a_j \text{Cov}(X_i, X_j)$$

- ▶ Specifically, by taking all components of  $\mathbf{a}$  to be 1, we get

$$\text{Var}\left(\sum_{i=1}^n X_i\right) = \sum_{i,j=1}^n \text{Cov}(X_i, X_j)$$

- ▶  $Y = \mathbf{a}^T \mathbf{X} = \sum_i a_i X_i$  – linear combination of  $X_i$ 's.
- ▶ We know how to find its mean and variance

$$EY = \mathbf{a}^T E\mathbf{X} = \sum_i a_i EX_i;$$

$$\text{Var}(Y) = \mathbf{a}^T \Sigma_X \mathbf{a} = \sum_{i,j} a_i a_j \text{Cov}(X_i, X_j)$$

- ▶ Specifically, by taking all components of  $\mathbf{a}$  to be 1, we get

$$\text{Var}\left(\sum_{i=1}^n X_i\right) = \sum_{i,j=1}^n \text{Cov}(X_i, X_j) = \sum_{i=1}^n \text{Var}(X_i) + \sum_{i=1}^n \sum_{j \neq i} \text{Cov}(X_i, X_j)$$

- ▶  $Y = \mathbf{a}^T \mathbf{X} = \sum_i a_i X_i$  – linear combination of  $X_i$ 's.
- ▶ We know how to find its mean and variance

$$EY = \mathbf{a}^T E\mathbf{X} = \sum_i a_i EX_i;$$

$$\text{Var}(Y) = \mathbf{a}^T \Sigma_X \mathbf{a} = \sum_{i,j} a_i a_j \text{Cov}(X_i, X_j)$$

- ▶ Specifically, by taking all components of  $\mathbf{a}$  to be 1, we get

$$\text{Var}\left(\sum_{i=1}^n X_i\right) = \sum_{i,j=1}^n \text{Cov}(X_i, X_j) = \sum_{i=1}^n \text{Var}(X_i) + \sum_{i=1}^n \sum_{j \neq i} \text{Cov}(X_i, X_j)$$

- ▶ If  $X_i$  are independent, variance of sum is sum of variances.

- Covariance matrix  $\Sigma_X$  positive semidefinite because

$$\mathbf{a}^T \Sigma_X \mathbf{a} = \text{Var}(\mathbf{a}^T \mathbf{X}) \geq 0$$

- Covariance matrix  $\Sigma_X$  positive semidefinite because

$$\mathbf{a}^T \Sigma_X \mathbf{a} = \text{Var}(\mathbf{a}^T \mathbf{X}) \geq 0$$

- $\Sigma_X$  would be positive definite if  $\mathbf{a}^T \Sigma_X \mathbf{a} > 0, \forall \mathbf{a} \neq 0$



- Covariance matrix  $\Sigma_X$  positive semidefinite because

$$\mathbf{a}^T \Sigma_X \mathbf{a} = \text{Var}(\mathbf{a}^T \mathbf{X}) \geq 0$$

- $\Sigma_X$  would be positive definite if  $\mathbf{a}^T \Sigma_X \mathbf{a} > 0, \forall \mathbf{a} \neq 0$
- It would fail to be positive definite if  $\text{Var}(\mathbf{a}^T \mathbf{X}) = 0$  for some nonzero  $\mathbf{a}$ .

- Covariance matrix  $\Sigma_X$  positive semidefinite because

$$\mathbf{a}^T \Sigma_X \mathbf{a} = \text{Var}(\mathbf{a}^T \mathbf{X}) \geq 0$$

- $\Sigma_X$  would be positive definite if  $\mathbf{a}^T \Sigma_X \mathbf{a} > 0$ ,  $\forall \mathbf{a} \neq 0$
- It would fail to be positive definite if  $\text{Var}(\mathbf{a}^T \mathbf{X}) = 0$  for some nonzero  $\mathbf{a}$ .
- $\text{Var}(Z) = E[(Z - EZ)^2] = 0$  implies  $Z = EZ$ , a constant.

- Covariance matrix  $\Sigma_X$  positive semidefinite because

$$\mathbf{a}^T \Sigma_X \mathbf{a} = \text{Var}(\mathbf{a}^T \mathbf{X}) \geq 0$$

- $\Sigma_X$  would be positive definite if  $\mathbf{a}^T \Sigma_X \mathbf{a} > 0$ ,  $\forall \mathbf{a} \neq 0$
- It would fail to be positive definite if  $\text{Var}(\mathbf{a}^T \mathbf{X}) = 0$  for some nonzero  $\mathbf{a}$ .
- $\text{Var}(Z) = E[(Z - EZ)^2] = 0$  implies  $Z = EZ$ , a constant.
- Hence,  $\Sigma_X$  fails to be positive definite only if there is a non-zero linear combination of  $X_i$ 's that is a constant.

- ▶ Covariance matrix is a real symmetric positive semidefinite matrix

- ▶ Covariance matrix is a real symmetric positive semidefinite matrix
- ▶ It have real and non-negative eigen values.

- ▶ Covariance matrix is a real symmetric positive semidefinite matrix
- ▶ It have real and non-negative eigen values.
- ▶ It would have  $n$  linearly independent eigen vectors.

- ▶ Covariance matrix is a real symmetric positive semidefinite matrix
- ▶ It have real and non-negative eigen values.
- ▶ It would have  $n$  linearly independent eigen vectors.
- ▶ These also have some interesting roles.

- ▶ Covariance matrix is a real symmetric positive semidefinite matrix
- ▶ It have real and non-negative eigen values.
- ▶ It would have  $n$  linearly independent eigen vectors.
- ▶ These also have some interesting roles.
- ▶ We consider one simple example.



- ▶ Let  $Y = \mathbf{a}^T \mathbf{X}$  and assume  $\|\mathbf{a}\| = 1$

- ▶ Let  $Y = \mathbf{a}^T \mathbf{X}$  and assume  $\|\mathbf{a}\| = 1$
- ▶  $Y$  is projection of  $\mathbf{X}$  along the direction  $\mathbf{a}$ .

- ▶ Let  $Y = \mathbf{a}^T \mathbf{X}$  and assume  $\|\mathbf{a}\| = 1$
- ▶  $Y$  is projection of  $\mathbf{X}$  along the direction  $\mathbf{a}$ .
- ▶ Suppose we want to find a direction along which variance is maximized

- ▶ Let  $Y = \mathbf{a}^T \mathbf{X}$  and assume  $\|\mathbf{a}\| = 1$
- ▶  $Y$  is projection of  $\mathbf{X}$  along the direction  $\mathbf{a}$ .
- ▶ Suppose we want to find a direction along which variance is maximized
- ▶ We want to maximize  $\mathbf{a}^T \Sigma_X \mathbf{a}$  subject to  $\mathbf{a}^T \mathbf{a} = 1$

- ▶ Let  $Y = \mathbf{a}^T \mathbf{X}$  and assume  $\|\mathbf{a}\| = 1$
- ▶  $Y$  is projection of  $\mathbf{X}$  along the direction  $\mathbf{a}$ .
- ▶ Suppose we want to find a direction along which variance is maximized
- ▶ We want to maximize  $\mathbf{a}^T \Sigma_X \mathbf{a}$  subject to  $\mathbf{a}^T \mathbf{a} = 1$
- ▶ The lagrangian is  $\mathbf{a}^T \Sigma_X \mathbf{a} + \eta(1 - \mathbf{a}^T \mathbf{a})$

- ▶ Let  $Y = \mathbf{a}^T \mathbf{X}$  and assume  $\|\mathbf{a}\| = 1$
- ▶  $Y$  is projection of  $\mathbf{X}$  along the direction  $\mathbf{a}$ .
- ▶ Suppose we want to find a direction along which variance is maximized
- ▶ We want to maximize  $\mathbf{a}^T \Sigma_X \mathbf{a}$  subject to  $\mathbf{a}^T \mathbf{a} = 1$
- ▶ The lagrangian is  $\mathbf{a}^T \Sigma_X \mathbf{a} + \eta(1 - \mathbf{a}^T \mathbf{a})$
- ▶ Equating the gradient to zero, we get

$$\Sigma_X \mathbf{a} = \eta \mathbf{a}$$

- ▶ Let  $Y = \mathbf{a}^T \mathbf{X}$  and assume  $\|\mathbf{a}\| = 1$
- ▶  $Y$  is projection of  $\mathbf{X}$  along the direction  $\mathbf{a}$ .
- ▶ Suppose we want to find a direction along which variance is maximized
- ▶ We want to maximize  $\mathbf{a}^T \Sigma_X \mathbf{a}$  subject to  $\mathbf{a}^T \mathbf{a} = 1$
- ▶ The lagrangian is  $\mathbf{a}^T \Sigma_X \mathbf{a} + \eta(1 - \mathbf{a}^T \mathbf{a})$
- ▶ Equating the gradient to zero, we get

$$\Sigma_X \mathbf{a} = \eta \mathbf{a}$$

- ▶ So,  $\mathbf{a}$  should be an eigen vector (with eigen value  $\eta$ ).

- ▶ Let  $Y = \mathbf{a}^T \mathbf{X}$  and assume  $\|\mathbf{a}\| = 1$
- ▶  $Y$  is projection of  $\mathbf{X}$  along the direction  $\mathbf{a}$ .
- ▶ Suppose we want to find a direction along which variance is maximized
- ▶ We want to maximize  $\mathbf{a}^T \Sigma_X \mathbf{a}$  subject to  $\mathbf{a}^T \mathbf{a} = 1$
- ▶ The lagrangian is  $\mathbf{a}^T \Sigma_X \mathbf{a} + \eta(1 - \mathbf{a}^T \mathbf{a})$
- ▶ Equating the gradient to zero, we get

$$\Sigma_X \mathbf{a} = \eta \mathbf{a}$$

- ▶ So,  $\mathbf{a}$  should be an eigen vector (with eigen value  $\eta$ ).
- ▶ Then the variance would be  $\mathbf{a}^T \Sigma_X \mathbf{a} = \eta \mathbf{a}^T \mathbf{a} = \eta$



- ▶ Let  $Y = \mathbf{a}^T \mathbf{X}$  and assume  $\|\mathbf{a}\| = 1$
- ▶  $Y$  is projection of  $\mathbf{X}$  along the direction  $\mathbf{a}$ .
- ▶ Suppose we want to find a direction along which variance is maximized
- ▶ We want to maximize  $\mathbf{a}^T \Sigma_X \mathbf{a}$  subject to  $\mathbf{a}^T \mathbf{a} = 1$
- ▶ The lagrangian is  $\mathbf{a}^T \Sigma_X \mathbf{a} + \eta(1 - \mathbf{a}^T \mathbf{a})$
- ▶ Equating the gradient to zero, we get

$$\Sigma_X \mathbf{a} = \eta \mathbf{a}$$

- ▶ So,  $\mathbf{a}$  should be an eigen vector (with eigen value  $\eta$ ).
- ▶ Then the variance would be  $\mathbf{a}^T \Sigma_X \mathbf{a} = \eta \mathbf{a}^T \mathbf{a} = \eta$
- ▶ Hence the direction is the eigen vector corresponding to the highest eigen value.