

# Recap: Convergence in Probability

- ▶ A sequence of random variables,  $X_n$ , is said to **converge in probability** to a random variable  $X_0$  is

$$\lim_{n \rightarrow \infty} P[|X_n - X_0| > \epsilon] = 0, \forall \epsilon > 0$$

# Recap: Convergence in Probability

- ▶ A sequence of random variables,  $X_n$ , is said to **converge in probability** to a random variable  $X_0$  is

$$\lim_{n \rightarrow \infty} P[|X_n - X_0| > \epsilon] = 0, \forall \epsilon > 0$$

This is denoted as  $X_n \xrightarrow{P} X_0$

# Recap: Convergence in Probability

- ▶ A sequence of random variables,  $X_n$ , is said to **converge in probability** to a random variable  $X_0$  is

$$\lim_{n \rightarrow \infty} P[|X_n - X_0| > \epsilon] = 0, \forall \epsilon > 0$$

This is denoted as  $X_n \xrightarrow{P} X_0$

- ▶ By the definition of limit, the above means

$$\forall \delta > 0, \exists N < \infty, \text{ s.t. } P[|X_n - X_0| > \epsilon] < \delta, \forall n > N$$

# Recap: Convergence in Probability

- ▶ A sequence of random variables,  $X_n$ , is said to **converge in probability** to a random variable  $X_0$  is

$$\lim_{n \rightarrow \infty} P[|X_n - X_0| > \epsilon] = 0, \forall \epsilon > 0$$

This is denoted as  $X_n \xrightarrow{P} X_0$

- ▶ By the definition of limit, the above means

$$\forall \delta > 0, \exists N < \infty, \text{ s.t. } P[|X_n - X_0| > \epsilon] < \delta, \forall n > N$$

- ▶ We only need marginal distributions of individual  $X_n$  to decide whether a sequence converges to a constant in probability

# Recap: Weak Law of large numbers

- ▶  $X_i$  are iid,  $EX_i = \mu$ ,  $\text{Var}(X_i) = \sigma^2$ ,  $S_n = \sum_{i=1}^n X_i$

# Recap: Weak Law of large numbers

- $X_i$  are iid,  $EX_i = \mu$ ,  $\text{Var}(X_i) = \sigma^2$ ,  $S_n = \sum_{i=1}^n X_i$

$$E\left[\frac{S_n}{n}\right] = \mu; \quad \text{and} \quad \text{Var}\left(\frac{S_n}{n}\right) = \frac{\sigma^2}{n}$$

## Recap: Weak Law of large numbers

- $X_i$  are iid,  $EX_i = \mu$ ,  $\text{Var}(X_i) = \sigma^2$ ,  $S_n = \sum_{i=1}^n X_i$

$$E\left[\frac{S_n}{n}\right] = \mu; \quad \text{and} \quad \text{Var}\left(\frac{S_n}{n}\right) = \frac{\sigma^2}{n}$$

Weak law of large numbers states

$$\frac{S_n}{n} \xrightarrow{P} \mu$$

## Recap: almost sure convergence

- ▶ A sequence of random variables,  $X_n$ , is said to converge **almost surely** or **with probability one** to  $X$  if

$$P(\{\omega : X_n(\omega) \rightarrow X(\omega)\}) = 1$$



## Recap: almost sure convergence

- ▶ A sequence of random variables,  $X_n$ , is said to converge **almost surely** or **with probability one** to  $X$  if

$$P(\{\omega : X_n(\omega) \rightarrow X(\omega)\}) = 1$$

or equivalently

$$P(\{\omega : X_n(\omega) \not\rightarrow X(\omega)\}) = 0$$

## Recap: almost sure convergence

- ▶ A sequence of random variables,  $X_n$ , is said to converge **almost surely** or **with probability one** to  $X$  if

$$P(\{\omega : X_n(\omega) \rightarrow X(\omega)\}) = 1$$

or equivalently

$$P(\{\omega : X_n(\omega) \not\rightarrow X(\omega)\}) = 0$$

- ▶ Denoted as  $X_n \xrightarrow{a.s.} X$  or  $X_n \xrightarrow{w.p.1} X$  or  $X_n \rightarrow X_0$  (*w.p.1*)

## Recap: almost sure convergence

- ▶ A sequence of random variables,  $X_n$ , is said to converge **almost surely** or **with probability one** to  $X$  if

$$P(\{\omega : X_n(\omega) \rightarrow X(\omega)\}) = 1$$

or equivalently

$$P(\{\omega : X_n(\omega) \not\rightarrow X(\omega)\}) = 0$$

- ▶ Denoted as  $X_n \xrightarrow{a.s.} X$  or  $X_n \xrightarrow{w.p.1} X$  or  $X_n \rightarrow X$  (*w.p.1*)
- ▶ We can also write it as

$$P[X_n \rightarrow X] = 1$$

# Recap

- ▶ The sequence  $X_n$  converges to  $X$  almost surely iff

$$P\left(\bigcap_{N=1}^{\infty} \bigcup_{k=0}^{\infty} [|X_{N+k} - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

# Recap

- ▶ The sequence  $X_n$  converges to  $X$  almost surely iff

$$P(\cap_{N=1}^{\infty} \cup_{k=0}^{\infty} [|X_{N+k} - X| \geq \epsilon]) = 0, \quad \forall \epsilon > 0$$

Same as

$$P(\cap_{N=1}^{\infty} \cup_{k=N}^{\infty} [|X_k - X| \geq \epsilon]) = 0, \quad \forall \epsilon > 0$$

# Recap

- ▶ The sequence  $X_n$  converges to  $X$  almost surely iff

$$P\left(\bigcap_{N=1}^{\infty} \bigcup_{k=0}^{\infty} [|X_{N+k} - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

Same as

$$P\left(\bigcap_{N=1}^{\infty} \bigcup_{k=N}^{\infty} [|X_k - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

- ▶ Equivalently

$$\lim_{n \rightarrow \infty} P\left(\bigcup_{k=n}^{\infty} [|X_k - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

# Recap

- ▶ The sequence  $X_n$  converges to  $X$  almost surely iff

$$P\left(\bigcap_{N=1}^{\infty} \bigcup_{k=0}^{\infty} [|X_{N+k} - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

Same as

$$P\left(\bigcap_{N=1}^{\infty} \bigcup_{k=N}^{\infty} [|X_k - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

- ▶ Equivalently

$$\lim_{n \rightarrow \infty} P\left(\bigcup_{k=n}^{\infty} [|X_k - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

- ▶  $X_n \xrightarrow{P} X$  iff

$$\lim_{n \rightarrow \infty} P[|X_n - X| > \epsilon] = 0, \quad \forall \epsilon > 0$$

# Recap

- ▶ The sequence  $X_n$  converges to  $X$  almost surely iff

$$P\left(\bigcap_{N=1}^{\infty} \bigcup_{k=0}^{\infty} [|X_{N+k} - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

Same as

$$P\left(\bigcap_{N=1}^{\infty} \bigcup_{k=N}^{\infty} [|X_k - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

- ▶ Equivalently

$$\lim_{n \rightarrow \infty} P\left(\bigcup_{k=n}^{\infty} [|X_k - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

- ▶  $X_n \xrightarrow{P} X$  iff

$$\lim_{n \rightarrow \infty} P[|X_n - X| > \epsilon] = 0, \quad \forall \epsilon > 0$$

$$X_n \xrightarrow{a.s.} X \quad \Rightarrow \quad X_n \xrightarrow{P} X$$



## Recap: $\limsup$ and $\liminf$

- ▶ Let  $A_1, A_2, \dots$  be a sequence of events.

# Recap: $\limsup$ and $\liminf$

- ▶ Let  $A_1, A_2, \dots$  be a sequence of events.
- ▶ We define

$$\limsup A_n \triangleq \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k$$

# Recap: $\limsup$ and $\liminf$

- ▶ Let  $A_1, A_2, \dots$  be a sequence of events.
- ▶ We define

$$\begin{aligned}\limsup A_n &\triangleq \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \\ \liminf A_n &\triangleq \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k\end{aligned}$$

# Recap: lim sup and lim inf

- ▶ Let  $A_1, A_2, \dots$  be a sequence of events.
- ▶ We define

$$\begin{aligned}\limsup A_n &\triangleq \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \\ \liminf A_n &\triangleq \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k\end{aligned}$$

- ▶ If  $\limsup A_n = \liminf A_n$  then that is  $\lim A_n$ .  
Otherwise the sequence does not have a limit

# Recap: $\limsup$ and $\liminf$

- ▶ Let  $A_1, A_2, \dots$  be a sequence of events.
- ▶ We define

$$\begin{aligned}\limsup A_n &\triangleq \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \\ \liminf A_n &\triangleq \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k\end{aligned}$$

- ▶ If  $\limsup A_n = \liminf A_n$  then that is  $\lim A_n$ .  
Otherwise the sequence does not have a limit
- ▶  $\limsup A_n$  and  $\liminf A_n$  are events

## Recap: $\limsup$ and $\liminf$

- ▶ Let  $A_1, A_2, \dots$  be a sequence of events.
- ▶ We define

$$\begin{aligned}\limsup A_n &\triangleq \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \\ \liminf A_n &\triangleq \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k\end{aligned}$$

- ▶ If  $\limsup A_n = \liminf A_n$  then that is  $\lim A_n$ .  
Otherwise the sequence does not have a limit
- ▶  $\limsup A_n$  and  $\liminf A_n$  are events
- ▶  $\liminf A_n \subset \limsup A_n$

# Recap

►  $X_n \xrightarrow{a.s.} X$  iff

$$P\left(\bigcap_{N=1}^{\infty} \bigcup_{k=N}^{\infty} [|X_k - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

# Recap

►  $X_n \xrightarrow{a.s.} X$  iff

$$P\left(\bigcap_{N=1}^{\infty} \bigcup_{k=N}^{\infty} [|X_k - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

► Let  $A_k^\epsilon = [|X_k - X| \geq \epsilon]$ .



# Recap

- ▶  $X_n \xrightarrow{a.s.} X$  iff

$$P\left(\bigcap_{N=1}^{\infty} \bigcup_{k=N}^{\infty} [|X_k - X| \geq \epsilon]\right) = 0, \quad \forall \epsilon > 0$$

- ▶ Let  $A_k^\epsilon = [|X_k - X| \geq \epsilon]$ .
- ▶ Hence,  $X_n \xrightarrow{a.s.} X$  iff

$$P\left(\limsup A_n^\epsilon\right) = 0, \quad \forall \epsilon > 0$$

# Recall: Borel-Cantelli Lemma

# Recall: Borel-Cantelli Lemma

- ▶ **Borel-Cantelli lemma:** Given sequence of events,  $A_1, A_2, \dots$

# Recall: Borel-Cantelli Lemma

- ▶ **Borel-Cantelli lemma:** Given sequence of events,  $A_1, A_2, \dots$ 
  1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$

# Recall: Borel-Cantelli Lemma

- **Borel-Cantelli lemma:** Given sequence of events,  $A_1, A_2, \dots$
1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$
  2. If  $\sum_{i=1}^{\infty} P(A_i) = \infty$  and  $A_i$  are independent,  $P(\limsup A_n) = 1$

# Proof of Borel-Cantelli Lemma

# Proof of Borel-Cantelli Lemma

- ▶ **Borel-Cantelli lemma:** Given  $A_1, A_2, \dots$

# Proof of Borel-Cantelli Lemma

► **Borel-Cantelli lemma:** Given  $A_1, A_2, \dots$

1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$



# Proof of Borel-Cantelli Lemma

► **Borel-Cantelli lemma:** Given  $A_1, A_2, \dots$

1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$
2. If  $\sum_{i=1}^{\infty} P(A_i) = \infty$  and  $A_i$  are independent,  $P(\limsup A_n) = 1$

# Proof of Borel-Cantelli Lemma

- **Borel-Cantelli lemma:** Given  $A_1, A_2, \dots$
1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$
  2. If  $\sum_{i=1}^{\infty} P(A_i) = \infty$  and  $A_i$  are independent,  $P(\limsup A_n) = 1$

**Proof:**

# Proof of Borel-Cantelli Lemma

- ▶ **Borel-Cantelli lemma:** Given  $A_1, A_2, \dots$ 
  1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$
  2. If  $\sum_{i=1}^{\infty} P(A_i) = \infty$  and  $A_i$  are independent,  $P(\limsup A_n) = 1$

## Proof:

- ▶ We will first show:  $P(\cup_{i=n}^{\infty} A_i) \leq \sum_{i=n}^{\infty} P(A_i), \forall n$

# Proof of Borel-Cantelli Lemma

- ▶ **Borel-Cantelli lemma:** Given  $A_1, A_2, \dots$ 
  1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$
  2. If  $\sum_{i=1}^{\infty} P(A_i) = \infty$  and  $A_i$  are independent,  $P(\limsup A_n) = 1$

## Proof:

- ▶ We will first show:  $P(\cup_{i=n}^{\infty} A_i) \leq \sum_{i=n}^{\infty} P(A_i), \forall n$
- ▶ We have the result:  $P(\cup_{i=n}^N A_i) \leq \sum_{i=n}^N P(A_i), n \leq N$

# Proof of Borel-Cantelli Lemma

- ▶ **Borel-Cantelli lemma:** Given  $A_1, A_2, \dots$ 
  1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$
  2. If  $\sum_{i=1}^{\infty} P(A_i) = \infty$  and  $A_i$  are independent,  $P(\limsup A_n) = 1$

## Proof:

- ▶ We will first show:  $P(\cup_{i=n}^{\infty} A_i) \leq \sum_{i=n}^{\infty} P(A_i)$ ,  $\forall n$
- ▶ We have the result:  $P(\cup_{i=n}^N A_i) \leq \sum_{i=n}^N P(A_i)$ ,  $n \leq N$
- ▶ For any  $n$ , let  $B_N = \cup_{i=n}^N A_i$ . Then  $B_N \subset B_{N+1}$ .

# Proof of Borel-Cantelli Lemma

- ▶ **Borel-Cantelli lemma:** Given  $A_1, A_2, \dots$ 
  1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$
  2. If  $\sum_{i=1}^{\infty} P(A_i) = \infty$  and  $A_i$  are independent,  $P(\limsup A_n) = 1$

## Proof:

- ▶ We will first show:  $P(\cup_{i=n}^{\infty} A_i) \leq \sum_{i=n}^{\infty} P(A_i)$ ,  $\forall n$
- ▶ We have the result:  $P(\cup_{i=n}^N A_i) \leq \sum_{i=n}^N P(A_i)$ ,  $n \leq N$
- ▶ For any  $n$ , let  $B_N = \cup_{i=n}^N A_i$ . Then  $B_N \subset B_{N+1}$ .
- ▶  $\lim_{N \rightarrow \infty} B_N = \cup_{k=n}^{\infty} A_k$

# Proof of Borel-Cantelli Lemma

- ▶ **Borel-Cantelli lemma:** Given  $A_1, A_2, \dots$ 
  1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$
  2. If  $\sum_{i=1}^{\infty} P(A_i) = \infty$  and  $A_i$  are independent,  $P(\limsup A_n) = 1$

## Proof:

- ▶ We will first show:  $P(\cup_{i=n}^{\infty} A_i) \leq \sum_{i=n}^{\infty} P(A_i)$ ,  $\forall n$
- ▶ We have the result:  $P(\cup_{i=n}^N A_i) \leq \sum_{i=n}^N P(A_i)$ ,  $n \leq N$
- ▶ For any  $n$ , let  $B_N = \cup_{i=n}^N A_i$ . Then  $B_N \subset B_{N+1}$ .
- ▶  $\lim_{N \rightarrow \infty} B_N = \cup_{k=n}^{\infty} A_k$

$$P(\cup_{i=n}^{\infty} A_i) = P(\lim_{N \rightarrow \infty} \cup_{i=n}^N A_i)$$

# Proof of Borel-Cantelli Lemma

- ▶ **Borel-Cantelli lemma:** Given  $A_1, A_2, \dots$ 
  1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$
  2. If  $\sum_{i=1}^{\infty} P(A_i) = \infty$  and  $A_i$  are independent,  $P(\limsup A_n) = 1$

## Proof:

- ▶ We will first show:  $P(\cup_{i=n}^{\infty} A_i) \leq \sum_{i=n}^{\infty} P(A_i)$ ,  $\forall n$
- ▶ We have the result:  $P(\cup_{i=n}^N A_i) \leq \sum_{i=n}^N P(A_i)$ ,  $n \leq N$
- ▶ For any  $n$ , let  $B_N = \cup_{i=n}^N A_i$ . Then  $B_N \subset B_{N+1}$ .
- ▶  $\lim_{N \rightarrow \infty} B_N = \cup_{k=n}^{\infty} A_k$

$$P(\cup_{i=n}^{\infty} A_i) = P(\lim_{N \rightarrow \infty} \cup_{i=n}^N A_i) = \lim_{N \rightarrow \infty} P(\cup_{i=n}^N A_i)$$



# Proof of Borel-Cantelli Lemma

- ▶ **Borel-Cantelli lemma:** Given  $A_1, A_2, \dots$ 
  1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$
  2. If  $\sum_{i=1}^{\infty} P(A_i) = \infty$  and  $A_i$  are independent,  $P(\limsup A_n) = 1$

## Proof:

- ▶ We will first show:  $P(\cup_{i=n}^{\infty} A_i) \leq \sum_{i=n}^{\infty} P(A_i)$ ,  $\forall n$
- ▶ We have the result:  $P(\cup_{i=n}^N A_i) \leq \sum_{i=n}^N P(A_i)$ ,  $n \leq N$
- ▶ For any  $n$ , let  $B_N = \cup_{i=n}^N A_i$ . Then  $B_N \subset B_{N+1}$ .
- ▶  $\lim_{N \rightarrow \infty} B_N = \cup_{k=n}^{\infty} A_k$

$$\begin{aligned} P(\cup_{i=n}^{\infty} A_i) &= P(\lim_{N \rightarrow \infty} \cup_{i=n}^N A_i) = \lim_{N \rightarrow \infty} P(\cup_{i=n}^N A_i) \\ &\leq \lim_{N \rightarrow \infty} \sum_{i=n}^N P(A_i) \end{aligned}$$

# Proof of Borel-Cantelli Lemma

- ▶ **Borel-Cantelli lemma:** Given  $A_1, A_2, \dots$ 
  1. If  $\sum_{i=1}^{\infty} P(A_i) < \infty$ , then,  $P(\limsup A_n) = 0$
  2. If  $\sum_{i=1}^{\infty} P(A_i) = \infty$  and  $A_i$  are independent,  $P(\limsup A_n) = 1$

## Proof:

- ▶ We will first show:  $P(\cup_{i=n}^{\infty} A_i) \leq \sum_{i=n}^{\infty} P(A_i)$ ,  $\forall n$
- ▶ We have the result:  $P(\cup_{i=n}^N A_i) \leq \sum_{i=n}^N P(A_i)$ ,  $n \leq N$
- ▶ For any  $n$ , let  $B_N = \cup_{i=n}^N A_i$ . Then  $B_N \subset B_{N+1}$ .
- ▶  $\lim_{N \rightarrow \infty} B_N = \cup_{k=n}^{\infty} A_k$

$$\begin{aligned} P(\cup_{i=n}^{\infty} A_i) &= P(\lim_{N \rightarrow \infty} \cup_{i=n}^N A_i) = \lim_{N \rightarrow \infty} P(\cup_{i=n}^N A_i) \\ &\leq \lim_{N \rightarrow \infty} \sum_{i=n}^N P(A_i) = \sum_{i=n}^{\infty} P(A_i) \end{aligned}$$

► By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

- By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

- Let  $\sum_{k=1}^{\infty} P(A_k) = C < \infty$

- By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

- Let  $\sum_{k=1}^{\infty} P(A_k) = C < \infty$
- It means given any  $\epsilon > 0$ ,  $\exists n$

$$\left| \sum_{k=1}^n P(A_k) - C \right| < \epsilon$$

- By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

- Let  $\sum_{k=1}^{\infty} P(A_k) = C < \infty$
- It means given any  $\epsilon > 0$ ,  $\exists n$

$$\left| \sum_{k=1}^n P(A_k) - C \right| < \epsilon \Rightarrow \left| \sum_{k=n}^{\infty} P(A_k) \right| < \epsilon$$

- By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

- Let  $\sum_{k=1}^{\infty} P(A_k) = C < \infty$
- It means given any  $\epsilon > 0$ ,  $\exists n$

$$\left| \sum_{k=1}^n P(A_k) - C \right| < \epsilon \Rightarrow \left| \sum_{k=n}^{\infty} P(A_k) \right| < \epsilon$$

- This implies

$$\lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

- By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

- Let  $\sum_{k=1}^{\infty} P(A_k) = C < \infty$

- It means given any  $\epsilon > 0$ ,  $\exists n$

$$\left| \sum_{k=1}^n P(A_k) - C \right| < \epsilon \Rightarrow \left| \sum_{k=n}^{\infty} P(A_k) \right| < \epsilon$$

- This implies

$$\lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

- Similarly we have

$$\sum_{k=1}^{\infty} P(A_k) = \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = \infty$$



► By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

► By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

$$0 \leq P(\limsup A_n) = P(\cap_{n=1}^{\infty} \cup_{k=n}^{\infty} A_k)$$

► By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

$$\begin{aligned} 0 \leq P(\limsup A_n) &= P\left(\bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k\right) \\ &= P\left(\lim_{n \rightarrow \infty} \bigcup_{k=n}^{\infty} A_k\right) \end{aligned}$$

► By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

$$\begin{aligned} 0 \leq P(\limsup A_n) &= P(\cap_{n=1}^{\infty} \cup_{k=n}^{\infty} A_k) \\ &= P\left(\lim_{n \rightarrow \infty} \cup_{k=n}^{\infty} A_k\right) \\ &= \lim_{n \rightarrow \infty} P(\cup_{k=n}^{\infty} A_k) \end{aligned}$$

► By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

$$\begin{aligned} 0 \leq P(\limsup A_n) &= P(\cap_{n=1}^{\infty} \cup_{k=n}^{\infty} A_k) \\ &= P\left(\lim_{n \rightarrow \infty} \cup_{k=n}^{\infty} A_k\right) \\ &= \lim_{n \rightarrow \infty} P(\cup_{k=n}^{\infty} A_k) \\ &\leq \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) \end{aligned}$$

► By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

$$\begin{aligned} 0 \leq P(\limsup A_n) &= P(\cap_{n=1}^{\infty} \cup_{k=n}^{\infty} A_k) \\ &= P\left(\lim_{n \rightarrow \infty} \cup_{k=n}^{\infty} A_k\right) \\ &= \lim_{n \rightarrow \infty} P(\cup_{k=n}^{\infty} A_k) \\ &\leq \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) \\ &= 0, \quad \text{if } \sum_{k=1}^{\infty} P(A_k) < \infty \end{aligned}$$

► By definition,

$$\sum_{k=1}^{\infty} P(A_k) < \infty, \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = 0$$

$$\begin{aligned} 0 \leq P(\limsup A_n) &= P(\cap_{n=1}^{\infty} \cup_{k=n}^{\infty} A_k) \\ &= P\left(\lim_{n \rightarrow \infty} \cup_{k=n}^{\infty} A_k\right) \\ &= \lim_{n \rightarrow \infty} P(\cup_{k=n}^{\infty} A_k) \\ &\leq \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) \\ &= 0, \quad \text{if } \sum_{k=1}^{\infty} P(A_k) < \infty \end{aligned}$$

► This completes proof of first part of Borel-Cantelli lemma

- For the second part of the lemma:



- For the second part of the lemma:

$$P(\limsup A_n) = P(\cap_{n=1}^{\infty} \cup_{k=n}^{\infty} A_k)$$

- For the second part of the lemma:

$$\begin{aligned} P(\limsup A_n) &= P(\cap_{n=1}^{\infty} \cup_{k=n}^{\infty} A_k) \\ &= P\left(\lim_{n \rightarrow \infty} \cup_{k=n}^{\infty} A_k\right) \end{aligned}$$

► For the second part of the lemma:

$$\begin{aligned} P(\limsup A_n) &= P(\cap_{n=1}^{\infty} \cup_{k=n}^{\infty} A_k) \\ &= P\left(\lim_{n \rightarrow \infty} \cup_{k=n}^{\infty} A_k\right) \\ &= \lim_{n \rightarrow \infty} P(\cup_{k=n}^{\infty} A_k) \end{aligned}$$

► For the second part of the lemma:

$$\begin{aligned} P(\limsup A_n) &= P(\cap_{n=1}^{\infty} \cup_{k=n}^{\infty} A_k) \\ &= P\left(\lim_{n \rightarrow \infty} \cup_{k=n}^{\infty} A_k\right) \\ &= \lim_{n \rightarrow \infty} P(\cup_{k=n}^{\infty} A_k) \\ &= \lim_{n \rightarrow \infty} (1 - P(\cap_{k=n}^{\infty} A_k^c)) \end{aligned}$$

► For the second part of the lemma:

$$\begin{aligned} P(\limsup A_n) &= P(\cap_{n=1}^{\infty} \cup_{k=n}^{\infty} A_k) \\ &= P\left(\lim_{n \rightarrow \infty} \cup_{k=n}^{\infty} A_k\right) \\ &= \lim_{n \rightarrow \infty} P(\cup_{k=n}^{\infty} A_k) \\ &= \lim_{n \rightarrow \infty} (1 - P(\cap_{k=n}^{\infty} A_k^c)) \\ &= \lim_{n \rightarrow \infty} \left(1 - \prod_{k=n}^{\infty} (1 - P(A_k))\right) \end{aligned}$$

► For the second part of the lemma:

$$\begin{aligned}P(\limsup A_n) &= P(\cap_{n=1}^{\infty} \cup_{k=n}^{\infty} A_k) \\&= P\left(\lim_{n \rightarrow \infty} \cup_{k=n}^{\infty} A_k\right) \\&= \lim_{n \rightarrow \infty} P(\cup_{k=n}^{\infty} A_k) \\&= \lim_{n \rightarrow \infty} (1 - P(\cap_{k=n}^{\infty} A_k^c)) \\&= \lim_{n \rightarrow \infty} \left(1 - \prod_{k=n}^{\infty} (1 - P(A_k))\right) \\&\quad \text{because } A_k \text{ are independent}\end{aligned}$$

► For the second part of the lemma:

$$\begin{aligned}P(\limsup A_n) &= P(\cap_{n=1}^{\infty} \cup_{k=n}^{\infty} A_k) \\&= P\left(\lim_{n \rightarrow \infty} \cup_{k=n}^{\infty} A_k\right) \\&= \lim_{n \rightarrow \infty} P(\cup_{k=n}^{\infty} A_k) \\&= \lim_{n \rightarrow \infty} (1 - P(\cap_{k=n}^{\infty} A_k^c)) \\&= \lim_{n \rightarrow \infty} \left(1 - \prod_{k=n}^{\infty} (1 - P(A_k))\right) \\&\quad \text{because } A_k \text{ are independent} \\&= 1 - \lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} (1 - P(A_k))\end{aligned}$$

- ▶ We can compute that limit as follows



► We can compute that limit as follows

$$\lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} (1 - P(A_k)) \leq \lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} e^{-P(A_k)}, \quad \text{since } 1 - x \leq e^{-x}$$

► We can compute that limit as follows

$$\begin{aligned}\lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} (1 - P(A_k)) &\leq \lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} e^{-P(A_k)}, \quad \text{since } 1 - x \leq e^{-x} \\ &= \lim_{n \rightarrow \infty} e^{-\sum_{k=n}^{\infty} P(A_k)}\end{aligned}$$

► We can compute that limit as follows

$$\begin{aligned}\lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} (1 - P(A_k)) &\leq \lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} e^{-P(A_k)}, \quad \text{since } 1 - x \leq e^{-x} \\ &= \lim_{n \rightarrow \infty} e^{-\sum_{k=n}^{\infty} P(A_k)} \\ &= 0\end{aligned}$$

► We can compute that limit as follows

$$\begin{aligned}\lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} (1 - P(A_k)) &\leq \lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} e^{-P(A_k)}, \quad \text{since } 1 - x \leq e^{-x} \\ &= \lim_{n \rightarrow \infty} e^{-\sum_{k=n}^{\infty} P(A_k)} \\ &= 0\end{aligned}$$

because

$$\sum_{k=1}^{\infty} P(A_k) = \infty \quad \Rightarrow \quad \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = \infty$$

- We can compute that limit as follows

$$\begin{aligned}\lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} (1 - P(A_k)) &\leq \lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} e^{-P(A_k)}, \quad \text{since } 1 - x \leq e^{-x} \\ &= \lim_{n \rightarrow \infty} e^{-\sum_{k=n}^{\infty} P(A_k)} \\ &= 0\end{aligned}$$

because

$$\sum_{k=1}^{\infty} P(A_k) = \infty \quad \Rightarrow \quad \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = \infty$$

- This finally gives us

$$P(\limsup A_n) = 1 - \lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} (1 - P(A_k)) = 1$$

- We can compute that limit as follows

$$\begin{aligned}\lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} (1 - P(A_k)) &\leq \lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} e^{-P(A_k)}, \quad \text{since } 1 - x \leq e^{-x} \\ &= \lim_{n \rightarrow \infty} e^{-\sum_{k=n}^{\infty} P(A_k)} \\ &= 0\end{aligned}$$

because

$$\sum_{k=1}^{\infty} P(A_k) = \infty \Rightarrow \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} P(A_k) = \infty$$

- This finally gives us

$$P(\limsup A_n) = 1 - \lim_{n \rightarrow \infty} \prod_{k=n}^{\infty} (1 - P(A_k)) = 1$$

- That completes the proof.

- ▶ Given a sequence  $X_n$  we want to know whether it converges to  $X$

- ▶ Given a sequence  $X_n$  we want to know whether it converges to  $X$
- ▶ Let  $A_k^\epsilon = [|X_k - X| \geq \epsilon]$



- ▶ Given a sequence  $X_n$  we want to know whether it converges to  $X$
- ▶ Let  $A_k^\epsilon = [|X_k - X| \geq \epsilon]$
- ▶  $X_n \xrightarrow{P} X$  if

$$\lim_{k \rightarrow \infty} P[|X_k - X| \geq \epsilon] = 0$$

- ▶ Given a sequence  $X_n$  we want to know whether it converges to  $X$
- ▶ Let  $A_k^\epsilon = [|X_k - X| \geq \epsilon]$
- ▶  $X_n \xrightarrow{P} X$  if

$$\lim_{k \rightarrow \infty} P[|X_k - X| \geq \epsilon] = 0 \quad \text{same as} \quad \lim_{k \rightarrow \infty} P(A_k^\epsilon) = 0, \quad \forall \epsilon > 0$$

- ▶ Given a sequence  $X_n$  we want to know whether it converges to  $X$
- ▶ Let  $A_k^\epsilon = [|X_k - X| \geq \epsilon]$
- ▶  $X_n \xrightarrow{P} X$  if

$$\lim_{k \rightarrow \infty} P[|X_k - X| \geq \epsilon] = 0 \quad \text{same as} \quad \lim_{k \rightarrow \infty} P(A_k^\epsilon) = 0, \quad \forall \epsilon > 0$$

- ▶ By Borel-Cantelli lemma, if  $\sum_{k=1}^{\infty} P(A_k^\epsilon) < \infty$ ,

$$\sum_{k=1}^{\infty} P(A_k^\epsilon) < \infty$$

- ▶ Given a sequence  $X_n$  we want to know whether it converges to  $X$
- ▶ Let  $A_k^\epsilon = [|X_k - X| \geq \epsilon]$
- ▶  $X_n \xrightarrow{P} X$  if

$$\lim_{k \rightarrow \infty} P[|X_k - X| \geq \epsilon] = 0 \quad \text{same as} \quad \lim_{k \rightarrow \infty} P(A_k^\epsilon) = 0, \quad \forall \epsilon > 0$$

- ▶ By Borel-Cantelli lemma, if  $\sum_{k=1}^{\infty} P(A_k^\epsilon) < \infty$ , then  $P(\limsup A_k^\epsilon) = 0$ , which implies  $X_k \xrightarrow{a.s.} X$ .

$$\sum_{k=1}^{\infty} P(A_k^\epsilon) < \infty \Rightarrow P(\limsup A_k^\epsilon) = 0 \Rightarrow X_k \xrightarrow{a.s.} X$$

- ▶ Given a sequence  $X_n$  we want to know whether it converges to  $X$
- ▶ Let  $A_k^\epsilon = [|X_k - X| \geq \epsilon]$
- ▶  $X_n \xrightarrow{P} X$  if

$$\lim_{k \rightarrow \infty} P[|X_k - X| \geq \epsilon] = 0 \quad \text{same as} \quad \lim_{k \rightarrow \infty} P(A_k^\epsilon) = 0, \quad \forall \epsilon > 0$$

- ▶ By Borel-Cantelli lemma, if  $\sum_{k=1}^{\infty} P(A_k^\epsilon) < \infty$ , then  $P(\limsup A_k^\epsilon) = 0 \Rightarrow X_k \xrightarrow{a.s.} X$

$$\sum_{k=1}^{\infty} P(A_k^\epsilon) < \infty \Rightarrow P(\limsup A_k^\epsilon) = 0 \Rightarrow X_k \xrightarrow{a.s.} X$$

If  $X_k$  are ind

$$\sum_{k=1}^{\infty} P(A_k^\epsilon) = \infty \Rightarrow P(\limsup A_k^\epsilon) = 1 \Rightarrow X_k \not\xrightarrow{a.s.} X$$

## Recall Example

- Consider a sequence  $X_n$  with

$$P[X_n = 0] = 1 - \frac{1}{n}; \quad P[X_n = 1] = \frac{1}{n}$$

## Recall Example

- ▶ Consider a sequence  $X_n$  with

$$P[X_n = 0] = 1 - \frac{1}{n}; \quad P[X_n = 1] = \frac{1}{n}$$

- ▶ Since  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow \infty$ , we conclude  $X_n \xrightarrow{P} 0$ .

## Recall Example

- ▶ Consider a sequence  $X_n$  with

$$P[X_n = 0] = 1 - \frac{1}{n}; \quad P[X_n = 1] = \frac{1}{n}$$

- ▶ Since  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow \infty$ , we conclude  $X_n \xrightarrow{P} 0$ .
- ▶ But since,  $\sum_n \frac{1}{n} = \infty$ , Borel-Cantelli is not useful



## Recall Example

- ▶ Consider a sequence  $X_n$  with

$$P[X_n = 0] = 1 - \frac{1}{n}; \quad P[X_n = 1] = \frac{1}{n}$$

- ▶ Since  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow \infty$ , we conclude  $X_n \xrightarrow{P} 0$ .
- ▶ But since,  $\sum_n \frac{1}{n} = \infty$ , Borel-Cantelli is not useful
- ▶ We saw one example of such  $X_n$  with  $X_k \xrightarrow{a.s.} 0$

## Recall Example

- ▶ Consider a sequence  $X_n$  with

$$P[X_n = 0] = 1 - \frac{1}{n}; \quad P[X_n = 1] = \frac{1}{n}$$

- ▶ Since  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow \infty$ , we conclude  $X_n \xrightarrow{P} 0$ .
- ▶ But since,  $\sum_n \frac{1}{n} = \infty$ , Borel-Cantelli is not useful
- ▶ We saw one example of such  $X_n$  with  $X_k \xrightarrow{a.s.} 0$
- ▶ But, if  $X_n$  are independent, then by Borel-Cantelli lemma, they do not converge

## Recall Example

- ▶ Consider a sequence  $X_n$  with

$$P[X_n = 0] = 1 - \frac{1}{n}; \quad P[X_n = 1] = \frac{1}{n}$$

- ▶ Since  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow \infty$ , we conclude  $X_n \xrightarrow{P} 0$ .
- ▶ But since,  $\sum_n \frac{1}{n} = \infty$ , Borel-Cantelli is not useful
- ▶ We saw one example of such  $X_n$  with  $X_k \xrightarrow{a.s.} 0$
- ▶ But, if  $X_n$  are independent, then by Borel-Cantelli lemma, they do not converge
- ▶ Almost sure convergence is stronger than convergence in probability

## Recall Example

- ▶ Consider a sequence  $X_n$  with

$$P[X_n = 0] = 1 - \frac{1}{n}; \quad P[X_n = 1] = \frac{1}{n}$$

- ▶ Since  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow \infty$ , we conclude  $X_n \xrightarrow{P} 0$ .
- ▶ But since,  $\sum_n \frac{1}{n} = \infty$ , Borel-Cantelli is not useful
- ▶ We saw one example of such  $X_n$  with  $X_k \xrightarrow{a.s.} 0$
- ▶ But, if  $X_n$  are independent, then by Borel-Cantelli lemma, they do not converge
- ▶ Almost sure convergence is stronger than convergence in probability
- ▶ Convergence (to a constant) in probability depends only on distribution of individual  $X_n$ .

## Recall Example

- ▶ Consider a sequence  $X_n$  with

$$P[X_n = 0] = 1 - \frac{1}{n}; \quad P[X_n = 1] = \frac{1}{n}$$

- ▶ Since  $\frac{1}{n} \rightarrow 0$  as  $n \rightarrow \infty$ , we conclude  $X_n \xrightarrow{P} 0$ .
- ▶ But since,  $\sum_n \frac{1}{n} = \infty$ , Borel-Cantelli is not useful
- ▶ We saw one example of such  $X_n$  with  $X_k \xrightarrow{a.s.} 0$
- ▶ But, if  $X_n$  are independent, then by Borel-Cantelli lemma, they do not converge
- ▶ Almost sure convergence is stronger than convergence in probability
- ▶ Convergence (to a constant) in probability depends only on distribution of individual  $X_n$ .
- ▶ Convergence almost surely depends on the joint distribution

# Strong Law of Large Numbers

- ▶ Let  $X_n$  be iid,  $EX_n = \mu$ ,  $\text{Var}(X_n) = \sigma^2$ ,  $S_n = \sum_{i=1}^n X_i$

# Strong Law of Large Numbers

- ▶ Let  $X_n$  be iid,  $EX_n = \mu$ ,  $\text{Var}(X_n) = \sigma^2$ ,  $S_n = \sum_{i=1}^n X_i$
- ▶ We saw weak law of large numbers:

$$\frac{S_n}{n} \xrightarrow{P} \mu$$

# Strong Law of Large Numbers

- ▶ Let  $X_n$  be iid,  $EX_n = \mu$ ,  $\text{Var}(X_n) = \sigma^2$ ,  $S_n = \sum_{i=1}^n X_i$
- ▶ We saw weak law of large numbers:

$$\frac{S_n}{n} \xrightarrow{P} \mu$$

- ▶ Strong law of large numbers says:

$$\frac{S_n}{n} \xrightarrow{\text{a.s.}} \mu$$



# Strong Law of Large Numbers

- ▶ Let  $X_n$  be iid,  $EX_n = \mu$ ,  $\text{Var}(X_n) = \sigma^2$ ,  $S_n = \sum_{i=1}^n X_i$
- ▶ We saw weak law of large numbers:

$$\frac{S_n}{n} \xrightarrow{P} \mu$$

- ▶ Strong law of large numbers says:

$$\frac{S_n}{n} \xrightarrow{\text{a.s.}} \mu$$

- ▶ This is true for **all** random variables

# Strong Law of Large Numbers

- ▶ Let  $X_n$  be iid,  $EX_n = \mu$ ,  $\text{Var}(X_n) = \sigma^2$ ,  $S_n = \sum_{i=1}^n X_i$
- ▶ We saw weak law of large numbers:

$$\frac{S_n}{n} \xrightarrow{P} \mu$$

- ▶ Strong law of large numbers says:

$$\frac{S_n}{n} \xrightarrow{a.s.} \mu$$

- ▶ This is true for **all** random variables
- ▶ We prove only a restricted version

► Let  $A_n^\epsilon = [|\frac{S_n}{n} - \mu| > \epsilon]$

- ▶ Let  $A_n^\epsilon = [|\frac{S_n}{n} - \mu| > \epsilon]$
- ▶ As we saw, by Chebyshev inequality

$$P \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right] \leq \frac{\sigma^2}{n\epsilon^2}$$

- ▶ Let  $A_n^\epsilon = \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right]$
- ▶ As we saw, by Chebyshev inequality

$$P \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right] \leq \frac{\sigma^2}{n\epsilon^2}$$

- ▶ This shows  $P(A_n^\epsilon) \rightarrow 0$  and thus we get weak law

- ▶ Let  $A_n^\epsilon = \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right]$
- ▶ As we saw, by Chebyshev inequality

$$P \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right] \leq \frac{\sigma^2}{n\epsilon^2}$$

- ▶ This shows  $P(A_n^\epsilon) \rightarrow 0$  and thus we get weak law
- ▶ To prove strong law using Borel-Cantelli lemma, we need  $\sum P(A_n^\epsilon) < \infty$

- ▶ Let  $A_n^\epsilon = \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right]$
- ▶ As we saw, by Chebyshev inequality

$$P \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right] \leq \frac{\sigma^2}{n\epsilon^2}$$

- ▶ This shows  $P(A_n^\epsilon) \rightarrow 0$  and thus we get weak law
- ▶ To prove strong law using Borel-Cantelli lemma, we need  $\sum P(A_n^\epsilon) < \infty$
- ▶ Since  $\sum_n \frac{\sigma^2}{n\epsilon^2} = \infty$ , the Chebyshev bound is not useful

- ▶ Let  $A_n^\epsilon = [|\frac{S_n}{n} - \mu| > \epsilon]$
- ▶ As we saw, by Chebyshev inequality

$$P \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right] \leq \frac{\sigma^2}{n\epsilon^2}$$

- ▶ This shows  $P(A_n^\epsilon) \rightarrow 0$  and thus we get weak law
- ▶ To prove strong law using Borel-Cantelli lemma, we need  $\sum P(A_n^\epsilon) < \infty$
- ▶ Since  $\sum_n \frac{\sigma^2}{n\epsilon^2} = \infty$ , the Chebyshev bound is not useful
- ▶ We need a bound:  $P[|\frac{S_n}{n} - \mu|] \leq c_n$  such that  $\sum_n c_n < \infty$ .



- ▶ Let us assume  $X_i$  have finite fourth moment

► Let us assume  $X_i$  have finite fourth moment

$$\left( \sum_{i=1}^n (X_i - \mu) \right)^4 =$$

► Let us assume  $X_i$  have finite fourth moment

$$\left( \sum_{i=1}^n (X_i - \mu) \right)^4 = \sum_{i=1}^n (X_i - \mu)^4 + \sum_i \sum_{j>i} \frac{4!}{2!2!} (X_i - \mu)^2 (X_j - \mu)^2 + T$$

► Let us assume  $X_i$  have finite fourth moment

$$\left( \sum_{i=1}^n (X_i - \mu) \right)^4 = \sum_{i=1}^n (X_i - \mu)^4 + \sum_i \sum_{j>i} \frac{4!}{2!2!} (X_i - \mu)^2 (X_j - \mu)^2 + T$$

Where  $T$  represent a number of terms such that every term in it contains a factor like  $(X_i - \mu)$

► Let us assume  $X_i$  have finite fourth moment

$$\left( \sum_{i=1}^n (X_i - \mu) \right)^4 = \sum_{i=1}^n (X_i - \mu)^4 + \sum_i \sum_{j>i} \frac{4!}{2!2!} (X_i - \mu)^2 (X_j - \mu)^2 + T$$

Where  $T$  represent a number of terms such that every term in it contains a factor like  $(X_i - \mu)$

Note that  $E[(X_i - \mu)(X_j - \mu)^3] = 0$  etc. because  $X_i$  are independent.

- Let us assume  $X_i$  have finite fourth moment

$$\left( \sum_{i=1}^n (X_i - \mu) \right)^4 = \sum_{i=1}^n (X_i - \mu)^4 + \sum_i \sum_{j>i} \frac{4!}{2!2!} (X_i - \mu)^2 (X_j - \mu)^2 + T$$

Where  $T$  represent a number of terms such that every term in it contains a factor like  $(X_i - \mu)$

Note that  $E[(X_i - \mu)(X_j - \mu)^3] = 0$  etc. because  $X_i$  are independent.

- Hence we get

$$E \left[ \left( \sum_{i=1}^n (X_i - \mu) \right)^4 \right] = nE[(X_i - \mu)^4] + 3n(n-1)\sigma^4 \leq C'n^2$$

- ▶ Now we can get, using Markov inequality

- Now we can get, using Markov inequality

$$P \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right] = P [|S_n - n\mu| > n\epsilon]$$



- Now we can get, using Markov inequality

$$\begin{aligned} P \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right] &= P [|S_n - n\mu| > n\epsilon] \\ &= P \left[ \left| \sum_{i=1}^n (X_i - \mu) \right| > n\epsilon \right] \end{aligned}$$

- Now we can get, using Markov inequality

$$\begin{aligned} P \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right] &= P [|S_n - n\mu| > n\epsilon] \\ &= P \left[ \left| \sum_{i=1}^n (X_i - \mu) \right| > n\epsilon \right] \\ &\leq \frac{E (\sum_{i=1}^n (X_i - \mu))^4}{(n\epsilon)^4} \end{aligned}$$

- Now we can get, using Markov inequality

$$\begin{aligned} P \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right] &= P [|S_n - n\mu| > n\epsilon] \\ &= P \left[ \left| \sum_{i=1}^n (X_i - \mu) \right| > n\epsilon \right] \\ &\leq \frac{E (\sum_{i=1}^n (X_i - \mu))^4}{(n\epsilon)^4} \\ &\leq \frac{C' n^2}{n^4 \epsilon^4} = \frac{C}{n^2} \end{aligned}$$

- Now we can get, using Markov inequality

$$\begin{aligned} P \left[ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right] &= P [|S_n - n\mu| > n\epsilon] \\ &= P \left[ \left| \sum_{i=1}^n (X_i - \mu) \right| > n\epsilon \right] \\ &\leq \frac{E (\sum_{i=1}^n (X_i - \mu))^4}{(n\epsilon)^4} \\ &\leq \frac{C' n^2}{n^4 \epsilon^4} = \frac{C}{n^2} \end{aligned}$$

- Since  $\sum_n \frac{C}{n^2} < \infty$ , we get  $\frac{S_n}{n} \xrightarrow{a.s.} \mu$

- Strong law of large numbers says

$$\frac{S_n}{n} \xrightarrow{a.s.} \mu \quad \text{where } S_n = \sum_{i=1}^n X_i, \quad X_i \text{ iid, } EX_i = \mu$$

- ▶ Strong law of large numbers says

$$\frac{S_n}{n} \xrightarrow{a.s.} \mu \quad \text{where } S_n = \sum_{i=1}^n X_i, \quad X_i \text{ iid}, \quad EX_i = \mu$$

- ▶ We proved it assuming finite fourth moment of  $X_i$ .

- ▶ Strong law of large numbers says

$$\frac{S_n}{n} \xrightarrow{a.s.} \mu \quad \text{where} \quad S_n = \sum_{i=1}^n X_i, \quad X_i \text{ iid}, \quad EX_i = \mu$$

- ▶ We proved it assuming finite fourth moment of  $X_i$ .
- ▶ This is only for illustration

- ▶ Strong law of large numbers says

$$\frac{S_n}{n} \xrightarrow{a.s.} \mu \quad \text{where } S_n = \sum_{i=1}^n X_i, \quad X_i \text{ iid, } EX_i = \mu$$

- ▶ We proved it assuming finite fourth moment of  $X_i$ .
- ▶ This is only for illustration
- ▶ Strong law holds without any such assumptions on moments



- ▶ Strong law of large numbers says

$$\frac{S_n}{n} \xrightarrow{a.s.} \mu \quad \text{where } S_n = \sum_{i=1}^n X_i, \quad X_i \text{ iid}, \quad EX_i = \mu$$

- ▶ We proved it assuming finite fourth moment of  $X_i$ .
- ▶ This is only for illustration
- ▶ Strong law holds without any such assumptions on moments
- ▶ Strong law of large numbers says that sample mean converges to the expectation with probability one.

# Convergence in probability Vs Almost sure convergence

►  $X_n \xrightarrow{P} X_0$   
 $\Leftrightarrow \lim_{n \rightarrow \infty} P[|X_n - X_0| > \epsilon] = 0, \forall \epsilon > 0$

# Convergence in probability Vs Almost sure convergence

- ▶  $X_n \xrightarrow{P} X_0$   
 $\Leftrightarrow \lim_{n \rightarrow \infty} P[|X_n - X_0| > \epsilon] = 0, \forall \epsilon > 0$
- ▶  $X_n \xrightarrow{a.s.} X_0$   
 $\Leftrightarrow \lim_{n \rightarrow \infty} P(\cup_{k=n}^{\infty} [|X_k - X_0| \geq \epsilon]) = 0, \forall \epsilon > 0$

# Convergence in probability Vs Almost sure convergence

- ▶  $X_n \xrightarrow{P} X_0$   
 $\Leftrightarrow \lim_{n \rightarrow \infty} P[|X_n - X_0| > \epsilon] = 0, \forall \epsilon > 0$
- ▶  $X_n \xrightarrow{a.s.} X_0$   
 $\Leftrightarrow \lim_{n \rightarrow \infty} P(\cup_{k=n}^{\infty} [|X_k - X_0| \geq \epsilon]) = 0, \forall \epsilon > 0$
- ▶ One can intuitively see why convergence almost surely is a much stronger notion of convergence.

# Example

- $\Omega = [0, 1]$ . Sequence of binary random variables:  
 $X_{nk}$ ,  $k = 1, \dots, n$ ,  $n = 1, 2, \dots$ , defined by

$$X_{nk}(\omega) = 1 \text{ iff } \frac{k-1}{n} \leq \omega < \frac{k}{n}, \quad 1 \leq k \leq n, n = 1, 2, \dots$$

That is, the sequence is  $X_{11}, X_{21}, X_{22}, X_{31}, X_{32}, X_{33}, \dots$ .

# Example

- ▶  $\Omega = [0, 1]$ . Sequence of binary random variables:  
 $X_{nk}$ ,  $k = 1, \dots, n$ ,  $n = 1, 2, \dots$ , defined by

$$X_{nk}(\omega) = 1 \text{ iff } \frac{k-1}{n} \leq \omega < \frac{k}{n}, \quad 1 \leq k \leq n, n = 1, 2, \dots$$

That is, the sequence is  $X_{11}, X_{21}, X_{22}, X_{31}, X_{32}, X_{33}, \dots$ .

- ▶ One can show the sequence converges to zero in probability.

# Example

- ▶  $\Omega = [0, 1]$ . Sequence of binary random variables:  
 $X_{nk}$ ,  $k = 1, \dots, n$ ,  $n = 1, 2, \dots$ , defined by

$$X_{nk}(\omega) = 1 \text{ iff } \frac{k-1}{n} \leq \omega < \frac{k}{n}, \quad 1 \leq k \leq n, n = 1, 2, \dots$$

That is, the sequence is  $X_{11}, X_{21}, X_{22}, X_{31}, X_{32}, X_{33}, \dots$ .

- ▶ One can show the sequence converges to zero in probability.
- ▶ But,  $P[X_{nk} \rightarrow 0] = 0!$

$$X_{nk}(\omega) = 1 \quad \text{iff} \quad \frac{k-1}{n} \leq \omega < \frac{k}{n}, \quad 1 \leq k \leq n$$



## Convergence in $r^{th}$ mean

- We say that a sequence  $X_n$  converges in  $r^{th}$  mean to  $X$  if  $E[|X_n|^r] < \infty$ ,  $\forall n$ ,  $E[|X|^r] < \infty$  and

$$E[|X_n - X|^r] \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

## Convergence in $r^{th}$ mean

- ▶ We say that a sequence  $X_n$  converges in  $r^{th}$  mean to  $X$  if  $E[|X_n|^r] < \infty$ ,  $\forall n$ ,  $E[|X|^r] < \infty$  and

$$E[|X_n - X|^r] \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

- ▶ Denoted as  $X_n \xrightarrow{r} X$

## Convergence in $r^{th}$ mean

- ▶ We say that a sequence  $X_n$  converges in  $r^{th}$  mean to  $X$  if  $E[|X_n|^r] < \infty, \forall n, E[|X|^r] < \infty$  and

$$E[|X_n - X|^r] \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

- ▶ Denoted as  $X_n \xrightarrow{r} X$
- ▶ Consider our old example of binary random variables

$$P[X_n = 1] = \frac{1}{n} \quad P[X_n = 0] = 1 - \frac{1}{n}$$

## Convergence in $r^{th}$ mean

- ▶ We say that a sequence  $X_n$  converges in  $r^{th}$  mean to  $X$  if  $E[|X_n|^r] < \infty, \forall n, E[|X|^r] < \infty$  and

$$E[|X_n - X|^r] \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

- ▶ Denoted as  $X_n \xrightarrow{r} X$
- ▶ Consider our old example of binary random variables

$$P[X_n = 1] = \frac{1}{n} \quad P[X_n = 0] = 1 - \frac{1}{n}$$

- ▶ All moments of  $X_n$  are finite and we have

## Convergence in $r^{th}$ mean

- ▶ We say that a sequence  $X_n$  converges in  $r^{th}$  mean to  $X$  if  $E[|X_n|^r] < \infty$ ,  $\forall n$ ,  $E[|X|^r] < \infty$  and

$$E[|X_n - X|^r] \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

- ▶ Denoted as  $X_n \xrightarrow{r} X$
- ▶ Consider our old example of binary random variables

$$P[X_n = 1] = \frac{1}{n} \quad P[X_n = 0] = 1 - \frac{1}{n}$$

- ▶ All moments of  $X_n$  are finite and we have

$$E[|X_n - 0|^2] = \frac{1}{n} \rightarrow 0$$

## Convergence in $r^{th}$ mean

- ▶ We say that a sequence  $X_n$  converges in  $r^{th}$  mean to  $X$  if  $E[|X_n|^r] < \infty$ ,  $\forall n$ ,  $E[|X|^r] < \infty$  and

$$E[|X_n - X|^r] \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

- ▶ Denoted as  $X_n \xrightarrow{r} X$
- ▶ Consider our old example of binary random variables

$$P[X_n = 1] = \frac{1}{n} \quad P[X_n = 0] = 1 - \frac{1}{n}$$

- ▶ All moments of  $X_n$  are finite and we have

$$E[|X_n - 0|^2] = \frac{1}{n} \rightarrow 0$$

- ▶ Hence  $X_n \xrightarrow{2} 0$ .

## Convergence in $r^{th}$ mean

- ▶ We say that a sequence  $X_n$  converges in  $r^{th}$  mean to  $X$  if  $E[|X_n|^r] < \infty, \forall n, E[|X|^r] < \infty$  and

$$E[|X_n - X|^r] \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

- ▶ Denoted as  $X_n \xrightarrow{r} X$
- ▶ Consider our old example of binary random variables

$$P[X_n = 1] = \frac{1}{n} \quad P[X_n = 0] = 1 - \frac{1}{n}$$

- ▶ All moments of  $X_n$  are finite and we have

$$E[|X_n - 0|^2] = \frac{1}{n} \rightarrow 0$$

- ▶ Hence  $X_n \xrightarrow{2} 0$ .
- ▶ In this example  $X_n$  converges in  $r^{th}$  mean for all  $r$

► Suppose  $X_n \xrightarrow{r} X$ . Then, by Markov inequality

$$P[|X_n - X| > \epsilon] \leq \frac{E[|X_n - X|^r]}{\epsilon^r} \rightarrow 0$$



- Suppose  $X_n \xrightarrow{r} X$ . Then, by Markov inequality

$$P[|X_n - X| > \epsilon] \leq \frac{E[|X_n - X|^r]}{\epsilon^r} \rightarrow 0$$

- Hence

$$X_n \xrightarrow{r} X \Rightarrow X_n \xrightarrow{P} X$$

- Suppose  $X_n \xrightarrow{r} X$ . Then, by Markov inequality

$$P[|X_n - X| > \epsilon] \leq \frac{E[|X_n - X|^r]}{\epsilon^r} \rightarrow 0$$

- Hence

$$X_n \xrightarrow{r} X \Rightarrow X_n \xrightarrow{P} X$$

- Consider the sequence  $X_n$  with

$$P[X_n = 0] = 1 - \frac{1}{n}; \quad P[X_n = e^n] = \frac{1}{n}$$

- Suppose  $X_n \xrightarrow{r} X$ . Then, by Markov inequality

$$P[|X_n - X| > \epsilon] \leq \frac{E[|X_n - X|^r]}{\epsilon^r} \rightarrow 0$$

- Hence

$$X_n \xrightarrow{r} X \Rightarrow X_n \xrightarrow{P} X$$

- Consider the sequence  $X_n$  with

$$P[X_n = 0] = 1 - \frac{1}{n}; \quad P[X_n = e^n] = \frac{1}{n}$$

We have  $P[|X_n - 0| > \epsilon] = \frac{1}{n} \Rightarrow X_n \xrightarrow{P} 0$

- ▶ Suppose  $X_n \xrightarrow{r} X$ . Then, by Markov inequality

$$P[|X_n - X| > \epsilon] \leq \frac{E[|X_n - X|^r]}{\epsilon^r} \rightarrow 0$$

- ▶ Hence

$$X_n \xrightarrow{r} X \Rightarrow X_n \xrightarrow{P} X$$

- ▶ Consider the sequence  $X_n$  with

$$P[X_n = 0] = 1 - \frac{1}{n}; \quad P[X_n = e^n] = \frac{1}{n}$$

We have  $P[|X_n - 0| > \epsilon] = \frac{1}{n} \Rightarrow X_n \xrightarrow{P} 0$

- ▶ But,  $E|X_n - 0|^r = \frac{e^{rn}}{n}$  and hence  $X_n$  does not converge in  $r^{th}$  mean.

- ▶ In general, neither of convergence almost surely and in  $r^{th}$  mean imply the other.

- ▶ In general, neither of convergence almost surely and in  $r^{th}$  mean imply the other.
- ▶ We can generate counter examples for this easily.

- ▶ In general, neither of convergence almost surely and in  $r^{th}$  mean imply the other.
- ▶ We can generate counter examples for this easily.
- ▶ However, if all  $X_n$  take values in a bounded interval, then almost sure convergence implies  $r^{th}$  mean convergence

- Consider sequence  $X_n$  where  $X_n$  are independent with

$$P[X_n = 0] = 1 - a_n; \quad P[X_n = c_n] = a_n$$



- Consider sequence  $X_n$  where  $X_n$  are independent with

$$P[X_n = 0] = 1 - a_n; \quad P[X_n = c_n] = a_n$$

- ▶ Consider sequence  $X_n$  where  $X_n$  are independent with

$$P[X_n = 0] = 1 - a_n; \quad P[X_n = c_n] = a_n$$

- ▶ Assume  $a_n \rightarrow 0$  so that  $X_n \xrightarrow{P} 0$

- ▶ Consider sequence  $X_n$  where  $X_n$  are independent with

$$P[X_n = 0] = 1 - a_n; \quad P[X_n = c_n] = a_n$$

- ▶ Assume  $a_n \rightarrow 0$  so that  $X_n \xrightarrow{P} 0$
- ▶ By Borel-Cantelli lemma

$$X_n \xrightarrow{a.s.} 0 \quad \Leftrightarrow \quad \sum_n a_n < \infty$$

- ▶ Consider sequence  $X_n$  where  $X_n$  are independent with

$$P[X_n = 0] = 1 - a_n; \quad P[X_n = c_n] = a_n$$

- ▶ Assume  $a_n \rightarrow 0$  so that  $X_n \xrightarrow{P} 0$
- ▶ By Borel-Cantelli lemma

$$X_n \xrightarrow{a.s.} 0 \quad \Leftrightarrow \quad \sum_n a_n < \infty$$

- ▶ For convergence in  $r^{th}$  mean we need

$$E[|X_n - 0|^r] = (c_n)^r a_n \rightarrow 0$$

- ▶ Consider sequence  $X_n$  where  $X_n$  are independent with

$$P[X_n = 0] = 1 - a_n; \quad P[X_n = c_n] = a_n$$

- ▶ Assume  $a_n \rightarrow 0$  so that  $X_n \xrightarrow{P} 0$
- ▶ By Borel-Cantelli lemma

$$X_n \xrightarrow{a.s.} 0 \quad \Leftrightarrow \quad \sum_n a_n < \infty$$

- ▶ For convergence in  $r^{th}$  mean we need

$$E[|X_n - 0|^r] = (c_n)^r a_n \rightarrow 0$$

- ▶ Take  $a_n = \frac{1}{n}$  and  $c_n = 1$ . Then  $X_n \xrightarrow{r} 0$  but the sequence does not converge almost surely.

- Consider sequence  $X_n$  where  $X_n$  are independent with

$$P[X_n = 0] = 1 - a_n; \quad P[X_n = c_n] = a_n$$

- Assume  $a_n \rightarrow 0$  so that  $X_n \xrightarrow{P} 0$
- By Borel-Cantelli lemma

$$X_n \xrightarrow{a.s.} 0 \quad \Leftrightarrow \quad \sum_n a_n < \infty$$

- For convergence in  $r^{th}$  mean we need

$$E[|X_n - 0|^r] = (c_n)^r a_n \rightarrow 0$$

- Take  $a_n = \frac{1}{n}$  and  $c_n = 1$ . Then  $X_n \xrightarrow{r} 0$  but the sequence does not converge almost surely.
- Take  $a_n = \frac{1}{n^2}$  and  $c_n = e^n$ . Then  $X_n \xrightarrow{a.s.} 0$  but the sequence does not converge in  $r^{th}$  mean for any  $r$ .

► Let  $X_n \xrightarrow{r} X$ . Then

- Let  $X_n \xrightarrow{r} X$ . Then
1.  $E[|X_n|^r] \rightarrow E[|X|^r]$



- Let  $X_n \xrightarrow{r} X$ . Then
1.  $E[|X_n|^r] \rightarrow E[|X|^r]$
  2.  $X_n \xrightarrow{s} X, \forall s < r$

- ▶ Let  $X_n \xrightarrow{r} X$ . Then
  1.  $E[|X_n|^r] \rightarrow E[|X|^r]$
  2.  $X_n \xrightarrow{s} X, \forall s < r$

- ▶ The proofs are straight-forward but we omit the proofs

# Convergence in distribution

- ▶ Let  $F_n$  be the df of  $X_n$ ,  $n = 1, 2, \dots$ . Let  $X$  be a rv with df  $F$ .

# Convergence in distribution

- ▶ Let  $F_n$  be the df of  $X_n$ ,  $n = 1, 2, \dots$ . Let  $X$  be a rv with df  $F$ .
- ▶ Sequence  $X_n$  is said to converge to  $X$  **in distribution** if

$$F_n(x) \rightarrow F(x), \quad \forall x \text{ where } F \text{ is continuous}$$

# Convergence in distribution

- ▶ Let  $F_n$  be the df of  $X_n$ ,  $n = 1, 2, \dots$ . Let  $X$  be a rv with df  $F$ .
- ▶ Sequence  $X_n$  is said to converge to  $X$  **in distribution** if

$$F_n(x) \rightarrow F(x), \quad \forall x \text{ where } F \text{ is continuous}$$

- ▶ We denote this as

$$X_n \xrightarrow{d} X, \quad \text{or} \quad X_n \xrightarrow{L} X, \quad \text{or} \quad F_n \xrightarrow{w} F$$

# Convergence in distribution

- ▶ Let  $F_n$  be the df of  $X_n$ ,  $n = 1, 2, \dots$ . Let  $X$  be a rv with df  $F$ .
- ▶ Sequence  $X_n$  is said to converge to  $X$  **in distribution** if

$$F_n(x) \rightarrow F(x), \quad \forall x \text{ where } F \text{ is continuous}$$

- ▶ We denote this as

$$X_n \xrightarrow{d} X, \quad \text{or} \quad X_n \xrightarrow{L} X, \quad \text{or} \quad F_n \xrightarrow{w} F$$

- ▶ This is also known as **convergence in law** or weak convergence

# Convergence in distribution

- ▶ Let  $F_n$  be the df of  $X_n$ ,  $n = 1, 2, \dots$ . Let  $X$  be a rv with df  $F$ .
- ▶ Sequence  $X_n$  is said to converge to  $X$  **in distribution** if

$$F_n(x) \rightarrow F(x), \quad \forall x \text{ where } F \text{ is continuous}$$

- ▶ We denote this as

$$X_n \xrightarrow{d} X, \quad \text{or} \quad X_n \xrightarrow{L} X, \quad \text{or} \quad F_n \xrightarrow{w} F$$

- ▶ This is also known as **convergence in law** or weak convergence
- ▶ Note that here we are essentially talking about convergence of distribution functions.

# Convergence in distribution

- ▶ Let  $F_n$  be the df of  $X_n$ ,  $n = 1, 2, \dots$ . Let  $X$  be a rv with df  $F$ .
- ▶ Sequence  $X_n$  is said to converge to  $X$  **in distribution** if

$$F_n(x) \rightarrow F(x), \quad \forall x \text{ where } F \text{ is continuous}$$

- ▶ We denote this as

$$X_n \xrightarrow{d} X, \quad \text{or} \quad X_n \xrightarrow{L} X, \quad \text{or} \quad F_n \xrightarrow{w} F$$

- ▶ This is also known as **convergence in law** or weak convergence
- ▶ Note that here we are essentially talking about convergence of distribution functions.
- ▶ Convergence in probability implies convergence in distribution



# Convergence in distribution

- ▶ Let  $F_n$  be the df of  $X_n$ ,  $n = 1, 2, \dots$ . Let  $X$  be a rv with df  $F$ .

- ▶ Sequence  $X_n$  is said to converge to  $X$  **in distribution** if

$$F_n(x) \rightarrow F(x), \quad \forall x \text{ where } F \text{ is continuous}$$

- ▶ We denote this as

$$X_n \xrightarrow{d} X, \quad \text{or} \quad X_n \xrightarrow{L} X, \quad \text{or} \quad F_n \xrightarrow{w} F$$

- ▶ This is also known as **convergence in law** or weak convergence
- ▶ Note that here we are essentially talking about convergence of distribution functions.
- ▶ Convergence in probability implies convergence in distribution
- ▶ The converse is not true. (e.g., sequence of iid random variables)

# Examples

- ▶  $X_1, X_2, \dots$  be iid; uniform over  $(0, 1)$

# Examples

- ▶  $X_1, X_2, \dots$  be iid; uniform over  $(0, 1)$
- ▶  $N_n = \min(X_1, \dots, X_n)$ ,  $Y_n = nN_n$ .

# Examples

- ▶  $X_1, X_2, \dots$  be iid; uniform over  $(0, 1)$
- ▶  $N_n = \min(X_1, \dots, X_n)$ ,  $Y_n = nN_n$ .  
Does  $Y_n$  converge in distribution?

# Examples

- ▶  $X_1, X_2, \dots$  be iid; uniform over  $(0, 1)$
- ▶  $N_n = \min(X_1, \dots, X_n)$ ,  $Y_n = nN_n$ .  
Does  $Y_n$  converge in distribution?

$$P[N_n > a] = (P[X_i > a])^n$$

# Examples

- ▶  $X_1, X_2, \dots$  be iid; uniform over  $(0, 1)$
- ▶  $N_n = \min(X_1, \dots, X_n)$ ,  $Y_n = nN_n$ .

Does  $Y_n$  converge in distribution?

$$P[N_n > a] = (P[X_i > a])^n = (1 - a)^n, \quad 0 < a < 1$$

# Examples

- ▶  $X_1, X_2, \dots$  be iid; uniform over  $(0, 1)$
- ▶  $N_n = \min(X_1, \dots, X_n)$ ,  $Y_n = nN_n$ .

Does  $Y_n$  converge in distribution?

$$P[N_n > a] = (P[X_i > a])^n = (1 - a)^n, \quad 0 < a < 1$$

$$P[Y_n > y] = P[N_n > y/n]$$

# Examples

- ▶  $X_1, X_2, \dots$  be iid; uniform over  $(0, 1)$
- ▶  $N_n = \min(X_1, \dots, X_n)$ ,  $Y_n = nN_n$ .

Does  $Y_n$  converge in distribution?

$$P[N_n > a] = (P[X_i > a])^n = (1 - a)^n, \quad 0 < a < 1$$

$$P[Y_n > y] = P[N_n > y/n] = \left(1 - \frac{y}{n}\right)^n, \quad \text{if } n > y$$



# Examples

- ▶  $X_1, X_2, \dots$  be iid; uniform over  $(0, 1)$
- ▶  $N_n = \min(X_1, \dots, X_n)$ ,  $Y_n = nN_n$ .

Does  $Y_n$  converge in distribution?

$$P[N_n > a] = (P[X_i > a])^n = (1 - a)^n, \quad 0 < a < 1$$

$$P[Y_n > y] = P[N_n > y/n] = \left(1 - \frac{y}{n}\right)^n, \quad \text{if } n > y$$

- ▶ Hence for any  $y$

$$\lim_{n \rightarrow \infty} P[Y_n > y] = \lim_{n \rightarrow \infty} \left(1 - \frac{y}{n}\right)^n$$

# Examples

- ▶  $X_1, X_2, \dots$  be iid; uniform over  $(0, 1)$
- ▶  $N_n = \min(X_1, \dots, X_n)$ ,  $Y_n = nN_n$ .

Does  $Y_n$  converge in distribution?

$$P[N_n > a] = (P[X_i > a])^n = (1 - a)^n, \quad 0 < a < 1$$

$$P[Y_n > y] = P[N_n > y/n] = \left(1 - \frac{y}{n}\right)^n, \quad \text{if } n > y$$

- ▶ Hence for any  $y$

$$\lim_{n \rightarrow \infty} P[Y_n > y] = \lim_{n \rightarrow \infty} \left(1 - \frac{y}{n}\right)^n = e^{-y}$$

# Examples

- ▶  $X_1, X_2, \dots$  be iid; uniform over  $(0, 1)$
- ▶  $N_n = \min(X_1, \dots, X_n)$ ,  $Y_n = nN_n$ .

Does  $Y_n$  converge in distribution?

$$P[N_n > a] = (P[X_i > a])^n = (1 - a)^n, \quad 0 < a < 1$$

$$P[Y_n > y] = P[N_n > y/n] = \left(1 - \frac{y}{n}\right)^n, \quad \text{if } n > y$$

- ▶ Hence for any  $y$

$$\lim_{n \rightarrow \infty} P[Y_n > y] = \lim_{n \rightarrow \infty} \left(1 - \frac{y}{n}\right)^n = e^{-y}$$

- ▶ The sequence converges in distribution to an exponential rv

►  $X_n \xrightarrow{d} X$   
 $\Leftrightarrow F_n(x) \rightarrow F(x), \quad \forall x$  where  $F$  is continuous

- ▶  $X_n \xrightarrow{d} X$   
 $\Leftrightarrow F_n(x) \rightarrow F(x), \forall x$  where  $F$  is continuous
- ▶ This means that the sequence of functions  $F_n$  converge point-wise and the limit function is a distribution function.

- ▶  $X_n \xrightarrow{d} X$   
 $\Leftrightarrow F_n(x) \rightarrow F(x), \forall x$  where  $F$  is continuous
- ▶ This means that the sequence of functions  $F_n$  converge point-wise and the limit function is a distribution function.
- ▶ In general,  $X_n \xrightarrow{d} X$  does not imply that the pdf's or pmf's converge point-wise to the limit pdf or pmf.

- ▶  $X_n \xrightarrow{d} X$   
 $\Leftrightarrow F_n(x) \rightarrow F(x), \forall x$  where  $F$  is continuous
- ▶ This means that the sequence of functions  $F_n$  converge point-wise and the limit function is a distribution function.
- ▶ In general,  $X_n \xrightarrow{d} X$  does not imply that the pdf's or pmf's converge point-wise to the limit pdf or pmf.
- ▶ However if the sequence of pmf's (or pdf's) converge point-wise and the limit is a pmf (or pdf) then we have  $X_n \xrightarrow{d} X$ .

- ▶  $X_n \xrightarrow{d} X$   
 $\Leftrightarrow F_n(x) \rightarrow F(x), \forall x$  where  $F$  is continuous
- ▶ This means that the sequence of functions  $F_n$  converge point-wise and the limit function is a distribution function.
- ▶ In general,  $X_n \xrightarrow{d} X$  does not imply that the pdf's or pmf's converge point-wise to the limit pdf or pmf.
- ▶ However if the sequence of pmf's (or pdf's) converge point-wise and the limit is a pmf (or pdf) then we have  $X_n \xrightarrow{d} X$ .
- ▶  $X_n \xrightarrow{P} X \Rightarrow X_n \xrightarrow{d} X$



- ▶  $X_n \xrightarrow{d} X$   
 $\Leftrightarrow F_n(x) \rightarrow F(x), \forall x$  where  $F$  is continuous
- ▶ This means that the sequence of functions  $F_n$  converge point-wise and the limit function is a distribution function.
- ▶ In general,  $X_n \xrightarrow{d} X$  does not imply that the pdf's or pmf's converge point-wise to the limit pdf or pmf.
- ▶ However if the sequence of pmf's (or pdf's) converge point-wise and the limit is a pmf (or pdf) then we have  $X_n \xrightarrow{d} X$ .
- ▶  $X_n \xrightarrow{P} X \Rightarrow X_n \xrightarrow{d} X$
- ▶  $X_n \xrightarrow{d} k \Rightarrow X_n \xrightarrow{P} k$ , where  $k$  is a constant