Math for EE

CBSE Class 12 2015

CONTENTS

1 CBSE Class XII Mathematics 3
1.1 Formulae . . . 3
1.1.1 Vectors and three-dimensional geometry . . . 3
1.1.2 Matrices and determinants . . . 3
1.1.3 Differentiation and differential equations . . . 3
1.1.4 Integration and area . . . 4
1.1.5 Functions, relations and operations . . . 4
1.1.6 Probability . . . 5
1.1.7 Linear programming . . . 5
1.1.8 Coordinate geometry and optimisation . . . 5
1.1.9 Engineering methods: choosing the right tool . . . 5
1.2 Worked solutions . . . 6
1.2.1 Vectors and three-dimensional geometry . . . 6
1.2.2 Lines and planes . . . 8
1.2.3 Differential equations . . . 10
1.2.4 Matrices and determinants . . . 12
1.2.5 Integration and area . . . 13
1.2.6 Functions and relations . . . 17
1.2.7 Probability . . . 20
1.2.8 Linear programming . . . 24
1.2.9 Coordinate geometry and optimisation . . . 25

1. CBSE CLASS XII MATHEMATICS

A. Formulae

1) Vectors and three-dimensional geometry: For vectors $\vec{a}, \vec{b} \in \mathbb{R}^3$,

$$\vec{a} \cdot \vec{b} = a_1b_1 + a_2b_2 + a_3b_3, \quad \vec{a} \perp \vec{b} \iff \vec{a} \cdot \vec{b} = 0. \tag{1.1.1.1}$$

$$\vec{a} \times \vec{b} = \begin{pmatrix} a_2b_3 - a_3b_2 \\ a_3b_1 - a_1b_3 \\ a_1b_2 - a_2b_1 \end{pmatrix}. \tag{1.1.1.2}$$

$$\text{unit vector along } \vec{v} = \frac{\vec{v}}{\|\vec{v}\|}, \qquad \|c\vec{v}\| = |c| \, \|\vec{v}\| \, . \tag{1.1.1.3}$$

The vector product is normal to both vectors in the product. Distances and line/plane relations are

$$d(\text{parallel lines}) = \frac{\left\| (\vec{b} - \vec{a}) \times \vec{v} \right\|}{\|\vec{v}\|}. \tag{1.1.2.1}$$

$$d(\text{skew lines}) = \frac{|(\vec{b} - \vec{a}) \cdot (\vec{v}_1 \times \vec{v}_2)|}{\|\vec{v}_1 \times \vec{v}_2\|}. \tag{1.1.2.2}$$

$$\text{line parallel to plane} \iff \vec{d} \cdot \vec{n} = 0. \tag{1.1.2.3}$$

$$\text{plane through } \vec{r}_0 \text{ with normal } \vec{n}: \quad \vec{n} \cdot (\vec{r} - \vec{r}_0) = 0. \tag{1.1.2.4}$$

$$\text{line with point } \vec{A} \text{ and direction } \vec{d}: \quad \vec{r} = \vec{A} + t\vec{d}. \tag{1.1.2.5}$$

2) Matrices and determinants:

$$\det(AB) = \det(A)\det(B), \qquad \det(A^n) = (\det A)^n. \tag{1.2.1.1}$$

$$A^{-1} = \frac{\operatorname{adj}(A)}{\det A}, \qquad \det A \neq 0. \tag{1.2.1.2}$$

$$A\vec{x} = \vec{b} \implies \vec{x} = A^{-1}\vec{b} \quad (A \text{ invertible}). \tag{1.2.1.3}$$

$$[A \mid \vec{b}] \xrightarrow{\text{elementary row operations}} [I \mid \vec{x}] \quad \text{(Gaussian elimination)}. \tag{1.2.1.4}$$

The identity used to factor the determinant in Q36 is

$$a^2 + b^2 + c^2 - ab - bc - ca = \frac{1}{2}\big((a-b)^2 + (b-c)^2 + (c-a)^2\big). \tag{1.2.2.1}$$

3) Differentiation and differential equations:

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt}, \qquad \frac{d^2y}{dx^2} = \frac{d}{dt}(dy/dx) \Big/ \frac{dx}{dt}. \tag{1.3.1.1}$$

$$y' + P(x)y = Q(x), \quad \mu(x) = e^{\int P(x)\,dx}, \quad \frac{d}{dx}(\mu y) = \mu Q. \tag{1.3.1.2}$$

$$y = vx \implies y' = v + xv' \quad \text{(homogeneous first-order ODE)}. \tag{1.3.1.3}$$

$$(1 + (y')^2)^3 = r^2(y'')^2 \quad \text{for a circle family of fixed radius } r. \tag{1.3.1.4}$$

$$\frac{d}{dx}\log|u(x)| = \frac{u'(x)}{u(x)}, \qquad u(x) \neq 0. \tag{1.3.1.5}$$

4) Integration and area:

$$\int \frac{du}{a^2-u^2} = \frac{1}{2a}\log\left|\frac{a+u}{a-u}\right| + C. \tag{1.4.1.1}$$

$$\int \frac{du}{1+u^2} = \tan^{-1} u + C. \tag{1.4.1.2}$$

$$\int \frac{u\,du}{\sqrt{1+u^2}} = \sqrt{1+u^2} + C. \tag{1.4.1.3}$$

$$A = \int_a^b |y_{\text{upper}} - y_{\text{lower}}|\,dx. \tag{1.4.1.4}$$

$$\frac{P(x)}{(x-a)(x^2+b^2)} = Q(x) + \frac{A}{x-a} + \frac{Bx+C}{x^2+b^2} \quad \text{(partial fractions after division).} \tag{1.4.1.5}$$

$$\int u\,dv = uv - \int v\,du. \tag{1.4.1.6}$$

5) Functions, relations and operations: At an interior point, continuity requires matching one-sided limits; differentiability additionally requires matching one-sided derivatives:

$$\lim_{x\to a^-} f(x) = f(a) = \lim_{x\to a^+} f(x), \qquad f'_-(a) = f'_+(a) = f'(a). \tag{1.5.1.1}$$

$$\begin{aligned} &\text{reflexive: } (a,a) \in R, \\ &\text{symmetric: } (a,b) \in R \Rightarrow (b,a) \in R, \\ &\text{transitive: } (a,b),(b,c) \in R \Rightarrow (a,c) \in R. \end{aligned} \tag{1.5.1.2}$$

$$\text{For componentwise addition on } \mathbb{R}^2: \quad e = (0,0), \quad (a,b)^{-1} = (-a,-b). \tag{1.5.1.3}$$

$$a * b = (a+b) \bmod n, \quad 0 \text{ is the identity}, \quad (-a) \bmod n \text{ is the additive inverse.} \tag{1.5.1.4}$$

6) Probability:

$$P(A \mid E) = \frac{P(A \cap E)}{P(E)}. \tag{1.6.1.1}$$

$$P(A \mid E) = \frac{P(E \mid A)P(A)}{P(E)}. \tag{1.6.1.2}$$

$$X \sim \mathrm{Bin}(n,p): \quad P(X=k) = \binom{n}{k} p^k (1-p)^{n-k}. \tag{1.6.1.3}$$

$$G(z) = \sum_j P(D=j) z^j. \tag{1.6.1.4}$$

$$G_5(z) = (0.4z + 0.6z^{-1})^5 \text{ for the five-step walk.}$$

$$E[X] = \sum_x x P(X=x), \quad \mathrm{Var}(X) = E[X^2] - E[X]^2. \tag{1.6.1.5}$$

$$P(X=k) = \frac{\binom{K}{k}\binom{N-K}{n-k}}{\binom{N}{n}}. \tag{1.6.1.6}$$

$$P(E) = \sum_i P(E \mid A_i) P(A_i). \tag{1}$$

(law of total probability). (1.6.1.7)

7) Linear programming:

Evaluate the objective at each feasible corner; an optimum occurs at a feasible vertex. (1.7.1.1)

8) Coordinate geometry and optimisation: The tangent to $x^2 + y^2 = r^2$ at $(r\cos\theta, r\sin\theta)$ is

$$x\cos\theta + y\sin\theta = r. \tag{1.8.1.1}$$

For the ellipse $x^2/a^2 + y^2/b^2 = 1$, a parametrisation is $(a\cos t, b\sin t)$, and triangle area is one half base times perpendicular height:

$$\text{ellipse point } (a\cos t, b\sin t), \quad A_\triangle = \frac{1}{2} bh. \tag{1.8.1.2}$$

9) Engineering methods: choosing the right tool: Matrix methods are used for vector and matrix conditions and for Gaussian elimination. The five-step random walk and the small binomial problem use probability-generating polynomial coefficients where it makes the probability structure clearer. Laplace transforms are useful for suitable linear time-domain differential equations with initial conditions; they are not forced into the nonlinear or variable-coefficient differential equations in this set where they would not simplify the solution.

B. Worked solutions

1) Vectors and three-dimensional geometry:

Q1. Find a vector of magnitude $\sqrt{171}$ perpendicular to both $\vec{a} = \begin{bmatrix} 1 \\ -2 \\ -3 \end{bmatrix}$ and $\vec{b} = \begin{bmatrix} 3 \\ -1 \\ -2 \end{bmatrix}$.

Formula reference: `1.1.1.2, 1.1.1.3`


Solution. A vector perpendicular to both is parallel to their cross product:

$$\vec{a} \times \vec{b} = \begin{bmatrix} 1 \\ -7 \\ 5 \end{bmatrix}, \qquad \left\| \vec{a} \times \vec{b} \right\| = \sqrt{75} = 5\sqrt{3}. \tag{1.2.1.1}$$

Scale this direction to the required magnitude:

$$\vec{d} = \frac{\sqrt{171}}{5\sqrt{3}} \begin{bmatrix} 1 \\ -7 \\ 5 \end{bmatrix} = \boxed{\pm \frac{\sqrt{57}}{5} \begin{bmatrix} 1 \\ -7 \\ 5 \end{bmatrix}}.$$

Q1: the answer vector is normal to the plane spanned by a and b



Code reference – plot and numerical verification.


`codes/Q01.py`

Q5. Let $\vec{a} = \begin{bmatrix} 1 \\ 4 \\ 2 \end{bmatrix}$, $\vec{b} = \begin{bmatrix} 3 \\ -2 \\ 7 \end{bmatrix}$ and $\vec{c} = \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}$. Find $\vec{d}$ perpendicular to both $\vec{a}, \vec{b}$ with $\vec{c} \cdot \vec{d} = 27$.

Formula reference: `1.1.1.1, 1.1.1.2`

Solution. The required direction is $\vec{a} \times \vec{b} = (32, -1, -14)^T$. Set $\vec{d} = t(32, -1, -14)^T$. Since $\vec{c} \cdot (\vec{a} \times \vec{b}) = 9$, the constraint gives $9t = 27$, hence

$$\boxed{\vec{d} = \begin{bmatrix} 96 \\ -3 \\ -42 \end{bmatrix}}.$$



Code reference – plot and numerical verification.

`codes/Q05.py`

Q6. Find the shortest distance between $\vec{r} = (1, 2, 3) + \lambda(2, 3, 4)$ and $\vec{r} = (2, 4, 5) + \mu(4, 6, 8)$.

Formula reference: `1.1.2.1`


Solution. The direction vectors are parallel, since $(4, 6, 8) = 2(2, 3, 4)$. For parallel lines the distance is the perpendicular distance from one point to the other line:

$$d = \frac{\|(B - A) \times \vec{v}\|}{\|\vec{v}\|}, \quad B - A = (1, 2, 2), \quad \vec{v} = (2, 3, 4).$$

Here $(B-A)\times\vec{v}=(2,0,-1)$, so

$$\boxed{d=\frac{\sqrt{5}}{\sqrt{29}}}.$$

Q6: zoomed view of the shortest distance



Code reference – plot and numerical verification.

`codes/Q06.py`

2) Lines and planes:

Q10. Find the plane through the intersection of $2x+y-z=3$ and $5x-3y+4z+9=0$ that is parallel to $\frac{x-1}{2}=\frac{y-3}{-4}=\frac{5-z}{-5}$.

Formula reference: `1.1.2.3, 1.1.2.4`

Solution. A plane through the intersection has equation


$$(2x+y-z-3)+\lambda(5x-3y+4z+9)=0.$$

The line direction is $(2,-4,5)$. Its dot product with the plane normal $(2+5\lambda,1-3\lambda,-1+4\lambda)$ must vanish:

$$2(2+5\lambda)-4(1-3\lambda)+5(-1+4\lambda)=0,$$

so $\lambda=5/42$. Multiplying by 42 and simplifying gives

$$\boxed{109x+27y-22z-81=0}.$$



Code reference – plot and numerical verification.


`codes/Q10.py`

Q15. Find where the line through $A(3,4,1)$ and $B(5,1,6)$ meets the plane through $P(2,1,2)$, $Q(3,1,0)$ and $R(4,-2,1)$.

Formula reference: `1.1.2.5`

Solution. A normal to the plane is $(Q-P)\times(R-P) = (1,0,-2)\times(2,-3,-1) \propto (2,1,1)$. Thus the plane is $2x+y+z=7$. The line is $(3,4,1)+t(2,-3,5)$. Substitution into the plane yields $4+6t=0$, so $t=-2/3$. The intersection is

$$\boxed{\left(\frac{5}{3}, 6, -\frac{7}{3}\right)}.$$



Code reference – plot and numerical verification.

`codes/Q15.py`

3) Differential equations:

Q20. Find the integrating factor of $x \log x \, \frac{dy}{dx} + y = 2 \log x$.

Formula reference: `1.3.1.2`

Solution. In standard form,

$$y' + \frac{1}{x \log x} y = \frac{2}{x}.$$

Thus the integrating factor is

$$\exp\left(\int \frac{dx}{x \log x} \right) = \exp(\log |\log x|).$$

On a domain where $\log x > 0$, this is $\boxed{\log x}$ (more generally the factor can be taken as $|\log x|$ on an interval not crossing $x = 1$).

Q21. Find the differential equation of the family $(x - h)^2 + (y - k)^2 = r^2$, where h, k are arbitrary constants and r is fixed.

Formula reference: `1.3.1.4`

Solution. Differentiate once:

$$(x - h) + (y - k) y' = 0.$$

Differentiate again: $1+(y')^2+(y-k)y''=0$, giving $y-k=-(1+(y')^2)/y''$. Substitute this and $x-h=-(y-k)y'$ into the original circle equation. Then

$$\frac{(1+(y')^2)^3}{(y'')^2}=r^2, \quad \boxed{(1+(y')^2)^3=r^2(y'')^2}.$$

Q22. Show that $(x-y)\frac{dy}{dx}=x+2y$ is homogeneous and solve it.

Formula reference: `1.3.1.3`

Solution. The derivative is a ratio of homogeneous first-degree expressions. Put $y=vx$, so $y'=v+xv'$. Substitution gives

$$x\frac{dv}{dx}=\frac{1+v+v^2}{1-v}, \qquad \frac{1-v}{1+v+v^2}\,dv=\frac{dx}{x}.$$

Integrating,

$$-\frac{1}{2}\ln(v^2+v+1)+\sqrt{3}\tan^{-1}\left(\frac{2v+1}{\sqrt{3}}\right)=\ln|x|+C.$$

Using $v=y/x$ gives the implicit solution

$$\boxed{-\tfrac{1}{2}\ln\left(\frac{y^2}{x^2}+\frac{y}{x}+1\right)+\sqrt{3}\tan^{-1}\left(\frac{2y/x+1}{\sqrt{3}}\right)=\ln|x|+C}.$$

Q24. Find the particular solution of $y'+y\tan x=3x^2+x^3\tan x$, given $y=0$ at $x=\pi/3$ and $x\neq\pi/2$.

Formula reference: `1.3.1.2`

Solution. The integrating factor is $e^{\int\tan x\,dx}=\sec x$. Multiplication gives

$$\frac{d}{dx}(y\sec x)=3x^2\sec x+x^3\sec x\tan x=\frac{d}{dx}(x^3\sec x).$$

So $y\sec x=x^3\sec x+C$ and $y=x^3+C\cos x$. From $y(\pi/3)=0$, $C=-2\pi^3/27$. Therefore

$$\boxed{y=x^3-\frac{2\pi^3}{27}\cos x}.$$

Code reference – numerical verification.

`codes/Q24.py`

Q25. If $y=\log\left(\frac{x}{a+bx}\right)^x$, prove that $x^3y''=(xy'-y)^2$.

Formula reference: `1.3.1.5`

Solution. Interpret the expression as $y=x\log\left(\frac{x}{a+bx}\right)$. Then

$$y'=\log\left(\frac{x}{a+bx}\right)+\frac{a}{a+bx}, \qquad y''=\frac{a^2}{x(a+bx)^2}.$$

Thus $xy' - y = ax/(a+bx)$ and

$$x^3 y'' = \frac{a^2 x^2}{(a+bx)^2} = (xy' - y)^2.$$

4) Matrices and determinants:

Q28. If $A = \begin{bmatrix} \cos\theta & \sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$, find $\det(A^n)$ for natural n.

Formula reference: `1.2.1.1`

Solution. Determinants are multiplicative, so $\det(A^n) = (\det A)^n$. Now $\det A = \cos^2\theta - \sin^2\theta = \cos 2\theta$. Thus

$$\boxed{\det(A^n) = (\cos 2\theta)^n}.$$

Q30. Find the inverse of $\begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 0 \end{bmatrix}$ using elementary row operations.

Formula reference: `1.2.1.2`

Solution. Augment the matrix by I and row-reduce to $[I \mid A^{-1}]$. The determinant is -1, so the matrix is invertible. The result is

$$\boxed{A^{-1} = \begin{bmatrix} 3 & -2 & 1 \\ -9 & 6 & -2 \\ 5 & -3 & 1 \end{bmatrix}}.$$

The product check $AA^{-1} = I$ is carried out in code.

Code reference – numerical verification.

`codes/Q30.py`

Q36. If $a+b+c \neq 0$ and $\begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix} = 0$, prove that $a = b = c$.

Formula reference: `1.2.2.1`

Solution. Expanding and factorising the determinant gives

$$-(a+b+c)(a^2+b^2+c^2-ab-bc-ca) = 0.$$

The first factor is nonzero by assumption, so $a^2+b^2+c^2-ab-bc-ca = 0$. But

$$a^2+b^2+c^2-ab-bc-ca = \tfrac{1}{2}[(a-b)^2+(b-c)^2+(c-a)^2].$$

Each square must be zero, hence $\boxed{a = b = c}$.

Q37. A trust receives Rs. 30,000 per month. Half is spent on care, and 2% of the amount spent is collected from beneficiaries. The unspent balance is deposited in a bank. Find the monthly interest rate needed for total monthly income of Rs. 1800.

Formula reference: `1.2.1.4`

Solution. Let u be the monthly contribution collected from beneficiaries and v the bank interest. Half of Rs. $30,000$ is spent, so

$$u = 0.02(15000) = 300.$$

The total monthly income is Rs. 1800, so

$$u + v = 1800, \qquad u = 300.$$

Write this system as an augmented matrix:

$$\left[\begin{array}{cc|c} 1 & 1 & 1800 \\ 1 & 0 & 300 \end{array}\right] \xrightarrow{R_2 \leftarrow R_2 - R_1} \left[\begin{array}{cc|c} 1 & 1 & 1800 \\ 0 & -1 & -1500 \end{array}\right] \xrightarrow{R_2 \leftarrow -R_2} \left[\begin{array}{cc|c} 1 & 1 & 1800 \\ 0 & 1 & 1500 \end{array}\right].$$

Eliminate v from the first row:

$$R_1 \leftarrow R_1 - R_2: \qquad \left[\begin{array}{cc|c} 1 & 0 & 300 \\ 0 & 1 & 1500 \end{array}\right].$$

Thus $u = 300$ and $v = 1500$. The deposit is Rs. 15000, therefore

$$r = \frac{1500}{15000} \times 100\% = \boxed{10\% \text{ per month}}.$$

Code reference – numerical verification.

`codes/Q37.py`

5) Integration and area:

Q38. Evaluate $I = \displaystyle\int_0^{\pi/2} \frac{\cos^2 x}{1 + 3\sin^2 x}\, dx.$

Formula reference: `1.4.1.1, 1.4.1.2`

Solution. Let $t = \tan x$. Then $dx = dt/(1+t^2)$ and the integral becomes

$$I = \int_0^{\infty} \frac{dt}{(1+t^2)(1+4t^2)} = \int_0^{\infty} \left[-\frac{1}{3(1+t^2)} + \frac{4}{3(1+4t^2)}\right] dt.$$

Evaluating the arctangent limits gives $I = -\pi/6 + \pi/3 = \boxed{\pi/6}$.

Q39. Evaluate $\displaystyle\int_0^{\pi/4} \frac{\sin x + \cos x}{3 + \sin 2x}\, dx.$

Formula reference: `1.4.1.1`

Solution. Set $u = \sin x - \cos x$, so $du = (\sin x + \cos x)dx$ and $\sin 2x = 1 - u^2$. The limits become $u = -1$ and $u = 0$. Hence

$$I = \int_{-1}^{0} \frac{du}{4 - u^2} = \left[\frac{1}{4} \ln \left|\frac{2+u}{2-u}\right|\right]_{-1}^{0} = \boxed{\frac{1}{4}\ln 3}.$$

Q40. Evaluate $\displaystyle\int \frac{\sin x - x\cos x}{x(x + \sin x)}\, dx.$

Formula reference: `1.3.1.5`

Solution. Since

$$\frac{d}{dx}\ln\left|\frac{x}{x+\sin x}\right| = \frac{\sin x - x\cos x}{x(x+\sin x)},$$

the integral is $\boxed{\ln\left|\dfrac{x}{x+\sin x}\right| + C}$.

Q41. Evaluate $\displaystyle\int \frac{x^3}{(x-1)(x^2+1)}\,dx$.

Formula reference: `1.4.1.5`

Solution. Polynomial division and partial fractions give

$$\frac{x^3}{(x-1)(x^2+1)} = 1 + \frac{1}{2(x-1)} + \frac{x-1}{2(x^2+1)}.$$

Therefore

$$\boxed{x + \frac{1}{2}\ln|x-1| + \frac{1}{4}\ln(x^2+1) - \frac{1}{2}\tan^{-1}x + C}.$$

Q42. Find $\displaystyle\int \frac{x\,dx}{1+x\tan x}$.

Formula reference: `1.3.1.5`

Solution. Rewrite the integrand as $x\cos x/(\cos x + x\sin x)$. Since $d(\cos x + x\sin x)/dx = x\cos x$,

$$\boxed{\int \frac{x\,dx}{1+x\tan x} = \ln|\cos x + x\sin x| + C}.$$

Q43. Evaluate $\displaystyle\int \frac{x^4}{(x-1)(x^2+1)}\,dx$.

Formula reference: `1.4.1.5`

Solution. Division gives $x^4/[(x-1)(x^2+1)] = x + 1 + 1/[(x-1)(x^2+1)]$. Decompose the last term as

$$\frac{1}{(x-1)(x^2+1)} = \frac{1}{2(x-1)} - \frac{x+1}{2(x^2+1)}.$$

Thus

$$\boxed{\frac{x^2}{2} + x + \frac{1}{2}\ln|x-1| - \frac{1}{4}\ln(x^2+1) - \frac{1}{2}\tan^{-1}x + C}.$$

Q44. Evaluate $\displaystyle\int_0^{\pi/2} \frac{5\sin x + 3\cos x}{\sin x + \cos x}\,dx$.

Formula reference: `1.3.1.5`

Solution. Write the numerator as $4(\sin x + \cos x) + (\sin x - \cos x)$. Then

$$I = 4\int_0^{\pi/2} dx - [\ln(\sin x + \cos x)]_0^{\pi/2} = \boxed{2\pi}.$$

Q45. Find $\displaystyle\int \left[\log(\log x) + \frac{1}{(\log x)^2}\right] dx$.

Formula reference: `1.3.1.5, 1.4.1.6`

Solution. Differentiate $x\log(\log x) - x/\log x$ to check that it equals the integrand. Hence

$$\boxed{x\log(\log x) - \frac{x}{\log x} + C}, \qquad x > 1 \text{ for a real-valued } \log(\log x).$$

Q46. If the area bounded by $y^2 = 16ax$ and $y = 4mx$ is $a^2/2$, find m.

Formula reference: `1.4.1.4`

Solution. Use the upper branch $y = 4\sqrt{ax}$. At an intersection with $y = 4mx$,

$$4\sqrt{ax} = 4mx \quad\Longrightarrow\quad x = 0 \quad \text{or} \quad x = \frac{a}{m^2}.$$


For positive slope, the area is the integral of the parabola minus the line:

$$\begin{aligned} A &= \int_0^{a/m^2} \left(4\sqrt{ax} - 4mx\right) dx \\ &= \left[\frac{8\sqrt{a}}{3}x^{3/2} - 2mx^2\right]_0^{a/m^2} \\ &= \frac{8a^2}{3m^3} - \frac{2a^2}{m^3} = \frac{2a^2}{3m^3}. \end{aligned}$$

Set the result equal to the given area:

$$\frac{2a^2}{3m^3} = \frac{a^2}{2} \quad\Longrightarrow\quad 4 = 3m^3 \quad\Longrightarrow\quad \boxed{m = \sqrt[3]{\frac{4}{3}}}.$$

The graph uses $a = 1$ for visibility; the value of m is independent of that scale.



Code reference – plot and numerical verification.

`codes/Q46.py`


Q47. Find the area of $\{(x, y) : x^2 + y^2 \le 4,\ x + y \ge 2\}$ by integration.

Formula reference: `1.4.1.4`

Solution. The chord $x + y = 2$ meets the circle at $(0, 2)$ and $(2, 0)$. For $0 \le x \le 2$, the upper circle boundary is $y = \sqrt{4 - x^2}$ and the line is $y = 2 - x$. Therefore

$$\begin{aligned} A &= \int_0^2 \left(\sqrt{4 - x^2} - (2 - x) \right) dx \\ &= \underbrace{\int_0^2 \sqrt{4 - x^2}\, dx}_{\text{quarter-circle area } = \pi} - \underbrace{\int_0^2 (2 - x)\, dx}_{\text{triangle area } = 2} \\ &= \boxed{\pi - 2} \quad \text{square units.} \end{aligned}$$

The graph shades the circular segment above the chord.



Code reference – plot and numerical verification.

`codes/Q47.py`

6) Functions and relations:

Q48. Discuss continuity and differentiability of $f(x) = |x| + |x - 1|$ on $(-1, 2)$.

Formula reference: `1.5.1.1`

Solution. First split the absolute values at $x = 0$ and $x = 1$:

$$f(x) = \begin{cases} 1 - 2x, & -1 < x < 0, \\ 1, & 0 \leq x < 1, \\ 2x - 1, & 1 \leq x < 2. \end{cases}$$

Continuity. At $x = 0$, the left limit, right limit and function value are all 1. The same is true at $x = 1$. Each branch is continuous on its interval; hence f is continuous throughout $(-1, 2)$.

Differentiability at $x = 0$. The left-hand slope is -2 and the right-hand slope is 0:


$$f'_-(0) = -2 \neq 0 = f'_+(0).$$

Thus f is not differentiable at 0.

Differentiability at $x = 1$. The left-hand slope is 0 and the right-hand slope is 2:

$$f'_-(1) = 0 \neq 2 = f'_+(1).$$

Thus f is not differentiable at 1. At other points the function is linear and differentiable. The plot shows these one-sided slopes: differentiability requires equal one-sided slopes, and the two corners fail that test.



Code reference – plot and slope verification.

`codes/Q48.py`

Q49. On $\mathbb{R}$, define $(a, b) \in R$ when $a - b + \sqrt{3}$ is irrational. Determine whether R is reflexive, symmetric and transitive.

Formula reference: `1.5.1.2`

Solution. We test each property separately.

1. Reflexive. Reflexivity requires $(a, a) \in R$ for every real a. Substituting $b = a$ gives

$$a - a + \sqrt{3} = \sqrt{3},$$

which is irrational. Therefore R is reflexive.

2. Symmetric. Symmetry requires $(a, b) \in R$ to imply $(b, a) \in R$. Choose $a = \sqrt{3}$ and $b = 0$. Then

$$a - b + \sqrt{3} = 2\sqrt{3}$$

is irrational, so $(a, b) \in R$. Reversing the pair gives

$$b - a + \sqrt{3} = 0,$$

which is rational, so $(b, a) \notin R$. Hence R is not symmetric.

3. Transitive. Transitivity requires $(a,b),(b,c) \in R$ to imply $(a,c) \in R$. Choose $a = \sqrt{3}$, $b = 0$, and $c = 2\sqrt{3}$. Then

$$a - b + \sqrt{3} = 2\sqrt{3}, \qquad b - c + \sqrt{3} = -\sqrt{3},$$

both irrational, so $(a,b),(b,c) \in R$. However,

$$a - c + \sqrt{3} = \sqrt{3} - 2\sqrt{3} + \sqrt{3} = 0,$$

which is rational. Thus $(a,c) \notin R$, and R is not transitive.
Conclusion. $\boxed{R \text{ is reflexive, but neither symmetric nor transitive.}}$

Q50. On $\mathbb{R} \times \mathbb{R}$, define $(a,b) * (c,d) = (a + c, b + d)$. Prove commutativity and associativity, and find the identity and the inverse of $(3,-5)$.

Formula reference: `1.5.1.3`

Solution. Addition of real components is commutative and associative, so $*$ has both properties. Its identity is $\boxed{(0,0)}$, since $(a,b) * (0,0) = (a,b)$. The inverse of $(3,-5)$ is $\boxed{(-3,5)}$.

Q51. On $\{0,1,2,3,4,5,6\}$ define $a * b = a + b$ if $a + b < 7$, and $a * b = a + b - 7$ otherwise. Give the operation table and show that 0 is the identity and $7 - a$ is the inverse for nonzero a.

Formula reference: `1.5.1.4`

Solution. This is addition modulo 7. Its table is

$*$	0	1	2	3	4	5	6
0	0	1	2	3	4	5	6
1	1	2	3	4	5	6	0
2	2	3	4	5	6	0	1
3	3	4	5	6	0	1	2
4	4	5	6	0	1	2	3
5	5	6	0	1	2	3	4
6	6	0	1	2	3	4	5

Row and column 0 reproduce every element, hence 0 is the identity. For each $a = 1,\ldots,6$, $a + (7 - a) = 7 \equiv 0 \pmod 7$, so $7 - a$ is its inverse.


Q52. Find any point of discontinuity of $f(x) = x - |x - x^2|$, $x \in [-1,1]$.

Formula reference: `1.5.1.1`

Solution. On $[-1,1]$, $x - x^2 = x(1 - x)$, whose sign changes only at 0. Thus

$$f(x) = \begin{cases} 2x - x^2, & -1 \le x \le 0, \\ x^2, & 0 \le x \le 1. \end{cases}$$

Both pieces take value 0 at $x = 0$ and meet continuously. The endpoints are also included continuously. Therefore there are $\boxed{\text{no points of discontinuity}}$.



Code reference – plot.

`codes/Q52.py`

7) Probability:

Q53. A man steps forward with probability 0.4 and backward with probability 0.6. Find the probability that after 5 steps he is one step away from his starting point.

Formula reference: `1.6.1.3, 1.6.1.4`

Solution. Let k be the number of forward steps. The net displacement after five steps is

$$D = k - (5 - k) = 2k - 5.$$

One step away means $D = +1$ or $D = -1$. These occur when $k = 3$ or $k = 2$, respectively.
Step 1: Net displacement $+1$. Three steps are forward and two backward, so the binomial formula gives

$$\begin{aligned} P(D = 1) &= \binom{5}{3}(0.4)^3(0.6)^2 \\ &= 10(0.064)(0.36) = 0.2304. \end{aligned}$$

Step 2: Net displacement -1. Two steps are forward and three backward:

$$\begin{aligned} P(D = -1) &= \binom{5}{2}(0.4)^2(0.6)^3 \\ &= 10(0.16)(0.216) = 0.3456. \end{aligned}$$


Step 3: Add the disjoint events.

$$P(|D| = 1) = P(D = 1) + P(D = -1) = 0.2304 + 0.3456 = \boxed{0.576}.$$

Generating function / Z-transform. For one step, the Laurent probability generating function is $G(z) = 0.4z + 0.6z^{-1}$. For five independent steps,

$$G_5(z) = (0.4z + 0.6z^{-1})^5 = \sum_{k=0}^{5} \binom{5}{k} (0.4)^k (0.6)^{5-k} z^{2k-5}.$$

The coefficient of z^1 is 0.2304 and the coefficient of z^{-1} is 0.3456. Their sum is the required probability. The plot shows the theoretical PMF for every possible displacement.



Code reference – plot and numerical verification.

`codes/Q53.py`

Q54. A girl throws a die. If it shows 1 or 2, she tosses a coin three times; otherwise she tosses it once. Given that exactly one tail was obtained, find the probability that the die showed 3, 4, 5 or 6.

Formula reference: `1.6.1.2, 1.6.1.7`

Solution. Let A mean that the die shows 1 or 2, B mean that it shows 3, 4, 5 or 6, and E mean that exactly one tail is observed.

Step 1: Find the prior probabilities.

$$P(A) = \frac{2}{6} = \frac{1}{3}, \qquad P(B) = \frac{4}{6} = \frac{2}{3}.$$

Step 2: Find the likelihood of E. If A occurs, three fair coin tosses are made. Exactly one tail can occur in $\binom{3}{1}$ of the 8 equally likely outcomes:

$$P(E \mid A) = \binom{3}{1} \left(\frac{1}{2}\right)^3 = \frac{3}{8}.$$

If B occurs, only one coin toss is made, so

$$P(E \mid B) = \frac{1}{2}.$$

Step 3: Calculate $P(E)$ by total probability.

$$\begin{aligned} P(E) &= P(E \mid A)P(A) + P(E \mid B)P(B) \\ &= \frac{3}{8} \cdot \frac{1}{3} + \frac{1}{2} \cdot \frac{2}{3} \\ &= \frac{1}{8} + \frac{1}{3} = \frac{11}{24}. \end{aligned}$$

Step 4: Apply Bayes' theorem.

$$P(B \mid E) = \frac{P(E \mid B)P(B)}{P(E)} = \frac{(1/2)(2/3)}{11/24} = \boxed{\frac{8}{11}}.$$

Q55. An urn contains 5 red and 2 black balls. Two balls are drawn without replacement. Let X be the number of black balls. State its possible values and find its mean and variance.


Formula reference: `1.6.1.5, 1.6.1.6`

Solution. The possible values are $0, 1, 2$. The probabilities from the hypergeometric model are

$$P(X = 0) = \frac{10}{21}, \quad P(X = 1) = \frac{10}{21}, \quad P(X = 2) = \frac{1}{21}.$$

So $E[X] = 0(10/21) + 1(10/21) + 2(1/21) = \boxed{4/7}$. Also $E[X^2] = 10/21 + 4/21 = 2/3$, hence

$$\mathrm{Var}(X) = E[X^2] - E[X]^2 = \frac{2}{3} - \frac{16}{49} = \boxed{\frac{50}{147}}.$$



Code reference – plot and numerical verification.

codes/Q55.py

Q56. In 3 trials of a binomial distribution, the probability of exactly 2 successes is 9 times the probability of 3 successes. Find the success probability.

Formula reference: `1.6.1.3`

Solution. Let the success probability be p and $q = 1 - p$. The probability-generating polynomial is

$$B(z) = (q + pz)^3 = q^3 + 3pq^2z + 3p^2qz^2 + p^3z^3.$$

The coefficient of z^2 is

$$P(X = 2) = \binom{3}{2} p^2 q = 3p^2q,$$

and the coefficient of z^3 is

$$P(X = 3) = p^3.$$

Use $P(X = 2) = 9P(X = 3)$:

$$3p^2q = 9p^3,$$
$$3p^2(1 - p) = 9p^3.$$

For the non-degenerate binomial case $p > 0$, divide by $3p^2$: $1 - p = 3p$. Thus $4p = 1$ and

$$\boxed{p = \frac{1}{4}}.$$

The polynomial's coefficients directly identify the two probabilities; no separate Z-transform is needed beyond this generating-polynomial viewpoint.

Code reference – numerical verification.

codes/Q56.py

Q57. An urn contains 3 red and 5 black balls. A ball is drawn, its colour recorded and returned, and 2 more balls of that colour are added. Then two balls are drawn without replacement. Find the probability both are red.

Formula reference: `1.6.1.7`

Solution. If the first ball is red, with probability $3/8$, the urn becomes 5 red and 5 black; the chance of two reds is $(5/10)(4/9) = 2/9$. If the first ball is black, with probability $5/8$, the urn becomes 3 red and 7 black; the chance of two reds is $(3/10)(2/9) = 1/15$. Thus

$$P(\text{two red}) = \frac{3}{8}\frac{2}{9} + \frac{5}{8}\frac{1}{15} = \boxed{\frac{1}{8}}.$$

Code reference – numerical verification.

codes/Q57.py

Q58. A man speaks truth 3 out of 5 times. He throws a die and reports that it is 4. Find the probability that it actually is 4.

Formula reference: `1.6.1.2`

Solution. Let A be the event that the die shows 4, and R the event that he reports 4. Assuming that when he lies each of the five wrong numbers is equally likely, $P(A) = 1/6$, $P(R|A) = 3/5$, and $P(R|A^c) = (2/5)(1/5) = 2/25$. Hence

$$P(A|R) = \frac{(1/6)(3/5)}{(1/6)(3/5) + (5/6)(2/25)} = \boxed{\frac{3}{5}}.$$


8) Linear programming:

Q59. Solve graphically: minimise $z = 3x + 5y$ subject to $x + 2y \geq 10$, $x + y \geq 6$, $3x + y \geq 8$, $x, y \geq 0$.

Formula reference: `1.7.1.1`

Solution. The lower-boundary vertices are $(0, 8)$, $(1, 5)$, $(2, 4)$ and $(10, 0)$. Their objective values are respectively 40, 28, 26 and 30. The smallest value is attained at $(2, 4)$:

$$\boxed{x = 2,\ y = 4,\ z_{\min} = 26}.$$



Code reference – plot and numerical verification.


`codes/Q59.py`

Q60. A dealer has Rs. 57,600 and space for at most 20 sewing machines. An electronic machine costs Rs. 3600 and earns Rs. 220 profit; a manual machine costs Rs. 2400 and earns Rs. 180 profit. How should he invest to maximise profit?

Formula reference: `1.7.1.1`

Solution. Let x and y be electronic and manual machines. The constraints are $3x+2y \le 48$, $x+y \le 20$, $x, y \ge 0$, and profit is $P = 220x + 180y$. At the intersection of the two active limits, $3x + 2y = 48$ and $x + y = 20$, we get $(x, y) = (8, 12)$ and $P = 3920$. The axis vertices give profits Rs. 3520 and Rs. 3600. Therefore the best feasible integer plan is

$$\boxed{8 \text{ electronic and } 12 \text{ manual machines; maximum profit Rs. } 3920}.$$



Code reference – plot and numerical verification.

`codes/Q60.py`


9) Coordinate geometry and optimisation:

Q64. A tangent to $x^2 + y^2 = 4$ at a point in the first quadrant meets the positive axes at A and B. Find the minimum of $OA + OB$.

Formula reference: `1.8.1.1`

Solution. At the point $(2\cos\theta, 2\sin\theta)$, the tangent is $x\cos\theta + y\sin\theta = 2$. Thus $OA = 2\sec\theta$ and $OB = 2\csc\theta$. By symmetry (or differentiating), their sum is minimised when $\sin\theta = \cos\theta = 1/\sqrt{2}$. Hence

$$\boxed{\min(OA + OB) = 2(\sqrt{2} + \sqrt{2}) = 4\sqrt{2}}.$$



Code reference – plot and numerical verification.

`codes/Q64.py`

Q65. Find the maximum area of an isosceles triangle inscribed in $\frac{x^2}{16} + \frac{y^2}{9} = 1$ with its vertex at an end of the major axis.


Formula reference: `1.8.1.2`

Solution. By symmetry, take the fixed vertex as $(4, 0)$ and the other two vertices as (u, v) and $(u, -v)$. From the ellipse, $v = 3\sqrt{1 - u^2/16}$. The area is

$$A(u) = v(4 - u) = 3(4 - u)\sqrt{1 - u^2/16}.$$

Differentiation gives $A'(u) = 0$ when $u^2 - 2u - 8 = 0$, i.e. $u = 4$ or $u = -2$. The endpoint $u = 4$ has zero area, so the maximum is at $u = -2$, where $v = 3\sqrt{3}/2$. Thus

$$\boxed{A_{\max} = 6\left(\frac{3\sqrt{3}}{2}\right) = 9\sqrt{3}}.$$



Code reference – plot and numerical verification.

codes/Q65.py