Package leaflet: Information for the user

IE

Entecavir Viatris 0.5 mg

film-coated tablets

entecavir

Entecavir Viatris 1 mg

film-coated tablets

entecavir

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Entecavir Viatris is and what it is used for
2. What you need to know before you take Entecavir Viatris
3. How to take Entecavir Viatris
4. Possible side effects
5. How to store Entecavir Viatris
6. Contents of the pack and other information

1. What Entecavir Viatris is and what it is used for

Entecavir Viatris tablets are anti-viral medicines, used to treat chronic (long term) hepatitis B virus (HBV) infection in adults. Entecavir Viatris can be used in people whose liver is damaged but still functions properly (compensated liver disease) and in people whose liver is damaged and does not function properly (decompensated liver disease).

Entecavir Viatris tablets are also used to treat chronic (long term) HBV infection in children and adolescents aged 2 years to less than 18 years. Entecavir Viatris can be used in children whose liver is damaged but still functions properly (compensated liver disease).

Infection by the hepatitis B virus can lead to damage to the liver. Entecavir Viatris reduces the amount of virus in your body, and improves the condition of the liver.

2. What you need to know before you take Entecavir Viatris

Do not take Entecavir Viatris

- **if you are allergic (hypersensitive)** to entecavir or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor or pharmacist before taking Entecavir Viatris

- **if you have ever had problems with your kidneys**, tell your doctor. This is important because Entecavir Viatris is eliminated from your body through the kidneys and your dose or dosing schedule may need to be adjusted.
- **do not stop taking Entecavir Viatris without your doctor's advice** since your hepatitis may worsen after stopping treatment. When your treatment with Entecavir Viatris is stopped, your doctor will continue to monitor you and take blood tests for several months.
- **discuss with your doctor whether your liver functions properly** and, if not, what the possible effects on your Entecavir Viatris treatment may be.
- **if you are also infected with HIV** (human immunodeficiency virus) be sure to tell your doctor. You should not take Entecavir Viatris to treat your hepatitis B infection unless you are taking medicines for HIV at the same time, as the effectiveness of future HIV treatment may be reduced. Entecavir Viatris will not control your HIV infection.
- **taking Entecavir Viatris will not stop you from infecting other people with hepatitis B virus (HBV)** through sexual contact or body fluids (including blood contamination). So, it is important to take appropriate precautions to prevent others from becoming infected with HBV. A vaccine is available to protect those at risk from becoming infected with HBV.
- **Entecavir Viatris belongs to a class of medicines that can cause lactic acidosis** (excess of lactic acid in your blood) and enlargement of the liver. Symptoms such as nausea, vomiting and stomach pain might indicate the development of lactic acidosis. This rare but serious side effect has occasionally been fatal. Lactic acidosis occurs more often in women, particularly if they are very overweight. Your doctor will monitor you regularly while you are receiving Entecavir Viatris.
- **if you have previously received treatment for chronic hepatitis B**, please inform your doctor.

Children and adolescents

Entecavir Viatris should not be used for children below 2 years of age or weighing less than 10 kg.

Other medicines and Entecavir Viatris

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Entecavir Viatris with food and drink

In most cases you may take Entecavir Viatris with or without food. However, if you have had a previous treatment with a medicine containing the active substance lamivudine you should consider the following. If you were switched over to Entecavir Viatris because the treatment with lamivudine was not successful, you should take Entecavir Viatris on an empty stomach once daily. If your liver disease is very advanced, your doctor will also instruct you to take Entecavir Viatris on an empty stomach. Empty stomach means at least 2 hours after a meal and at least 2 hours before your next meal.

Children and adolescents (from 2 to less than 18 years of age) can take Entecavir Viatris with or without food.

Pregnancy, breast-feeding and fertility

Tell your doctor if you are pregnant or planning to become pregnant. It has not been demonstrated that entecavir is safe to use during pregnancy. Entecavir Viatris must not be used during pregnancy unless specifically directed by your doctor. It is important that women of childbearing age receiving treatment with Entecavir Viatris use an effective method of contraception to avoid becoming pregnant.

You should not breast-feed during treatment with Entecavir Viatris. Tell your doctor if you are breastfeeding. It is not known whether entecavir, the active substance in Entecavir Viatris, is excreted in human breast milk.

Driving and using machines

Dizziness, tiredness (fatigue) and sleepiness (somnolence) are common side effects which may impair your ability to drive and use machines. If you have any concerns consult your doctor.

Entecavir Viatris contains lactose

This medicine contains lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Entecavir Viatris

Not all patients need to take the same dose of Entecavir Viatris.

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Mylan				Date of Final	dd / mm / yy	Date of Issue	dd / mm / yy Page Count 1 of 2
Date/Time of Creation	29/01/26 16:40	TrackWise / GLAMS ID	4036385	Client Market	Ireland	Software / Ver. No.	Adobe InDesign 2025
Vendor Job No.	NA	Affiliate New Code	4036385	Product Description	LIT(BF) ENTECAVIR VIA TABLETS(COM)IE V8		
Artwork Proof No.	1	Aff. Superseded Code	3744903				
Pharma Code	NA	Barcode Information	NA	New Material Code	75111617	Actual A/w Size	Flat-170 x 400 mm
Product Render ID No.	NA	MA Number	EU/1/17/1227/002	ITF Barcode		Other Sizes (if any)	Folded-170 x 40mm
Main Font Type - 1	Helvetica Neue LT Pro	SAP or SPC Code	400539223	Superseded Code	75108802	Pitch After Every	NA
Main Font Type - 2	NA	Packaging Site	Mylan Laboratories Ltd. , FDF 3 (Pithampur, Dhar Dist., M.P. - IN)	Component Type	Printed Literature	Eye / Sensor Mark	NA
Min. Font Size	10 pt	Mfg. Lic. No. / Code No.	NA	No. of Printed Colors	1	ITF Barcode Size	NA
Reason for Revision / Issuance	Labeling			Printed Colors CMYK / PMS Coated / PMS Color Bridge Coated	BLACK 100%		
	NA			Non Printed Colors	Keyline		
	NA			Equivalent with CMYK (Pantones Ref. Code)	NA		
Artwork Implementation Schedule, (✓) whichever is applicable	New Component			Material of Construction	40gsm ITC Tribeni Paper		
	Immediately (Stock of supperseded component to be destroyed, if applicable)			Design & Style	Supply in Folded form as Proposed Size		
	After Consumption of Existing (supperseded) stock						
	Others (specify)						

LABEL CONTROL / BUSINESS DEVELOPMENT / REGULATORY / MARKETING				Prepared By	Checked By	Approved By		
				Packaging Technical Services		Regulatory Affairs	Production	Quality Assurance
Sign Offs	Digital Signature	Digital Signature	Digital Signature	Digital Signature	Sign, Date OR Digital Signature	Sign, Date OR Digital Signature	Sign, Date OR Digital Signature	Sign, Date OR Digital Signature	Sign, Date OR Digital Signature
Revision History	23.01.2026- Keyline Revised Date / Remarks Date / Remarks		Date / Remarks Date / Remarks Date / Remarks			Date / Remarks Date / Remarks Date / Remarks		

Reference SOP-000459814 FORM-000644250 Version 3.0

#Note: This artwork is electronically signed through TrackWise/GLAMS, hence wet ink/digital signatures are not done/required.

For adults the recommended dose is either 0.5 mg or 1 mg once daily orally (by mouth).

Your dose will depend on:
- whether you have been treated for HBV infection before, and what medicine you received.
- whether you have kidney problems. Your doctor may prescribe a lower dose for you or instruct you to take it less often than once a day.
- the condition of your liver.

For children and adolescents (from 2 to less than 18 years of age), your child's doctor will decide the right dose based on your child's weight. Children weighing at least 32.6 kg may take the 0.5 mg tablet or an entecavir oral solution may be available. For patients weighing from 10 kg to 32.5 kg, an entecavir oral solution is recommended. All doses should be taken once daily orally (by mouth). There are no recommendations for entecavir in children less than 2 years of age or weighing less than 10 kg.

For children and adolescents (from 2 to less than 18 years of age), Entecavir Viatris 0.5 mg Film-coated Tablets are available or an oral solution may be available. Your child's doctor will decide the right dose based on your child's weight.

Your doctor will advise you on the dose that is right for you.
Always take the dose recommended by your doctor to ensure that your medicine is fully effective and to reduce the development of resistance to treatment. Take Entecavir Viatris as long as your doctor has told you. Your doctor will tell you if and when you should stop the treatment.

Some patients must take Entecavir Viatris on an empty stomach (see **Entecavir Viatris with food and drink in Section 2**). If your doctor instructs you to take Entecavir Viatris on an empty stomach, empty stomach means at least 2 hours after a meal and at least 2 hours before your next meal.

If you take more Entecavir Viatris than you should
Contact your doctor at once.

If you forget to take Entecavir Viatris
It is important that you do not miss any doses. If you miss a dose of Entecavir Viatris, take it as soon as possible, and then take your next scheduled dose at its regular time. If it is almost time for your next dose, do not take the missed dose. Wait and take the next dose at the regular time. Do not take a double dose to make up for a forgotten dose.

Do not stop Entecavir Viatris without your doctor's advice
Some people get very serious hepatitis symptoms when they stop taking entecavir. Tell your doctor immediately about any changes in symptoms that you notice after stopping treatment.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Patients treated with entecavir have reported the following side effects:
Adults
- Common (at least 1 in 100 patients): headache, insomnia (inability to sleep), fatigue (extreme tiredness), dizziness, somnolence (sleepiness), vomiting, diarrhoea, nausea, dyspepsia (indigestion), and increased blood levels of liver enzymes.
- Uncommon (at least 1 in 1,000 patients): rash, hair loss.
- Rare (at least 1 in 10,000 patients): severe allergic reaction.

Children and adolescents
The side effects experienced in children and adolescents are similar to those experienced in adults as described above with the following difference:

Very common (at least 1 in 10 patients): low levels of neutrophils (one type of white blood cells, which are important in fighting infection).

Reporting of side effects
If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRA Pharmacovigilance, Website: www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Entecavir Viatris

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the bottle, blister or carton after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Entecavir Viatris contains

Entecavir Viatris 0.5 mg film-coated tablets
The active substance is entecavir. Each film-coated tablet contains entecavir monohydrate equivalent to 0.5 mg entecavir.

Entecavir Viatris 1 mg film-coated tablets
The active substance is entecavir. Each film-coated tablet contains entecavir monohydrate equivalent to 1 mg entecavir.

The other ingredients are:
Tablet core: microcrystalline cellulose, crospovidone, lactose monohydrate (see section 2, "Entacavir Viatris contains lactose"), magnesium stearate
Tablet coating: titanium dioxide (E171), hypromellose, macrogol 400, polysorbate 80.

What Entecavir Viatris looks like and contents of the pack

Entecavir Viatris 0.5 mg film-coated tablets
The film-coated tablets (tablets) are white, film-coated, round, biconvex, beveled edge tablet debossed with "M" on one side and "EA" on the other side.

Entecavir Viatris 1 mg film-coated tablets
The film-coated tablets (tablets) are white, film-coated, round, biconvex, beveled edge tablet debossed with "M" on one side and "EB" on the other side.

Entecavir Viatris film-coated tablets are supplied in blisters packs containing 30 tablets, perforated blister packs containing 30 x 1 or 90 x 1 tablets, and in bottles containing 30 or 90 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder
Viatris Limited
Damastown Industrial Park,
Mulhuddart,
Dublin 15,
DUBLIN
Ireland

Manufacturer

Mylan Hungary Kft.
Mylan utca 1,
Komárom - 2900
Hungary

Mylan Germany GmbH
Zweigniederlassung Bad Homburg v. d. Hoehe,
Benzstrasse 1
Bad Homburg v. d. Hoehe
Hessen, 61352,
Germany

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

Ireland
Viatris Limited
Tel: +353 1 8711600

This leaflet was last revised in 16 January 2026

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency web site: http://www.ema.europa.eu.

VIATRIS
4036385
75111617

Mylan				Date of Final	dd / mm / yy	Date of Issue	dd / mm / yy	Page Count	2 of 2
Date/Time of Creation	29/01/26 16:40	TrackWise / GLAMS ID	4036385	Client Market	Ireland	Software / Ver. No.	Adobe InDesign 2025		
Vendor Job No.	NA	Affiliate New Code	4036385	Product Description	LIT(BF) ENTECAVIR VIA TABLETS(COM)IE V8				
Artwork Proof No.	1	Aff. Superseded Code	3744903						
Pharma Code	NA	Barcode Information	NA	New Material Code	75111617	Actual A/w Size	Flat-170 x 400 mm		
Product Render ID No.	NA	MA Number	EU/1/17/1227/002	ITF Barcode		Other Sizes (if any)	Folded-170 x 40mm		
Main Font Type - 1	Helvetica Neue LT Pro	SAP or SPC Code	400539223	Superseded Code	75108802	Pitch After Every	NA		
Main Font Type - 2	NA	Packaging Site	Mylan Laboratories Ltd. , FDF 3 (Pithampur, Dhar Dist., M.P. - IN)	Component Type	Printed Literature	Eye / Sensor Mark	NA		
Min. Font Size	10 pt	Mfg. Lic. No. / Code No.	NA	No. of Printed Colors	1	ITF Barcode Size	NA		
Reason for Revision / Issuance	Labeling			Printed Colors CMYK / PMS Coated / PMS Color Bridge Coated	BLACK 100%				
	NA			Non Printed Colors	Keyline				
	NA			Equivalent with CMYK (Pantones Ref. Code)	NA				
Artwork Implementation Schedule, (✓) whichever is applicable	New Component			Material of Construction	40gsm ITC Tribeni Paper				
	Immediately (Stock of supperseded component to be destroyed, if applicable)			Design & Style	Supply in Folded form as Proposed Size				
	After Consumption of Existing (supperseded) stock								
	Others (specify)								

LABEL CONTROL / BUSINESS DEVELOPMENT / REGULATORY / MARKETING				Prepared By	Checked By	Approved By		
				Packaging Technical Services		Regulatory Affairs	Production	Quality Assurance
Sign Offs	Digital Signature	Digital Signature	Digital Signature	Digital Signature	Sign, Date OR Digital Signature	Sign, Date OR Digital Signature	Sign, Date OR Digital Signature	Sign, Date OR Digital Signature	Sign, Date OR Digital Signature
Revision History	23.01.2026- Keyline Revised Date / Remarks Date / Remarks		Date / Remarks Date / Remarks Date / Remarks			Date / Remarks Date / Remarks Date / Remarks		

Reference SOP-000459814 FORM-000644250 Version 3.0

#Note: This artwork is electronically signed through TrackWise/GLAMS, hence wet ink/digital signatures are not done/required.