



## Chapter – 13


### Hydrocarbons

#### NCERT Back Exercises :


**Ques 13.1:** How do you account for the formation of ethane during chlorination of methane?

**Ans 13.1:** Chlorination of methane proceeds via a free radical chain mechanism. The whole reaction takes place in the given three steps.


Step 1: Initiation: The reaction begins with the homolytic cleavage of Cl – Cl bond as:





Step 2: Propagation: In the second step, chlorine free radicals attack methane molecules and break down the C–H bond to generate methyl radicals as:





These methyl radicals react with other chlorine free radicals to form methyl chloride along with the liberation of a chlorine free radical.



Hence, methyl free radicals and chlorine free radicals set up a chain reaction. While HCl and CH<sub>3</sub>Cl are the major products formed, other higher halogenated compounds are also formed as:



Step 3: Termination: Formation of ethane is a result of the termination of chain reactions taking place as a result of the consumption of reactants as:



Hence, by this process, ethane is obtained as a by-product of chlorination of methane.




Ques 13.2: Write IUPAC names of the following compounds:




(iii)




(iv)




(v)



(vi)




(vii)






Ans 13.2:

(i)



IUPAC name : 2-Methylbut-2-ene

(ii)




IUPAC name : Pent-1-ene-3-yne

(iii)




can be written as:




IUPAC name : 1, 3-Butadiene or Buta-1, 3-diene

(iv)



IUPAC name : 4-Phenyl but-1-ene


(v)



IUPAC name : 2-Methyl phenol




(vi)



IUPAC name : 5-(2- Methylpropyl)decane

(vii)



IUPAC name: 4-Ethyldeca-1, 5, 8-triene




**Ques 13.3:** For the following compounds, write structural formulas and IUPAC names for all possible isomers having the number of double or triple bond as indicated:

(i)  $\text{C}_4\text{H}_8$  (one double bond)

(ii)  $\text{C}_5\text{H}_8$  (one triple bond)

**Ans 13.3:**

(i) The following structural isomers are possible for  $\text{C}_4\text{H}_8$  with one double bond



The IUPAC name of




Compound (i) is But-1-ene,

Compound (ii) is But-2-ene, and

Compound (iii) is 2-Methylprop-1-ene.



- (ii) The following structural isomers are possible for  $C_5C_8$  with one triple bond:




The IUPAC name of  
Compound (i) is Pent-1-yne,  
Compound (ii) is Pent-2-yne, and  
Compound (iii) is 3-Methylbut-1-ene.

**Ques 13.4:** Write IUPAC names of the products obtained by the ozonolysis of the following compounds:

- (i) Pent-2-ene
- (ii) 3,4-Dimethyl-hept-3-ene
- (iii) 2-Ethylbut-1-ene
- (iv) 1-Phenylbut-1-ene

**Ans 13.4:**


- (i) Pent-2-ene undergoes ozonolysis as:



The IUPAC name of Product (I) is propanal and Product (II) is ethanal.




- (ii) 3, 4-Dimethylhept-3-ene undergoes ozonolysis as:



The IUPAC name of Product (I) is Pentane-2-one and Product (II) is Butane-2-one.


- (iii) 2-Ethylbut-1-ene undergoes ozonolysis as:



The IUPAC name of Product (I) is pentan-3-one and Product (II) is methanal.




(iv) 1-Phenylbut-1-ene undergoes ozonolysis as:



The IUPAC name of Product (I) is Propanal and Product (II) is Benzaldehyde.


**Ques 13.5:** An alkene 'A' on ozonolysis gives a mixture of ethanal and pentan-3-one. Write structure and IUPAC name of 'A'.

**Ans 13.5:**




During ozonolysis, an ozonide having a cyclic structure is formed as an intermediate which undergoes cleavage to give the final products. Ethanal and pentan-3-one are obtained from the intermediate ozonide.

Hence, the expected structure of the ozonide is:





This ozonide is formed as an addition of ozone to 'A'. The desired structure of 'A' can be obtained by the removal of ozone from the ozonide. Hence, the structural formula of 'A' is:




The IUPAC name of 'A' is 3-Ethylpent-2-ene.

**Ques 13.6:** An alkene 'A' contains three C – C, eight C – H  $\sigma$  bonds and one C – C  $\pi$  bond. 'A' on ozonolysis gives two moles of an aldehyde of molar mass 44 u. Write IUPAC name of 'A'.

**Ans 13.6:** As per the given information, 'A' on ozonolysis gives two moles of an aldehyde of molar mass 44 u. The formation of two moles of an aldehyde indicates the presence of identical structural units on both sides of the double bond containing carbon atoms.


Hence, the structure of 'A' can be represented as:



There are eight C–H  $\sigma$  bonds. Hence, there are 8 hydrogen atoms in 'A'. Also, there are three C–C bonds.

Hence, there are four carbon atoms present in the structure of 'A'.


Combining the inferences, the structure of 'A' can be represented as:



'A' has 3 C–C bonds, 8 C–H  $\sigma$  bonds, and one C–C  $\pi$  bond.

Hence, the IUPAC name of 'A' is But-2-ene.

Ozonolysis of 'A' takes place as:








The final product is ethanal with molecular mass =  $[(2 \times 12) + (4 \times 1) + (1 \times 16)]$   
= 44 u


**Ques 13.7:** Propanal and pentan-3-one are the ozonolysis products of an alkene?  
What is the structural formula of the alkene?

**Ans 13.7:** As per the given information, propanal and pentan-3-one are the ozonolysis products of an alkene.


Let the given alkene be 'A'. Writing the reverse of the ozonolysis reaction, we get:



The products are obtained on the cleavage of ozonide 'X'. Hence, 'X' contains both products in the cyclic form. The possible structure of ozonide can be represented as:



Now, 'X' is an addition product of alkene 'A' with ozone. Therefore, the possible structure of alkene 'A' is:





**Ques 13.8:** Write chemical equations for combustion reaction of the following hydrocarbons:

(i) Butane


(ii) Pentene

(iii) Hexyne


(iv) Toluene

**Ans 13.8:** Combustion can be defined as a reaction of a compound with oxygen.


(i)




(ii)



(iii)




(iv)






**Ques 13.9:** Draw the cis and trans structures of hex-2-ene. Which isomer will have higher b.p. and why?


**Ans 13.9:** Hex-2-ene is represented as:



Geometrical isomers of hex-2-ene are:



Cis - isomer



Trans - isomer

The dipole moment of cis-compound is a sum of the dipole moments of C-CH<sub>3</sub> and C-CH<sub>2</sub>CH<sub>3</sub> bonds acting in the same direction.


The dipole moment of trans-compound is the resultant of the dipole moments of C-CH<sub>3</sub> and C-CH<sub>2</sub>CH<sub>3</sub> bonds acting in opposite directions.

Hence, cis-isomer is more polar than trans-isomer. The higher the polarity, the greater is the intermolecular dipole-dipole interaction and the higher will be the boiling point.

Hence, cis-isomer will have a higher boiling point than trans-isomer.

**Ques 13.10:** Why is benzene extra ordinarily stable though it contains three double bonds?



**Ans 13.10:** Benzene is a hybrid of resonating structures given as:



All six carbon atoms in benzene are  $sp^2$  hybridized. The two  $sp^2$  hybrid orbitals of each carbon atom overlap with the  $sp^2$  hybrid orbitals of adjacent carbon atoms to form six sigma bonds in the hexagonal plane.

The remaining  $sp^2$  hybrid orbital on each carbon atom overlaps with the  $s$ -orbital of hydrogen to form six sigma C-H bonds.

The remaining unhybridized  $p$ -orbital of carbon atoms has the possibility of forming three  $\pi$  bonds by the lateral overlap of



The six  $\pi$ 's are delocalized and can move freely about the six carbon nuclei. Even after the presence of three double bonds, these delocalized  $\pi$ -electrons stabilize benzene.


**Ques 13.11: What are the necessary conditions for any system to be aromatic?**

**Ans 13.11:** A compound is said to be aromatic if it satisfies the following three conditions:

- It should have a planar structure.
- The  $\pi$ -electrons of the compound are completely delocalized in the ring.
- The total number of  $\pi$ -electrons present in the ring should be equal to  $(4n + 2)$ , where  $n = 0, 1, 2 \dots$  etc. This is known as Huckel's rule.

**Ques 13.12: Explain why the following systems are not aromatic?**

(i)



(ii)




(iii)





Ans 13.12:

(i)



For the given compound, the number of  $\pi$ -electrons is 6.

By Huckel's rule,

$$4n + 2 = 6$$

$$4n = 4 \quad n = 1$$

For a compound to be aromatic, the value of  $n$  must be an integer ( $n = 0, 1, 2, \dots$ ).

Since the value of  $n$  is an integer, the given compound is aromatic in nature.

(ii)



$sp^3$ -carbon

For the given compound, the number of  $\pi$ -electrons is 4.

By Huckel's rule,

$$4n + 2 = 4$$

$$4n = 2$$

$$n = \frac{1}{2}$$

For a compound to be aromatic, the value of  $n$  must be an integer ( $n = 0, 1, 2, \dots$ ), which is not true for the given compound. Hence, it is not aromatic in nature.

(iii)



Planar



Tub shaped

For the given compound, the number of  $\pi$ -electrons is 8.

By Huckel's rule,

$$4n + 2 = 8$$

$$4n = 6$$

$$n = \frac{3}{2}$$

For a compound to be aromatic, the value of  $n$  must be an integer ( $n = 0, 1, 2, \dots$ ).

Since the value of  $n$  is not an integer, the given compound is not aromatic in nature.




**Ques 13.13:** How will you convert benzene into


- (i) *p*-Nitrobromobenzene
- (ii) *m*-Nitrochlorobenzene
- (iii) *p*-Nitrotoluene
- (iv) Acetophenone

**Ans 13.13:**

- (i) Benzene can be converted into *p*-nitrobromobenzene as:




- (ii) Benzene can be converted into *m*-nitrochlorobenzene as:






(iii) Benzene can be converted into *p*-nitrotoluene as:




(iv) Benzene can be converted into acetophenone as:



**Ques 13.14:** In the alkane  $\text{H}_3\text{C}-\text{CH}_2-\text{C}(\text{CH}_3)_2-\text{CH}_2-\text{CH}(\text{CH}_3)_2$ , identify  $1^\circ, 2^\circ, 3^\circ$  carbon atoms and give the number of H atoms bonded to each one of these.

**Ans 13.14:**



$1^\circ$  carbon atoms are those which are bonded to only one carbon atom i.e., they have only one carbon atom as their neighbour.

The given structure has five  $1^\circ$  carbon atoms and fifteen hydrogen atoms attached to it.

$2^\circ$  carbon atoms are those which are bonded to two carbon atoms i.e., they have two carbon atoms as their neighbours.

The given structure has two  $2^\circ$  carbon atoms and four hydrogen atoms attached to it.

$3^\circ$  carbon atoms are those which are bonded to three carbon atoms i.e., they have three carbon atoms as their neighbours.

The given structure has one  $3^\circ$  carbon atom and only one hydrogen atom is attached to it.



**Ques 13.15:** What effect does branching of an alkane chain has on its boiling point?


**Ans 13.15:** Alkanes experience inter-molecular Van der Waals forces. The stronger the force, the greater will be the boiling point of the alkane.

As branching increases, the surface area of the molecule decreases which results in a small area of contact. As a result, the Van der Waals force also decreases which can be overcome at a relatively lower temperature.

Hence, the boiling point of an alkane chain decreases with an increase in branching.

**Ques 13.16:** Addition of HBr to propene yields 2-bromopropane, while in the presence of benzoyl peroxide, the same reaction yields 1-bromopropane. Explain and give mechanism.

**Ans 13.16:** Addition of HBr to propene is an example of an electrophilic substitution reaction. Hydrogen bromide provides an electrophile,  $H^+$ . This electrophile attacks the double bond to form  $1^\circ$  and  $2^\circ$  carbocations as shown:



Secondary carbocations are more stable than primary carbocations. Hence, the former predominates since it will form at a faster rate.

Thus, in the next step,  $\text{Br}^-$  attacks the carbocation to form 2-bromopropane as the major product.



This reaction follows Markovnikov's rule where the negative part of the addendum is attached to the carbon atom having a lesser number of hydrogen atoms.






# Class 11 Chemistry NCERT Solutions

$$\begin{array}{c} \text{C}_6\text{H}_5\text{CO}-\text{O}-\text{O}-\text{COC}_6\text{H}_5 \xrightarrow[\text{(Homolytic fission)}]{\Delta} 2 \text{C}_6\text{H}_5\text{CO}\cdot \xrightarrow{\Delta} 2 \text{C}_6\text{H}_5\cdot + \text{CO}_2 \\ \text{Benzoyl peroxide} \qquad \qquad \qquad \text{Benzoyl radical} \qquad \qquad \qquad \text{Phenyl radical} \\ \qquad \qquad \qquad \text{(unstable)} \\ \text{C}_6\text{H}_5 + \text{H}-\text{Br} \longrightarrow \text{C}_6\text{H}_5\cdot + \text{Br}\cdot \\ \text{Benzene} \end{array}$$

Thus, 1 – bromopropane is obtained as the major product.




In the presence of peroxide, Br free radical acts as an electrophile. Hence, two different products are obtained on addition of HBr to propene in the absence and presence of peroxide.

**Ques 13.17:** Write down the products of ozonolysis of 1, 2-dimethylbenzene (*o*-xylene). How does the result support Kekulé structure for benzene?

**Ans 13.17:**

*o*-xylene has two resonance structures:






All three products, i.e., methyl glyoxal, 1, 2-dimethylglyoxal, and glyoxal are obtained from two Kekule structures.

Since all three products cannot be obtained from any one of the two structures, this proves that *o*-xylene is a resonance hybrid of two Kekule structures (I and II).

**Ques 13.18:** Arrange benzene, *n*-hexane and ethyne in decreasing order of acidic behaviour. Also give reason for this behaviour.

**Ans 13.18:** Acidic character of a species is defined on the basis of ease with which it can lose its H<sup>+</sup> atoms.

The hybridization state of carbon in the given compound is:



As the *s*-character increases, the electronegativity of carbon increases and the electrons of C–H bond pair lie closer to the carbon atom.

As a result, partial positive charge of H-atom increases and H<sup>+</sup> ions are set free.

The *s*-character increases in the order:

$$\text{Sp}^3 < \text{sp}^2 < \text{sp}$$

Hence, the decreasing order of acidic behaviour is Ethyne > Benzene > Hexane.

**Ques 13.19:** Why does benzene undergo electrophilic substitution reactions easily and nucleophilic substitutions with difficulty?

**Ans 13.19:** Benzene is a planar molecule having delocalized electrons above and below the plane of ring. Hence, it is electron-rich. As a result, it is highly attractive to electron deficient species i.e., electrophiles.

Therefore, it undergoes electrophilic substitution reactions very easily. Nucleophiles are electron-rich.

Hence, they are repelled by benzene. Hence, benzene undergoes nucleophilic substitutions with difficulty.




# Class 11 Chemistry NCERT Solutions


(i) Ethyne  
(ii) Ethene  
(iii) Hexane

**Ans 13.20:**


(i) Benzene from Ethyne:



(ii) Benzene from Ethene:




(iii) Hexane to Benzene :






**Ques 13.21:** Write structures of all the alkenes which on hydrogenation give 2-methylbutane.

**Ans 13.21:** The basic skeleton of 2-methylbutane is shown below:




On the basis of this structure, various alkenes that will give 2-methylbutane on hydrogenation are:


(i)



(ii)



(iii)



**Ques 13.22:** Arrange the following set of compounds in order of their decreasing relative reactivity with an electrophile,  $\text{E}^+$

- (i) Chlorobenzene, 2,4-dinitrochlorobenzene, *p*-nitrochlorobenzene  
(ii) Toluene, *p*- $\text{H}_3\text{C}-\text{C}_6\text{H}_4-\text{NO}_2$ , *p*- $\text{O}_2\text{N}-\text{C}_6\text{H}_4-\text{NO}_2$ .


**Ans 13.22:** Electrophiles are reagents that participate in a reaction by accepting an electron pair in order to bond to nucleophiles.

The higher the electron density on a benzene ring, the more reactive is the compound towards an electrophile,  $\text{E}^+$  (Electrophilic reaction).



(i) The presence of an electron withdrawing group (i.e.,  $\text{NO}_2$  and  $\text{Cl}$ ) deactivates the aromatic ring by decreasing the electron density.


Since  $\text{NO}_2$  group is more electron withdrawing (due to resonance effect) than the  $\text{Cl}$  group (due to inductive effect), the decreasing order of reactivity is as follows:



(ii) While  $\text{CH}_3$  is an electron donating group,  $\text{NO}_2$  group is electron withdrawing. Hence, toluene will have the maximum electron density and is most easily attacked by  $\text{E}^+$ .

$\text{NO}_2$  is an electron withdrawing group.

Hence, when the number of  $\text{NO}_2$  substituents is greater, the order is as follows:




**Ques 13.23: Out of benzene, *m*-dinitrobenzene and toluene which will undergo nitration most easily and why?**

**Ans 13.23:** The ease of nitration depends on the presence of electron density on the compound to form nitrates. Nitration reactions are examples of electrophilic substitution reactions where an electron-rich species is attacked by a nitronium ion ( $\text{NO}_2^+$ ).

Now,  $\text{CH}_3$  group is electron donating and  $\text{NO}_2$  is electron withdrawing. Therefore, toluene will have the maximum electron density among the three compounds followed by benzene.

On the other hand, *m*-Dinitrobenzene will have the least electron density. Hence, it will undergo nitration with difficulty. Hence, the increasing order of nitration is as follows:



**Ques 13.24: Suggest the name of a Lewis acid other than anhydrous aluminium chloride which can be used during ethylation of benzene.**

**Ans 13.24:** The ethylation reaction of benzene involves the addition of an ethyl group on the benzene ring. Such a reaction is called a Friedel-Craft alkylation reaction. This reaction takes place in the presence of a Lewis acid.

Any Lewis acid like anhydrous  $\text{FeCl}_3$ ,  $\text{SnCl}_4$ ,  $\text{BF}_3$  etc. can be used during the ethylation of benzene.




# Class 11 Chemistry NCERT Solutions

**Ans 13.25:** Wurtz reaction is limited for the synthesis of symmetrical alkanes (alkanes with an even number of carbon atoms) In the reaction, two similar alkyl halides are taken as reactants and an alkane, containing double the number of carbon atoms, are formed.

$$\text{CH}_3-\text{Br} + 2\text{Na} + \text{Br}-\text{CH}_3 \xrightarrow{\text{Dry ether}} \text{CH}_3-\text{CH}_3 + 2\text{NaBr}$$

BromomethaneEthyne

Since the reaction involves free radical species, a side reaction also occurs to produce an alkene. For example, the reaction of bromomethane and iodoethane gives a mixture of alkanes.



The boiling points of alkanes (obtained in the mixture) are very close. Hence, it becomes difficult to separate them.