



Chapter – 15

Plant Growth & Development

NCERT Back Exercises:

Ques 1: Define growth, differentiation, development, dedifferentiation, redifferentiation, determinate growth, meristem and growth rate.

Ans 1:

- (i) Growth: It is an irreversible and permanent process, accomplished by an increase in the size of an organ or organ parts or even of an individual cell.
- (ii) Differentiation: It is a process in which the cells derived from the apical meristem (root and shoot apex) and the cambium undergo structural changes in the cell wall and the protoplasm, becoming mature to perform specific functions.
- (iii) Development: It refers to the various changes occurring in an organism during its life cycle – from the germination of seeds to senescence.
- (iv) De-differentiation: It is the process in which permanent plant cells regain the power to divide under certain conditions.
- (v) Re-differentiation: It is the process in which de-differentiated cells become mature again and lose their capacity to divide.
- (vi) Determinate growth: It refers to limited growth. For example, animals and plant leaves stop growing after having reached maturity.
- (vii) Meristem: In plants, growth is restricted to specialised regions where active cell divisions take place. Such a region is called meristem. There are three types of meristems – apical meristem, lateral meristem, and intercalary meristem.
- (viii) Growth rate: It can be defined as the increased growth in plants per unit time.



Ques 2: Why is not any one parameter good enough to demonstrate growth throughout the life of a flowering plant?


Ans 2: In plants, growth is said to have taken place when the amount of protoplasm increases. Measuring the growth of protoplasm involves many parameters such as the weight of the fresh tissue sample, the weight of the dry tissue sample, the differences in length, area, volume, and cell number measured during the growth period. Measuring the growth of plants using only one parameter does not provide enough information and hence, is insufficient for demonstrating growth.

Ques 3: Describe briefly:

- (i) Arithmetic growth
- (ii) Geometric growth
- (iii) Sigmoid growth curve
- (iv) Absolute and relative growth rates


Ans 3:

- (i) Arithmetic growth: In arithmetic growth, one of the daughter cells continues to divide, while the other differentiates into maturity. The elongation of roots at a constant rate is an example of arithmetic growth.






- (ii) Geometric growth: Geometric growth is characterised by a slow growth in the initial stages and a rapid growth during the later stages. The daughter cells derived from mitosis retain the ability to divide, but slow down because of a limited nutrient supply.



- (iii) Sigmoid growth curve: The growth of living organisms in their natural environment is characterised by an S shaped curve called sigmoid growth curve. This curve is divided into three phases – lag phase, log phase or exponential phase of rapid growth, and stationary phase.



- (iv) Absolute and relative growth rates: Absolute growth rate refers to the measurement and comparison of total growth per unit time.

Relative growth rate refers to the growth of a particular system per unit time, expressed on a common basis.



Ques 4: List five main groups of natural plant growth regulators. Write a note on discovery, physiological functions and agricultural/horticultural applications of any one of them.

Ans 4: Plant growth regulators are the chemical molecules secreted by plants affecting the physiological attributes of a plant. There are five main plant growth regulators. These are:

(i) Auxins

Discovery:

The first observations regarding the effects of auxins were made by Charles Darwin and Francis Darwin wherein they saw the coleoptiles of canary grass bending toward a unilateral source of light.

It was concluded after a series of experiments that some substance produced at the tip of coleoptiles was responsible for the bending. Finally, this substance was extracted as auxins from the tips of coleoptiles in oat seedlings.

Physiological functions:

- (i) They control plant cell-growth.
- (ii) They cause the phenomenon of apical dominance.
- (iii) They control division in the vascular cambium and xylem differentiation.
- (iv) They induce parthenocarpy and prevent abscission of leaves and fruits.

Horticultural applications:

- (i) They are used as the rooting hormones in stem cuttings.
- (ii) 2-4 D is used as a weedicide to kill broadleaf, dicotyledonous weeds.
- (iii) They induce parthenocarpy in tomatoes.
- (iv) They promote flowering in pineapple and litchi.

(ii) Gibberellic acid

Discovery:

Bakane or the “foolish rice seedling” disease was first observed by Japanese farmers. In this disease, rice seedlings appear to grow taller than natural plants, and become slender and pale green. Later, after several experiments, it was found that this condition was caused by the infection from a certain fungus *Gibberella fujikuroi*. The active substance was isolated and identified as gibberellic acid.

Physiological functions:

- (i) It causes elongation of internodes.
- (ii) It promotes bolting in rosette plants.
- (iii) It helps in inducing seed germination by breaking seed dormancy and initiating the synthesis of hydrolases enzymes for digesting reserve food.



Horticultural applications:

- (i) It helps in increasing the sugar content in sugarcane by increasing the length of the internodes.
- (ii) It increases the length of grape stalks.
- (iii) It improves the shape of apple.
- (iv) It delays senescence.
- (v) It hastens maturity and induces seed-production in juvenile conifers.

(iii) Cytokinins

Discovery:

Through their experimental observations, F. Skoog and his co-workers found that the tobacco callus differentiated when extracts of vascular tissues, yeast extract, coconut milk, or DNA were added to the culture medium. This led to the discovery of cytokinins.

Physiological functions:

- (i) They promote the growth of lateral branches by inhibiting apical dominance.
- (ii) They help in the production of new leaves, chloroplasts, and adventitious shoots.
- (iii) They help in delaying senescence by promoting nutrient mobilisation.

Horticultural applications:

- (i) They are used for preventing apical dominance.
- (ii) They are used for delaying senescence in leaves.

(iv) Ethylene

Discovery:

It was observed that unripe bananas ripened faster when stored with ripe bananas. Later, the substance promoting the ripening was found to be ethylene.

Physiological functions:

- (i) It helps in breaking seed and bud dormancy.
- (ii) It promotes rapid internode-elongation in deep-water rice plants.
- (iii) It promotes root-growth and formation of root hairs.
- (iv) It promotes senescence and abscission of leaves and flowers.
- (v) It hastens the respiration rate in fruits and enhances fruit ripening.

Horticultural applications:

- (i) It is used to initiate flowering and synchronising the fruit set in pineapples.
- (ii) It induces flowering in mango.
- (iii) Ethephon is used to ripen the fruits in tomatoes and apples, and accelerate the abscission of flowers and leaves in cotton, cherry, and walnut.
- (iv) It promotes the number of female flowers in cucumbers.



◆ **(v) Absciscic acid** ◆

Discovery:

During the mid-1960s, inhibitor-B, abscission II, and dormin were discovered by three independent researchers. These were later on found to be chemically similar and were thereafter called ABA (Absciscic acid).

Physiological functions:

- (i) It acts as an inhibitor to plant metabolism.
- (ii) It stimulates stomatal closure during water stress.
- (iii) It induces seed dormancy.
- (iv) It induces abscission of leaves, fruits, and flowers.

Horticultural application:

- (i) It induces seed dormancy in stored seeds.

Ques 5: What do you understand by photoperiodism and vernalisation? Describe their significance.

Ans 5: Photoperiodism can be termed as the response of plants to periods of day/night. It is theorized that the hormonal substance that is responsible for flowering, is formed in the leaves which subsequently migrates to the shoot apices and alters them into flowering apices. This process of photoperiodism helps in studying the response of flowering in different crop plants when the duration of exposure of light is considered.

Vernalisation is the phenomena where the process of flowering in some plants is either quantitatively or qualitatively dependent on the exposure to lower temperatures. In particular, it refers to promoting the flowering process by a period of lower temperatures. The process prevents precocious reproductive development late in the growing season which thereby enables the plant to have sufficient time to attain maturity.

Ques 6: Why is Absciscic acid also known as stress hormone?

Ans 6: Absciscic acid is called stress hormones as it induces various responses in plants against stress conditions.

It increases the tolerance of plants toward various stresses. It induces the closure of the stomata during water stress. It promotes seed dormancy and ensures seed germination during favourable conditions. It helps seeds withstand desiccation. It also helps in inducing dormancy in plants at the end of the growing season and promotes abscission of leaves, fruits, and flowers.



Ques 7: 'Both growth and differentiation in higher plants are *open*'. Comment.


Ans 7: Growth and development in higher plants is referred to as being *open*. This is because various meristems, having the capacity for continuously dividing and producing new cells, are present at different locations in these plant bodies.

Ques 8: 'Both a short day plant and a long day plant can flower simultaneously in a given place'. Explain.

Ans 8: The flowering response in short-day plants and long-day plants is dependent on the durations for which these plants are exposed to light. The short-day plant and long day plant can flower at the same place, provided they have been given an adequate photoperiod.

Ques 9: Which one of the plant growth regulators would you use if you are asked to:

- (i) Induce rooting in a twig
- (ii) Quickly ripen a fruit
- (iii) Delay leaf senescence
- (iv) Induce growth in axillary buds
- (v) 'Bolt' a rosette plant
- (vi) Induce immediate stomatal closure in leaves.



Ans 9:

- | | |
|--|------------------|
| (i) Induce rooting in a twig | Auxins |
| (ii) Quickly ripen a fruit | Ethylene |
| (iii) Delay leaf senescence | Cytokinins |
| (iv) Induce growth in axillary buds | Cytokinins |
| (v) 'Bolt' a rosette plant | Gibberellic acid |
| (vi) Induce immediate stomatal closure in leaves | Absciscic acid |



Ques 10: Would a defoliated plant respond to photoperiodic cycle? Why?

Ans 10: A defoliated plant will not respond to the photoperiodic cycle.

It is hypothesised that the hormonal substance responsible for flowering is formed in the leaves, subsequently migrating to the shoot apices and modifying them into flowering apices. Therefore, in the absence of leaves, light perception would not occur, i.e., the plant would not respond to light.

Ques 11: What would be expected to happen if:

- (i) GA_3 is applied to rice seedlings
- (ii) Dividing cells stop differentiating
- (iii) A rotten fruit gets mixed with unripe fruits
- (iv) You forget to add cytokinin to the culture medium.

Ans 11:

- (i) If GA_3 is applied to rice seedlings, then the rice seedlings will exhibit internode elongation and increase in height.
- (ii) If dividing cells stop differentiating, then the plant organs such as leaves and stem will not be formed. The mass of undifferentiated cell is called callus.
- (iii) If a rotten fruit gets mixed with unripe fruits, then the ethylene produced from the rotten fruits will hasten the ripening of the unripe fruits.
- (iv) If you forget to add cytokinin to the culture medium, then cell division, growth, and differentiation will not be observed.