

Assignment 2

AI1110: Probability and Random Variables
Indian Institute of Technology Hyderabad

Yasir Usmani
AI22BTECH11031

12.13.6.18: Question. Consider the experiment of throwing a die, if a multiple of 3 comes up, throw the die again and if any other number comes, toss a coin. Find the conditional probability of the event ‘the coin shows a tail’, given that ‘at least one die shows a 3’.

Answer: 0.5

Solution: Using Markov Chain Approach,
Let the states **S** be,

- 1) 0 : The initial state.
- 2) 1 : 3 comes on die throw.
- 3) 2 : 6 comes on die throw.
- 4) 3 : non-multiple of 3 comes on die throw.
- 5) 4 : Heads comes up on coin toss.
- 6) 5 : Tails comes up on coin toss.

The Markov Chain for the given states is as follows,




Figure: Markov Chain

- Here, **States 4 and 5** are **Absorbing States** indicating that once they are reached, They cannot be left. While **States 0, 1, 2, 3** are **Transient States** whose probability will eventually become 0.

The State Transition Matrix is as follows,

$$\mathbf{A} = \begin{pmatrix} 0 & \frac{1}{6} & \frac{1}{6} & \frac{4}{6} & 0 & 0 \\ 0 & \frac{1}{6} & \frac{1}{6} & \frac{4}{6} & 0 & 0 \\ 0 & \frac{1}{6} & \frac{1}{6} & \frac{4}{6} & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \quad (1)$$

Now, Let the Probability of state i at time t be $P_i^{(t)}$. Then the state vector will be:

$$\mathbf{B}_t = (P_0^{(t)} \quad P_1^{(t)} \quad P_2^{(t)} \quad P_3^{(t)} \quad P_4^{(t)} \quad P_5^{(t)}) \quad (2)$$

From here we can write probability of states one step in time after t as,

$$P_0^{t+1} = 0 \quad (3)$$

$$P_1^{t+1} = \frac{1}{6} \times P_0^t + \frac{1}{6} \times P_1^t + \frac{1}{6} \times P_2^t \quad (4)$$

$$P_2^{t+1} = \frac{1}{6} \times P_0^t + \frac{1}{6} \times P_1^t + \frac{1}{6} \times P_2^t \quad (5)$$

$$P_3^{t+1} = \frac{4}{6} \times P_0^t + \frac{4}{6} \times P_1^t + \frac{4}{6} \times P_2^t \quad (6)$$

$$P_4^{t+1} = \frac{1}{2} \times P_3^t + 1 \times P_4^t \quad (7)$$

$$P_5^{t+1} = \frac{1}{2} \times P_3^t + 1 \times P_5^t \quad (8)$$

- These equations can be written directly as:

$$\mathbf{B}_{t+1} = \mathbf{B}_t \times \mathbf{A} \quad (9)$$

To Find,

$$\lim_{t \rightarrow \infty} P_5^{(t)} \quad (10)$$

From (9),

$$\mathbf{B}_t = \mathbf{B}_{t-1} \times \mathbf{A} = \mathbf{B}_{t-2} \times \mathbf{A}^2 \cdots = \mathbf{B}_0 \times \mathbf{A}^t \quad (11)$$

The Initial State \mathbf{B}_0 can be denoted by given that 3 occurs atleast once,

$$\mathbf{B}_0 = (0 \quad 1 \quad 0 \quad 0 \quad 0 \quad 0) \quad (12)$$

The last column of \mathbf{A}^t when $t \rightarrow \infty$ can be written as,

$$\mathbf{A}^t = \begin{pmatrix} \cdot & \cdot & \cdot & \cdot & \cdot & \frac{1}{2} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \frac{1}{2} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \frac{1}{2} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \frac{1}{2} \\ \cdot & \cdot & \cdot & \cdot & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1 \end{pmatrix} \quad (13)$$

$$\lim_{t \rightarrow \infty} \mathbf{B}_0 = (\cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad 0.5) \quad (14)$$

Hence From (2),

$$\implies \lim_{t \rightarrow \infty} P_5^{(t)} = 0.5 \quad (15)$$