

Near-optimal Quantum Algorithms for Multivariate Mean Estimation

arXiv:2111.09787

Arjan Cornelissen¹, Yassine Hamoudi², Sofiene Jerbi³

¹QuSoft, University of Amsterdam

²Simons Institute for the Theory of Computing, University of California, Berkeley

³Institute for Theoretical Physics, University of Innsbruck

June 20th, 2022



Problem statement

Problem statement

We have

- 1 A probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
- 2 A random variable $X : \Omega \rightarrow \mathbb{R}^d$.

Problem statement

We have

- 1 A probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
- 2 A random variable $X : \Omega \rightarrow \mathbb{R}^d$.

Properties:

- 1 *Mean:*

$$\mu = \mathbb{E}[X] = \sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega) \in \mathbb{R}^d.$$

Problem statement

We have

- 1 A probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
- 2 A random variable $X : \Omega \rightarrow \mathbb{R}^d$.

Properties:

- 1 *Mean:*

$$\mu = \mathbb{E}[X] = \sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega) \in \mathbb{R}^d.$$

- 2 *Covariance matrix:*

$$\Sigma = \begin{bmatrix} \text{Var}[X_1] & \text{Cov}[X_1, X_2] & \cdots & \text{Cov}[X_1, X_d] \\ \text{Cov}[X_1, X_2] & \text{Var}[X_2] & \cdots & \text{Cov}[X_2, X_d] \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}[X_1, X_d] & \text{Cov}[X_2, X_d] & \cdots & \text{Var}[X_d] \end{bmatrix}$$

Problem statement

We have

- 1 A probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
- 2 A random variable $X : \Omega \rightarrow \mathbb{R}^d$.

Properties:

- 1 *Mean:*

$$\mu = \mathbb{E}[X] = \sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega) \in \mathbb{R}^d.$$

- 2 *Covariance matrix:*

$$\Sigma = \begin{bmatrix} \text{Var}[X_1] & \text{Cov}[X_1, X_2] & \cdots & \text{Cov}[X_1, X_d] \\ \text{Cov}[X_1, X_2] & \text{Var}[X_2] & \cdots & \text{Cov}[X_2, X_d] \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}[X_1, X_d] & \text{Cov}[X_2, X_d] & \cdots & \text{Var}[X_d] \end{bmatrix}$$

Multivariate mean estimation:

- 1 *Goal:* Approximate $\mu \in \mathbb{R}^d$.

Problem statement

We have

- 1 A probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
- 2 A random variable $X : \Omega \rightarrow \mathbb{R}^d$.

Properties:

- 1 *Mean:*

$$\mu = \mathbb{E}[X] = \sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega) \in \mathbb{R}^d.$$

- 2 *Covariance matrix:*

$$\Sigma = \begin{bmatrix} \text{Var}[X_1] & \text{Cov}[X_1, X_2] & \cdots & \text{Cov}[X_1, X_d] \\ \text{Cov}[X_1, X_2] & \text{Var}[X_2] & \cdots & \text{Cov}[X_2, X_d] \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}[X_1, X_d] & \text{Cov}[X_2, X_d] & \cdots & \text{Var}[X_d] \end{bmatrix}$$

Multivariate mean estimation:

- 1 *Goal:* Approximate $\mu \in \mathbb{R}^d$.
- 2 *Assumption:*

$$\text{Tr}[\Sigma] = \sum_{j=1}^d \text{Var}[X_j] < \infty.$$

Problem statement

We have

- 1 A probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
- 2 A random variable $X : \Omega \rightarrow \mathbb{R}^d$.

Properties:

- 1 *Mean:*

$$\mu = \mathbb{E}[X] = \sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega) \in \mathbb{R}^d.$$

- 2 *Covariance matrix:*

$$\Sigma = \begin{bmatrix} \text{Var}[X_1] & \text{Cov}[X_1, X_2] & \cdots & \text{Cov}[X_1, X_d] \\ \text{Cov}[X_1, X_2] & \text{Var}[X_2] & \cdots & \text{Cov}[X_2, X_d] \\ \vdots & \vdots & \ddots & \vdots \\ \text{Cov}[X_1, X_d] & \text{Cov}[X_2, X_d] & \cdots & \text{Var}[X_d] \end{bmatrix}$$

Multivariate mean estimation:

- 1 *Goal:* Approximate $\mu \in \mathbb{R}^d$.
- 2 *Assumption:*

$$\text{Tr}[\Sigma] = \sum_{j=1}^d \text{Var}[X_j] < \infty.$$

Applications:

- 1 Physics/chemistry simulations
- 2 Computer graphics
- 3 Finance
- 4 Shadow tomography

Access models

We have

- ① A probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
- ② A random variable $X : \Omega \rightarrow \mathbb{R}^d$.

We want to approximate μ .

Access models

We have

- ① A probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
- ② A random variable $X : \Omega \rightarrow \mathbb{R}^d$.

We want to approximate μ .

Classical access model:

- ① Obtain outcome $\omega \sim \mathbb{P}$.
- ② Function $\omega \mapsto X(\omega)$.

Access models

We have

- ① A probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
- ② A random variable $X : \Omega \rightarrow \mathbb{R}^d$.

We want to approximate μ .

Classical access model:

- ① Obtain outcome $\omega \sim \mathbb{P}$.
- ② Function $\omega \mapsto X(\omega)$.

Quantum access model:

- ① Distribution oracle:

$$|0\rangle \mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle.$$

Access models

We have

- ① A probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
- ② A random variable $X : \Omega \rightarrow \mathbb{R}^d$.

We want to approximate μ .

Classical access model:

- ① Obtain outcome $\omega \sim \mathbb{P}$.
- ② Function $\omega \mapsto X(\omega)$.

Quantum access model:

- ① Distribution oracle:

$$|0\rangle \mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle.$$

- ② Random variable oracle:

$$|\omega\rangle |0\rangle \mapsto |\omega\rangle |X(\omega)\rangle.$$

Access models

We have

- ① A probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
- ② A random variable $X : \Omega \rightarrow \mathbb{R}^d$.

We want to approximate μ .

Classical access model:

- ① Obtain outcome $\omega \sim \mathbb{P}$.
- ② Function $\omega \mapsto X(\omega)$.

Quantum access model:

- ① Distribution oracle:

$$|0\rangle \mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle.$$

- ② Random variable oracle:

$$|\omega\rangle |0\rangle \mapsto |\omega\rangle |X(\omega)\rangle.$$

Think of

$$|X(\omega)\rangle = |X(\omega)_1\rangle \otimes |X(\omega)_2\rangle \otimes \cdots \otimes |X(\omega)_d\rangle$$

Access models

We have

- ① A probability space $(\Omega, \mathcal{F}, \mathbb{P})$,
- ② A random variable $X : \Omega \rightarrow \mathbb{R}^d$.

We want to approximate μ .

Classical access model:

- ① Obtain outcome $\omega \sim \mathbb{P}$.
- ② Function $\omega \mapsto X(\omega)$.

Quantum access model:

- ① Distribution oracle:

$$|0\rangle \mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle.$$

- ② Random variable oracle:

$$|\omega\rangle |0\rangle \mapsto |\omega\rangle |X(\omega)\rangle.$$

Think of

$$|X(\omega)\rangle = |X(\omega)_1\rangle \otimes |X(\omega)_2\rangle \otimes \cdots \otimes |X(\omega)_d\rangle$$


Calls to these routines are *samples*.

Results

Results

Goal: Construct estimator $\tilde{\mu}$, using n samples, s.t.


$$\mathbb{P}[\|\mu - \tilde{\mu}\|_2 \leq \varepsilon(n)] \geq \frac{2}{3}.$$



Results

Goal: Construct estimator $\tilde{\mu}$, using n samples, s.t.

$$\mathbb{P}[\|\mu - \tilde{\mu}\|_2 \leq \varepsilon(n)] \geq \frac{2}{3}.$$




		$\varepsilon(n)$	Remarks
Classically	$d = 1$		
	$d \geq 1$		
Quantumly	$d = 1$		
	$d \geq 1$		

Results

Goal: Construct estimator $\tilde{\mu}$, using n samples, s.t.

$$\mathbb{P}[\|\mu - \tilde{\mu}\|_2 \leq \varepsilon(n)] \geq \frac{2}{3}.$$




		$\varepsilon(n)$	Remarks
Classically	$d = 1$	$\Theta\left(\sqrt{\frac{\text{Var}[X]}{n}}\right)$	Monte Carlo sampling
	$d \geq 1$	$\Theta\left(\sqrt{\frac{\text{Tr}[\Sigma]}{n}}\right)$	Monte Carlo sampling
Quantumly	$d = 1$		
	$d \geq 1$		

Results

Goal: Construct estimator $\tilde{\mu}$, using n samples, s.t.

$$\mathbb{P}[\|\mu - \tilde{\mu}\|_2 \leq \varepsilon(n)] \geq \frac{2}{3}.$$




		$\varepsilon(n)$	Remarks
Classically	$d = 1$	$\Theta\left(\sqrt{\frac{\text{Var}[X]}{n}}\right)$	Monte Carlo sampling
	$d \geq 1$	$\Theta\left(\sqrt{\frac{\text{Tr}[\Sigma]}{n}}\right)$	Monte Carlo sampling
Quantumly	$d = 1$	$\tilde{\Theta}\left(\frac{\sqrt{\text{Var}[X]}}{n}\right)$	Known $\text{Var}[X]$ [Hei02; Mon15; HM19] Unknown $\text{Var}[X]$ [Ham21]
	$d \geq 1$		

Results

Goal: Construct estimator $\tilde{\mu}$, using n samples, s.t.

$$\mathbb{P}[\|\mu - \tilde{\mu}\|_2 \leq \varepsilon(n)] \geq \frac{2}{3}.$$




	$\varepsilon(n)$	Remarks
Classically	$d = 1$	$\Theta\left(\sqrt{\frac{\text{Var}[X]}{n}}\right)$ Monte Carlo sampling
	$d \geq 1$	$\Theta\left(\sqrt{\frac{\text{Tr}[\Sigma]}{n}}\right)$ Monte Carlo sampling
Quantumly	$d = 1$	$\tilde{\Theta}\left(\frac{\sqrt{\text{Var}[X]}}{n}\right)$ Known $\text{Var}[X]$ [Hei02; Mon15; HM19] Unknown $\text{Var}[X]$ [Ham21]
	$d \geq 1$	$\tilde{\Theta}\left(\begin{cases} \frac{\sqrt{d \text{Tr}[\Sigma]}}{n}, & \text{if } n \geq d \\ \sqrt{\frac{\text{Tr}[\Sigma]}{n}}, & \text{if } n < d \end{cases}\right)$ <i>Our work</i>


Results

Goal: Construct estimator $\tilde{\mu}$, using n samples, s.t.

$$\mathbb{P}[\|\mu - \tilde{\mu}\|_2 \leq \varepsilon(n)] \geq \frac{2}{3}.$$



	$\varepsilon(n)$	Remarks
Classically	$d = 1$	$\Theta\left(\sqrt{\frac{\text{Var}[X]}{n}}\right)$ Monte Carlo sampling
	$d \geq 1$	$\Theta\left(\sqrt{\frac{\text{Tr}[\Sigma]}{n}}\right)$ Monte Carlo sampling
Quantumly	$d = 1$	$\tilde{\Theta}\left(\frac{\sqrt{\text{Var}[X]}}{n}\right)$ Known Var[X] [Hei02; Mon15; HM19] Unknown Var[X] [Ham21]
	$d \geq 1$	$\tilde{\Theta}\left(\begin{cases} \frac{\sqrt{d \text{Tr}[\Sigma]}}{n}, & \text{if } n \geq d \\ \sqrt{\frac{\text{Tr}[\Sigma]}{n}}, & \text{if } n < d \end{cases}\right)$ <i>Our work</i>




Crucial observation: quantum speed-up only when $n \geq d$.

Results

Goal: Construct estimator $\tilde{\mu}$, using n samples, s.t.

$$\mathbb{P}[\|\mu - \tilde{\mu}\|_2 \leq \varepsilon(n)] \geq \frac{2}{3}.$$

		$\varepsilon(n)$	Remarks
Classically	$d = 1$	$\Theta\left(\sqrt{\frac{\text{Var}[X]}{n}}\right)$	Monte Carlo sampling
	$d \geq 1$	$\Theta\left(\sqrt{\frac{\text{Tr}[\Sigma]}{n}}\right)$	Monte Carlo sampling
Quantumly	$d = 1$	$\tilde{\Theta}\left(\sqrt{\frac{\text{Var}[X]}{n}}\right)$	Known $\text{Var}[X]$ [Hei02; Mon15; HM19] Unknown $\text{Var}[X]$ [Ham21]
	$d \geq 1$	$\tilde{\Theta}\left(\begin{cases} \sqrt{\frac{d \text{Tr}[\Sigma]}{n}}, & \text{if } n \geq d \\ \sqrt{\frac{\text{Tr}[\Sigma]}{n}}, & \text{if } n < d \end{cases}\right)$	<i>Our work</i>



Crucial observation: quantum speed-up only when $n \geq d$.

Quantum algorithm outline

Quantum algorithm outline

Goal: Estimate $\mu = \mathbb{E}[X] \in \mathbb{R}^d$.

μ

Quantum algorithm outline

Goal: Estimate $\mu = \mathbb{E}[X] \in \mathbb{R}^d$.

① *Get a crude estimate:* $\bar{\mu}$ s.t.

$$\|\mu - \bar{\mu}\|_2 \leq \sqrt{\text{Tr}[\Sigma]},$$

using $O(1)$ samples.

$$\begin{array}{c} \bullet \\ \mu \end{array} \quad \begin{array}{c} \bullet \\ \bar{\mu} \end{array}$$

Quantum algorithm outline

Goal: Estimate $\mu = \mathbb{E}[X] \in \mathbb{R}^d$.

- 1 *Get a crude estimate:* $\bar{\mu}$ s.t.

$$\|\mu - \bar{\mu}\|_2 \leq \sqrt{\text{Tr}[\Sigma]},$$

using $O(1)$ samples.

- 2 *Get an idea of the spread:*


$$\begin{array}{c} \bullet \\ \mu \end{array} \quad \begin{array}{c} \bullet \\ \bar{\mu} \end{array}$$

Quantum algorithm outline


Goal: Estimate $\mu = \mathbb{E}[X] \in \mathbb{R}^d$.

- 1 *Get a crude estimate:* $\bar{\mu}$ s.t.

$$\|\mu - \bar{\mu}\|_2 \leq \sqrt{\text{Tr}[\Sigma]},$$

using $O(1)$ samples.

- 2 *Get an idea of the spread:*



Quantum algorithm outline


Goal: Estimate $\mu = \mathbb{E}[X] \in \mathbb{R}^d$.

- 1 *Get a crude estimate:* $\bar{\mu}$ s.t.

$$\|\mu - \bar{\mu}\|_2 \leq \sqrt{\text{Tr}[\Sigma]},$$

using $O(1)$ samples.

- 2 *Get an idea of the spread:*



Quantum algorithm outline


Goal: Estimate $\mu = \mathbb{E}[X] \in \mathbb{R}^d$.

- 1 *Get a crude estimate:* $\bar{\mu}$ s.t.

$$\|\mu - \bar{\mu}\|_2 \leq \sqrt{\text{Tr}[\Sigma]},$$

using $O(1)$ samples.

- 2 *Get an idea of the spread:*



Quantum algorithm outline

Goal: Estimate $\mu = \mathbb{E}[X] \in \mathbb{R}^d$.

- ① *Get a crude estimate:* $\bar{\mu}$ s.t.


$$\|\mu - \bar{\mu}\|_2 \leq \sqrt{\text{Tr}[\Sigma]},$$

using $O(1)$ samples.

- ② *Get an idea of the spread:*
Estimate quantiles a_ℓ s.t.

$$\mathbb{P}[\|X - \bar{\mu}\|_2 \geq a_\ell] \approx \frac{1}{2^\ell},$$

for $\ell \in \{1, \dots, 2 \log(n)\}$.



Quantum algorithm outline

Goal: Estimate $\mu = \mathbb{E}[X] \in \mathbb{R}^d$.

- ① *Get a crude estimate:* $\bar{\mu}$ s.t.

$$\|\mu - \bar{\mu}\|_2 \leq \sqrt{\text{Tr}[\Sigma]},$$

using $O(1)$ samples.


- ② *Get an idea of the spread:*
Estimate quantiles a_ℓ s.t.

$$\mathbb{P}[\|X - \bar{\mu}\|_2 \geq a_\ell] \approx \frac{1}{2^\ell},$$

for $\ell \in \{1, \dots, 2 \log(n)\}$.

- ③ *Estimate truncated mean on every ring:*

$$\mathbb{E}[X] \approx \sum_{\ell=1}^k \mathbb{E}[X \cdot \mathbb{1}_{X \in R_\ell}].$$



Quantile estimation

Quantile estimation


We let $Y = \|X - \bar{\mu}\|_2$.

Want to find a_ℓ s.t. $\mathbb{P}[Y \geq a_\ell] \approx \frac{1}{2^\ell}$

Quantile estimation

We let $Y = \|X - \bar{\mu}\|_2$.

Want to find a_ℓ s.t. $\mathbb{P}[Y \geq a_\ell] \approx \frac{1}{2^\ell}$




Quantile estimation

We let $Y = \|X - \bar{\mu}\|_2$.

1. $\left| \begin{array}{l} \tilde{O}(1) \text{ samples from } Y \end{array} \right.$

Want to find a_ℓ s.t. $\mathbb{P}[Y \geq a_\ell] \approx \frac{1}{2^\ell}$

\hookrightarrow Let a_1 be the median
 $\mathbb{P}[Y \geq a_1] \approx \frac{1}{2}$




Quantile estimation

We let $Y = \|X - \bar{\mu}\|_2$.

1. $\left| \begin{array}{l} \tilde{O}(1) \text{ samples from } Y \end{array} \right.$

Want to find a_ℓ s.t. $\mathbb{P}[Y \geq a_\ell] \approx \frac{1}{2^\ell}$

\hookrightarrow Let a_1 be the median
 $\mathbb{P}[Y \geq a_1] \approx \frac{1}{2}$




Quantile estimation

We let $Y = \|X - \bar{\mu}\|_2$.

1. $\tilde{O}(1)$ samples from Y

Want to find a_ℓ s.t. $\mathbb{P}[Y \geq a_\ell] \approx \frac{1}{2^\ell}$

\hookrightarrow Let a_1 be the median
 $\mathbb{P}[Y \geq a_1] \approx \frac{1}{2}$




Quantile estimation

We let $Y = \|X - \bar{\mu}\|_2$.

1. $\left| \begin{array}{l} \tilde{O}(1) \text{ samples from } Y \end{array} \right.$

Want to find a_ℓ s.t. $\mathbb{P}[Y \geq a_\ell] \approx \frac{1}{2^\ell}$

\hookrightarrow Let a_1 be the median
 $\mathbb{P}[Y \geq a_1] \approx \frac{1}{2}$




Quantile estimation

We let $Y = \|X - \bar{\mu}\|_2$.

1. $\tilde{O}(1)$ samples from Y

Want to find a_ℓ s.t. $\mathbb{P}[Y \geq a_\ell] \approx \frac{1}{2^\ell}$

\hookrightarrow Let a_1 be the median
 $\mathbb{P}[Y \geq a_1] \approx \frac{1}{2}$




Quantile estimation

We let $Y = \|X - \bar{\mu}\|_2$.

1. $\left| \begin{array}{l} \tilde{O}(1) \text{ samples from } Y \end{array} \right.$

Want to find a_ℓ s.t. $\mathbb{P}[Y \geq a_\ell] \approx \frac{1}{2^\ell}$

\hookrightarrow Let a_1 be the median
 $\mathbb{P}[Y \geq a_1] \approx \frac{1}{2}$



Quantile estimation

We let $Y = \|X - \bar{\mu}\|_2$.


Want to find a_ℓ s.t. $\mathbb{P}[Y \geq a_\ell] \approx \frac{1}{2^\ell}$

1. $\tilde{O}(1)$ samples from Y
- \vdots
- \vdots
- ℓ . $\tilde{O}(1)$ samples from $Y|Y \geq a_{\ell-1}$
Requires $\tilde{O}(\sqrt{2^{\ell-1}})$ samples from Y
by *amplitude amplification*

\hookrightarrow Let a_1 be the median
 $\mathbb{P}[Y \geq a_1] \approx \frac{1}{2}$

\vdots

\hookrightarrow Let a_ℓ be the median
 $\mathbb{P}[Y \geq a_\ell] \approx \frac{1}{2^\ell}$




Quantile estimation

We let $Y = \|X - \bar{\mu}\|_2$.

Want to find a_ℓ s.t. $\mathbb{P}[Y \geq a_\ell] \approx \frac{1}{2^\ell}$

1.	$\tilde{O}(1)$ samples from Y	\hookrightarrow	Let a_1 be the median $\mathbb{P}[Y \geq a_1] \approx \frac{1}{2}$
\vdots	\vdots	\vdots	\vdots
ℓ .	$\tilde{O}(1)$ samples from $Y Y \geq a_{\ell-1}$ Requires $\tilde{O}(\sqrt{2^{\ell-1}})$ samples from Y by <i>amplitude amplification</i>	\hookrightarrow	Let a_ℓ be the median $\mathbb{P}[Y \geq a_\ell] \approx \frac{1}{2^\ell}$
\vdots	\vdots	\vdots	\vdots
$2 \log(n)$.	$\tilde{O}(1)$ samples from $Y Y \geq a_{2 \log(n)}$ Requires $\tilde{O}(n)$ samples from Y by <i>a.a.</i>	\hookrightarrow	Let $a_{2 \log(n)}$ be the median $\mathbb{P}[Y \geq a_{2 \log(n)}] \approx \frac{1}{n^2}$




Mean estimation on the rings

Mean estimation on the rings

We consider $X^{(\ell)} = X \cdot \mathbb{1}_{X \in R_\ell} - \bar{\mu}$.

Goal: Estimate $\mathbb{E}[X^{(\ell)}]$.




Mean estimation on the rings

We consider $X^{(\ell)} = X \cdot \mathbb{1}_{X \in R_\ell} - \bar{\mu}$.

Goal: Estimate $\mathbb{E}[X^{(\ell)}]$.

- 1 *Amplitude amplification on the ring:*
Requires $\tilde{O}(\sqrt{2^\ell})$ samples.




Mean estimation on the rings

We consider $X^{(\ell)} = X \cdot \mathbb{1}_{X \in R_\ell} - \bar{\mu}$.

Goal: Estimate $\mathbb{E}[X^{(\ell)}]$.

- ① *Amplitude amplification on the ring:*
Requires $\tilde{O}(\sqrt{2^\ell})$ samples.




Mean estimation on the rings

We consider $X^{(\ell)} = X \cdot \mathbb{1}_{X \in R_\ell} - \bar{\mu}$.

Goal: Estimate $\mathbb{E}[X^{(\ell)}]$.

- ① *Amplitude amplification on the ring:*
Requires $\tilde{O}(\sqrt{2^\ell})$ samples.



Mean estimation on the rings

We consider $X^{(\ell)} = X \cdot \mathbb{1}_{X \in R_\ell} - \bar{\mu}$.

Goal: Estimate $\mathbb{E}[X^{(\ell)}]$.


- 1 *Amplitude amplification on the ring:*

Requires $\tilde{O}(\sqrt{2^\ell})$ samples.

- 2 *Phase encoding techniques* [GSLW18]:

$$|\mathbf{u}\rangle \mapsto e^{i2^\ell \cdot \mathbf{u}^T \mathbb{E}[X^{(\ell)}]} |\mathbf{u}\rangle$$

Requires $\tilde{O}(1)$ calls to 1.



Mean estimation on the rings

We consider $X^{(\ell)} = X \cdot \mathbb{1}_{X \in R_\ell} - \bar{\mu}$.

Goal: Estimate $\mathbb{E}[X^{(\ell)}]$.


- 1 *Amplitude amplification on the ring:*

Requires $\tilde{O}(\sqrt{2^\ell})$ samples.

- 2 *Phase encoding techniques* [GSLW18]:

$$|\mathbf{u}\rangle \mapsto e^{i2^\ell \cdot \mathbf{u}^T \mathbb{E}[X^{(\ell)}]} |\mathbf{u}\rangle$$

Requires $\tilde{O}(1)$ calls to 1.



Mean estimation on the rings

We consider $X^{(\ell)} = X \cdot \mathbb{1}_{X \in R_\ell} - \bar{\mu}$.

Goal: Estimate $\mathbb{E}[X^{(\ell)}]$.

- 1 *Amplitude amplification on the ring:*

Requires $\tilde{O}(\sqrt{2^\ell})$ samples.

- 2 *Phase encoding techniques* [GSLW18]:


$$|\mathbf{u}\rangle \mapsto e^{i2^\ell \cdot \mathbf{u}^T \mathbb{E}[X^{(\ell)}]} |\mathbf{u}\rangle$$

Requires $\tilde{O}(1)$ calls to 1.

- 3 *Bernstein-Vazirani over the reals* [GAW18]:

$$\|\tilde{\mu}^{(\ell)} - \mathbb{E}[X^{(\ell)}]\|_\infty = \tilde{O}(a_\ell / (n\sqrt{2^\ell})).$$

Requires $\tilde{O}(n/\sqrt{2^\ell})$ calls to 2.




$$\mathbb{P}[X \in R_\ell] \approx \frac{1}{2^\ell}$$

For most \mathbf{u} :

$$|\mathbf{u}^T X^{(\ell)}(\omega)| \leq 1$$

(Naively: $\leq \sqrt{d}$)



Mean estimation on the rings

We consider $X^{(\ell)} = X \cdot \mathbb{1}_{X \in R_\ell} - \bar{\mu}$.

Goal: Estimate $\mathbb{E}[X^{(\ell)}]$.

- 1 *Amplitude amplification on the ring:*

Requires $\tilde{O}(\sqrt{2^\ell})$ samples.

- 2 *Phase encoding techniques* [GSLW18]:


$$|\mathbf{u}\rangle \mapsto e^{i2^\ell \cdot \mathbf{u}^T \mathbb{E}[X^{(\ell)}]} |\mathbf{u}\rangle$$

Requires $\tilde{O}(1)$ calls to 1.

- 3 *Bernstein-Vazirani over the reals* [GAW18]:

$$\begin{aligned} \|\tilde{\mu}^{(\ell)} - \mathbb{E}[X^{(\ell)}]\|_\infty &= \tilde{O}(a_\ell / (n\sqrt{2^\ell})). \\ &= \tilde{O}(\sqrt{\text{Tr}[\Sigma]}/n). \end{aligned}$$

Requires $\tilde{O}(n/\sqrt{2^\ell})$ calls to 2.




$$\mathbb{P}[X \in R_\ell] \approx \frac{1}{2^\ell}$$

For most \mathbf{u} :

$$|\mathbf{u}^T X^{(\ell)}(\omega)| \leq 1$$

(Naively: $\leq \sqrt{d}$)



Mean estimation on the rings

We consider $X^{(\ell)} = X \cdot \mathbb{1}_{X \in R_\ell} - \bar{\mu}$.

Goal: Estimate $\mathbb{E}[X^{(\ell)}]$.

- 1 *Amplitude amplification on the ring:*

Requires $\tilde{O}(\sqrt{2^\ell})$ samples.

- 2 *Phase encoding techniques* [GSLW18]:

$$|\mathbf{u}\rangle \mapsto e^{i2^\ell \cdot \mathbf{u}^T \mathbb{E}[X^{(\ell)}]} |\mathbf{u}\rangle$$

Requires $\tilde{O}(1)$ calls to 1.


- 3 *Bernstein-Vazirani over the reals* [GAW18]:

$$\begin{aligned} \|\tilde{\mu}^{(\ell)} - \mathbb{E}[X^{(\ell)}]\|_\infty &= \tilde{O}(a_\ell / (n\sqrt{2^\ell})). \\ &= \tilde{O}(\sqrt{\text{Tr}[\Sigma]}/n). \end{aligned}$$

Requires $\tilde{O}(n/\sqrt{2^\ell})$ calls to 2.

- 4 *Conversion to ℓ_2 -norm* (Hölder's inequality):

$$\|\tilde{\mu}^{(\ell)} - \mathbb{E}[X^{(\ell)}]\|_2 = \tilde{O}(\sqrt{d \text{Tr}[\Sigma]}/n).$$




$$\mathbb{P}[X \in R_\ell] \approx \frac{1}{2^\ell}$$

For most \mathbf{u} :

$$|\mathbf{u}^T X^{(\ell)}(\omega)| \leq 1$$

(Naively: $\leq \sqrt{d}$)



Concluding remarks

Concluding remarks


Main result:

Optimal estimator $\tilde{\mu}$ with n samples, s.t.

$$\mathbb{P} [\|\mu - \tilde{\mu}\|_2 \geq \varepsilon(n)] \leq \frac{1}{3},$$

has precision

$$\varepsilon(n) = \tilde{\Theta} \left(\begin{cases} \frac{\sqrt{d \operatorname{Tr}[\Sigma]}}{n}, & \text{if } n \geq d, \\ \sqrt{\frac{\operatorname{Tr}[\Sigma]}{n}}, & \text{if } n < d \end{cases} \right).$$



Concluding remarks

Open questions:

Main result:

Optimal estimator $\tilde{\mu}$ with n samples, s.t.

$$\mathbb{P} [\|\mu - \tilde{\mu}\|_2 \geq \varepsilon(n)] \leq \frac{1}{3},$$

has precision

$$\varepsilon(n) = \tilde{\Theta} \left(\begin{cases} \frac{\sqrt{d \operatorname{Tr}[\Sigma]}}{n}, & \text{if } n \geq d, \\ \sqrt{\frac{\operatorname{Tr}[\Sigma]}{n}}, & \text{if } n < d \end{cases} \right).$$

Concluding remarks

Main result:

Optimal estimator $\tilde{\mu}$ with n samples, s.t.

$$\mathbb{P} [\|\mu - \tilde{\mu}\|_2 \geq \varepsilon(n)] \leq \frac{1}{3},$$

has precision

$$\varepsilon(n) = \tilde{\Theta} \left(\begin{cases} \frac{\sqrt{d \operatorname{Tr}[\Sigma]}}{n}, & \text{if } n \geq d, \\ \sqrt{\frac{\operatorname{Tr}[\Sigma]}{n}}, & \text{if } n < d \end{cases} \right).$$

Open questions:

- Dependence on the failure probability δ ?
 - Classically: [LM19; Hop20]
$$\varepsilon(n) = O \left(\sqrt{\frac{\operatorname{Tr}[\Sigma]}{n}} + \sqrt{\frac{\|\Sigma\| \log \frac{1}{\delta}}{n}} \right)$$
 - Constant prefactors: [LV20; LV22].

Concluding remarks

Main result:

Optimal estimator $\tilde{\mu}$ with n samples, s.t.

$$\mathbb{P} [\|\mu - \tilde{\mu}\|_2 \geq \varepsilon(n)] \leq \frac{1}{3},$$

has precision

$$\varepsilon(n) = \tilde{\Theta} \left(\begin{cases} \frac{\sqrt{d \operatorname{Tr}[\Sigma]}}{n}, & \text{if } n \geq d, \\ \sqrt{\frac{\operatorname{Tr}[\Sigma]}{n}}, & \text{if } n < d \end{cases} \right).$$

Open questions:

- Dependence on the failure probability δ ?
 - Classically: [LM19; Hop20]
$$\varepsilon(n) = O \left(\sqrt{\frac{\operatorname{Tr}[\Sigma]}{n}} + \sqrt{\frac{\|\Sigma\| \log \frac{1}{\delta}}{n}} \right)$$
 - Constant prefactors: [LV20; LV22].
- Optimality in different norms?

Concluding remarks

Main result:

Optimal estimator $\tilde{\mu}$ with n samples, s.t.

$$\mathbb{P} [\|\mu - \tilde{\mu}\|_2 \geq \varepsilon(n)] \leq \frac{1}{3},$$

has precision

$$\varepsilon(n) = \tilde{\Theta} \left(\begin{cases} \frac{\sqrt{d \operatorname{Tr}[\Sigma]}}{n}, & \text{if } n \geq d, \\ \sqrt{\frac{\operatorname{Tr}[\Sigma]}{n}}, & \text{if } n < d \end{cases} \right).$$

Open questions:

- Dependence on the failure probability δ ?

- Classically: [LM19; Hop20]

$$\varepsilon(n) = O \left(\sqrt{\frac{\operatorname{Tr}[\Sigma]}{n}} + \sqrt{\frac{\|\Sigma\| \log \frac{1}{\delta}}{n}} \right)$$

- Constant prefactors: [LV20; LV22].

- Optimality in different norms?

- Different access models?

$$|\omega\rangle |0\rangle \mapsto |\omega\rangle |X(\omega)\rangle.$$

$$|\omega\rangle |j\rangle \mapsto e^{iX(\omega)_j} |\omega\rangle |j\rangle.$$

Concluding remarks

Main result:

Optimal estimator $\tilde{\mu}$ with n samples, s.t.

$$\mathbb{P} [\|\mu - \tilde{\mu}\|_2 \geq \varepsilon(n)] \leq \frac{1}{3},$$

has precision

$$\varepsilon(n) = \tilde{\Theta} \left(\begin{cases} \frac{\sqrt{d \operatorname{Tr}[\Sigma]}}{n}, & \text{if } n \geq d, \\ \sqrt{\frac{\operatorname{Tr}[\Sigma]}{n}}, & \text{if } n < d \end{cases} \right).$$

Open questions:

- Dependence on the failure probability δ ?
 - Classically: [LM19; Hop20]
$$\varepsilon(n) = O \left(\sqrt{\frac{\operatorname{Tr}[\Sigma]}{n}} + \sqrt{\frac{\|\Sigma\| \log \frac{1}{\delta}}{n}} \right)$$
 - Constant prefactors: [LV20; LV22].
- Optimality in different norms?
- Different access models?
$$\begin{aligned} |\omega\rangle |0\rangle &\mapsto |\omega\rangle |X(\omega)\rangle. \\ |\omega\rangle |j\rangle &\mapsto e^{iX(\omega)_j} |\omega\rangle |j\rangle. \end{aligned}$$
- Can prior knowledge on Σ help?

Concluding remarks

Main result:

Optimal estimator $\tilde{\mu}$ with n samples, s.t.

$$\mathbb{P} [\|\mu - \tilde{\mu}\|_2 \geq \varepsilon(n)] \leq \frac{1}{3},$$

has precision

$$\varepsilon(n) = \tilde{\Theta} \left(\begin{cases} \frac{\sqrt{d \operatorname{Tr}[\Sigma]}}{n}, & \text{if } n \geq d, \\ \sqrt{\frac{\operatorname{Tr}[\Sigma]}{n}}, & \text{if } n < d \end{cases} \right).$$

Open questions:

- Dependence on the failure probability δ ?
 - Classically: [LM19; Hop20]
$$\varepsilon(n) = O \left(\sqrt{\frac{\operatorname{Tr}[\Sigma]}{n}} + \sqrt{\frac{\|\Sigma\| \log \frac{1}{\delta}}{n}} \right)$$
 - Constant prefactors: [LV20; LV22].
- Optimality in different norms?
- Different access models?
$$\begin{aligned} |\omega\rangle |0\rangle &\mapsto |\omega\rangle |X(\omega)\rangle. \\ |\omega\rangle |j\rangle &\mapsto e^{iX(\omega)_j} |\omega\rangle |j\rangle. \end{aligned}$$
- Can prior knowledge on Σ help?

Thanks for your attention!
hamoudi@berkeley.edu

References

- [GAW18] A. Gilyén, S. Arunachalam, N. Wiebe. *Optimizing quantum optimization algorithms via faster quantum gradient computation*. arXiv:1711.00465.
- [GLSW18] A., Y. Su, G. H. Low, N. Wiebe. *Quantum singular value transformation and beyond: exponential improvements for quantum matrix arithmetics*. arXiv:1806.01838.
- [Ham22] Y. Hamoudi. *Quantum Sub-Gaussian Mean Estimator*. arXiv:2108.12172.
- [Hei02] S. Heinrich. *Quantum Summation with an Application to Integration*. arXiv:quant-ph/0105116.
- [HM19] Y. Hamoudi, F. Magniez. *Quantum Chebyshev's Inequality and Applications*. arXiv:1807.06456.
- [Hop20] S. B. Hopkins. *Mean Estimation with Sub-Gaussian Rates in Polynomial Time*. arXiv:1809.07425.
- [LM19] G. Lugosi, S. Mendelson. *Mean estimation and regression under heavy-tailed distributions – a survey*. arXiv:1906.04280.
- [LV20] J. C.H. Lee, P. Valiant. *Optimal Sub-Gaussian Mean Estimation in \mathbb{R}* . arXiv:2011.08384.
- [LV22] J. C.H. Lee, P. Valiant. *Optimal Sub-Gaussian Mean Estimation in Very High Dimensions*. arXiv:2011.08384.
- [Mon15] A. Montanaro. *Quantum speedup of Monte Carlo methods*. arXiv:1504.06987