


# Quantum Sub-Gaussian Mean Estimator


Yassine Hamoudi

ESA 2021


Experiment with unknown outcome distribution  $D$



Experiment with unknown outcome distribution  $D$




Experiment with unknown outcome distribution  $D$



$n = \#$  of runs of the experiment  $\left\{ \begin{array}{l} \text{classical} = n \text{ i.i.d. samples from } D \\ \text{quantum} = \text{see later} \end{array} \right.$

Experiment with unknown outcome distribution  $D$



$n = \#$  of runs of the experiment  $\begin{cases} \text{classical} = n \text{ i.i.d. samples from } D \\ \text{quantum} = \text{see later} \end{cases}$

**Goal:** compute an estimate  $\tilde{\mu}$  that minimizes the error  $\epsilon(n)$  such that

$$\Pr \left[ |\mu - \tilde{\mu}| > \epsilon(n) \right] < \delta \quad \text{given } \delta \in (0,1)$$

$$\Pr \left[ |\mu - \tilde{\mu}| > \varepsilon(n) \right] < \delta$$

$$\Pr \left[ |\mu - \tilde{\mu}| > \varepsilon(n) \right] < \delta$$

→ If  $D \in \left\{ \mathcal{N}(\mu, \sigma^2) \right\}_{\mu, \sigma}$  is Gaussian then the Empirical Mean is optimal and

$$\varepsilon(n) = O\left(\sqrt{\frac{\sigma^2 \log(1/\delta)}{n}}\right)$$

$$\Pr \left[ |\mu - \tilde{\mu}| > \varepsilon(n) \right] < \delta$$

→ If  $D \in \left\{ \mathcal{N}(\mu, \sigma^2) \right\}_{\mu, \sigma}$  is Gaussian then the Empirical Mean is optimal and

$$\varepsilon(n) = O\left(\sqrt{\frac{\sigma^2 \log(1/\delta)}{n}}\right)$$

→ An estimator is **sub-Gaussian** if it achieves the above bound for **any**  $D$



$$\Pr \left[ |\mu - \tilde{\mu}| > \varepsilon(n) \right] < \delta$$

→ If  $D \in \left\{ \mathcal{N}(\mu, \sigma^2) \right\}_{\mu, \sigma}$  is Gaussian then the Empirical Mean is optimal and

$$\varepsilon(n) = O\left(\sqrt{\frac{\sigma^2 \log(1/\delta)}{n}}\right)$$

→ An estimator is **sub-Gaussian** if it achieves the above bound for **any**  $D$

**✗** Empirical Mean is not sub-Gaussian

$$\Pr \left[ |\mu - \tilde{\mu}| > \varepsilon(n) \right] < \delta$$

→ If  $D \in \left\{ \mathcal{N}(\mu, \sigma^2) \right\}_{\mu, \sigma}$  is Gaussian then the Empirical Mean is optimal and

$$\varepsilon(n) = O\left(\sqrt{\frac{\sigma^2 \log(1/\delta)}{n}}\right)$$

→ An estimator is **sub-Gaussian** if it achieves the above bound for **any**  $D$

✗ Empirical Mean is not sub-Gaussian

✓ Median-of-Means, [Catoni'12], [Lee, Valiant'21], ... are sub-Gaussian

$$\Pr \left[ |\mu - \tilde{\mu}| > \varepsilon(n) \right] < \delta$$

→ If  $D \in \left\{ \mathcal{N}(\mu, \sigma^2) \right\}_{\mu, \sigma}$  is Gaussian then the Empirical Mean is optimal and

$$\varepsilon(n) = O\left(\sqrt{\frac{\sigma^2 \log(1/\delta)}{n}}\right)$$

→ An estimator is **sub-Gaussian** if it achieves the above bound for **any**  $D$

✗ Empirical Mean is not sub-Gaussian

✓ Median-of-Means, [Catoni'12], [Lee, Valiant'21], ... are sub-Gaussian

↑  
optimal  $\sqrt{2} + o(1)$  factor in  $O(\cdot)$

**Classical:**  $\varepsilon(n) = O\left(\sqrt{\frac{\sigma^2 \log(1/\delta)}{n}}\right)$

**Quantum:**  $\varepsilon(n) \approx \dots$

**Classical:**  $\varepsilon(n) = O\left(\sqrt{\frac{\sigma^2 \log(1/\delta)}{n}}\right)$

**Quantum:**  $\varepsilon(n) \approx \dots$

$$\dots \frac{\textcolor{red}{B} \log(1/\delta)}{n}$$

given  $B \geq \max\{|x| : p_x \neq 0\}$

[Grover'98] [Abrams,Williams'99]

[Brassard,Dupuis,Gambs,Tapp'11] ...

**Classical:**  $\varepsilon(n) = O\left(\sqrt{\frac{\sigma^2 \log(1/\delta)}{n}}\right)$

**Quantum:**  $\varepsilon(n) \approx \dots$

$$\dots \frac{B \log(1/\delta)}{n}$$

given  $B \geq \max\{|x| : p_x \neq 0\}$

[Grover'98] [Abrams, Williams'99]

[Brassard, Dupuis, Gambs, Tapp'11] ...

$$\dots \frac{\sqrt{B\mu} \log(1/\delta)}{n} + \frac{B \log(1/\delta)^2}{n^2}$$

[Brassard, Høyer, Mosca, Tapp'02]

**Amplitude Estimation**

**Classical:**  $\varepsilon(n) = O\left(\sqrt{\frac{\sigma^2 \log(1/\delta)}{n}}\right)$

**Quantum:**  $\varepsilon(n) \approx \dots$

$$\dots \frac{B \log(1/\delta)}{n}$$

given  $B \geq \max\{|x| : p_x \neq 0\}$

[Grover'98] [Abrams, Williams'99]  
[Brassard, Dupuis, Gambs, Tapp'11] ...

$$\dots \frac{\sqrt{B\mu} \log(1/\delta)}{n} + \frac{B \log(1/\delta)^2}{n^2}$$

[Brassard, Høyer, Mosca, Tapp'02]

**Amplitude Estimation**

$$\dots \frac{\Sigma \log(1/\delta)}{n}$$

given  $\Sigma \geq \sigma$

[Heinrich'02][Montanaro'15]

**Classical:**  $\varepsilon(n) = O\left(\sqrt{\frac{\sigma^2 \log(1/\delta)}{n}}\right)$

**Quantum:**  $\varepsilon(n) \approx \dots$

$$\dots \frac{B \log(1/\delta)}{n}$$

given  $B \geq \max\{|x| : p_x \neq 0\}$

[Grover'98] [Abrams, Williams'99]  
[Brassard, Dupuis, Gambs, Tapp'11] ...

$$\dots \frac{\sqrt{B\mu} \log(1/\delta)}{n} + \frac{B \log(1/\delta)^2}{n^2}$$

[Brassard, Høyer, Mosca, Tapp'02]

**Amplitude Estimation**

$$\dots \frac{\Sigma \log(1/\delta)}{n}$$

given  $\Sigma \geq \sigma$

[Heinrich'02][Montanaro'15]

$$\dots \frac{\mu \Delta \log(1/\delta)}{n}$$

given  $\Delta \geq \sigma/\mu$

[H., Magniez'19]



Quantum “sub-Gaussian” estimator:

$$\varepsilon(n) = \tilde{O}\left(\frac{\sigma \log(1/\delta)}{n}\right)$$

Quantum “sub-Gaussian” estimator:

$$\varepsilon(n) = \tilde{O}\left(\frac{\sigma \log(1/\delta)}{n}\right)$$

- ✓ Doesn't require any prior knowledge about  $D$
- ✓ Subsumes all previous quantum estimators
- ✓ Better than any classical sub-Gaussian estimator (recall:  $\sqrt{\frac{\sigma^2 \log(1/\delta)}{n}}$ )
- ✓ Optimal (lower bound by reduction from Quantum Search)

*\*up to log factors*

Unknown distribution  $D = (p_x)_x$

Unknown distribution  $D = (p_x)_x$

Model 1:  $n$  copies of the qsample  $\sum_x \sqrt{p_x} |x\rangle$

Unknown distribution  $D = (p_x)_x$

Model 1:  $n$  copies of the qsample  $\sum_x \sqrt{p_x} |x\rangle$

→ **classical** sub-Gaussian estimators are optimal in this model!

Unknown distribution  $D = (p_x)_x$

Model 1:  $n$  copies of the qsample  $\sum_x \sqrt{p_x} |x\rangle$

→ **classical** sub-Gaussian estimators are optimal in this model!

Model 2:  $n$  accesses to a **unitary**  $U_D$  such that  $U_D |0\rangle = \sum_x \sqrt{p_x} |x\rangle |\text{garbage}_x\rangle$

Unknown distribution  $D = (p_x)_x$

Model 1:  $n$  copies of the qsample  $\sum_x \sqrt{p_x} |x\rangle$

→ **classical** sub-Gaussian estimators are optimal in this model!

Model 2:  $n$  accesses to a **unitary**  $U_D$  such that  $U_D |0\rangle = \sum_x \sqrt{p_x} |x\rangle |\text{garbage}_x\rangle$

→  $U_D, U_D^{-1}$  and their controlled versions can be accessed

Unknown distribution  $D = (p_x)_x$

Model 1:  $n$  copies of the qsample  $\sum_x \sqrt{p_x} |x\rangle$

→ **classical** sub-Gaussian estimators are optimal in this model!

Model 2:  $n$  accesses to a **unitary**  $U_D$  such that  $U_D |0\rangle = \sum_x \sqrt{p_x} |x\rangle |\text{garbage}_x\rangle$

→  $U_D, U_D^{-1}$  and their controlled versions can be accessed

→ Grover Search, Phase/Amplitude Estimation... can be used on  $U_D$



Unknown distribution  $D = (p_x)_x$

Model 1:  $n$  copies of the qsample  $\sum_x \sqrt{p_x} |x\rangle$

→ **classical** sub-Gaussian estimators are optimal in this model!


Model 2:  $n$  accesses to a **unitary**  $U_D$  such that  $U_D |0\rangle = \sum_x \sqrt{p_x} |x\rangle |\text{garbage}_x\rangle$

→  $U_D, U_D^{-1}$  and their controlled versions can be accessed


→ Grover Search, Phase/Amplitude Estimation... can be used on  $U_D$

→ Similar models in other works on estimating/testing statistics  
[Nayak, Wu'99] [Bravyi, Harrow, Hassidim'11] [Montanaro'15]...

Empirical Mean and Amplitude Estimation are too sensitive to **outliers**




Empirical Mean and Amplitude Estimation are too sensitive to **outliers**




Natural approach: **truncate** the distribution at a certain level  $T$

Empirical Mean and Amplitude Estimation are too sensitive to **outliers**



Natural approach: **truncate** the distribution at a certain level  $T$


Empirical Mean and Amplitude Estimation are too sensitive to **outliers**



Natural approach: **truncate** the distribution at a certain level  $T$

→ How to choose  $T$  ?

Empirical Mean and Amplitude Estimation are too sensitive to **outliers**



Natural approach: **truncate** the distribution at a certain level  $T$


→ How to choose  $T$  ?

$$T \approx n\sigma$$

[Heinrich'02]  
[Montanaro'15]  
[H.,Magniez'19]

Require some  
knowledge about  $\sigma$

Empirical Mean and Amplitude Estimation are too sensitive to **outliers**



Natural approach: **truncate** the distribution at a certain level  $T$

→ How to choose  $T$ ?


$$T \approx n\sigma$$

[Heinrich'02]  
[Montanaro'15]  
[H.,Magniez'19]

Require some  
knowledge about  $\sigma$

Our work:  $T \approx$  quantile satisfying  $\Pr_{x \leftarrow D}[x > T] = 1/n^2$

Empirical Mean and Amplitude Estimation are too sensitive to **outliers**



Natural approach: **truncate** the distribution at a certain level  $T$

→ How to choose  $T$ ?

$$T \approx n\sigma$$

[Heinrich'02]  
[Montanaro'15]  
[H.,Magniez'19]

Require some  
knowledge about  $\sigma$


Our work:

$$T \approx \text{quantile satisfying } \Pr_{x \leftarrow D}[x > T] = 1/n^2$$


Computed in  $O(n)$  time  
by adapting Minimum  
Finding [Dürr,Høyer'96]



**Assumptions:**  $\delta = 1/3$  + non-negative outcomes +  $\sigma^2$  replaced with  $\mathbb{E}_D(x^2)$




**Assumptions:**  $\delta = 1/3$  + non-negative outcomes +  $\sigma^2$  replaced with  $\mathbb{E}_D(x^2)$




$T$  such that  $\Pr_D[x > T] = 1/n^2$


**Assumptions:**  $\delta = 1/3$  + non-negative outcomes +  $\sigma^2$  replaced with  $\mathbb{E}_D(x^2)$



**Assumptions:**  $\delta = 1/3$  + non-negative outcomes +  $\sigma^2$  replaced with  $\mathbb{E}_D(x^2)$




**Assumptions:**  $\delta = 1/3$  + non-negative outcomes +  $\sigma^2$  replaced with  $\mathbb{E}_D(x^2)$



$$\tilde{\mu} = \tilde{\mu}_1 + \tilde{\mu}_2 + \tilde{\mu}_3 + \tilde{\mu}_4 + \dots + 0$$


**Assumptions:**  $\delta = 1/3$  + non-negative outcomes +  $\sigma^2$  replaced with  $\mathbb{E}_D(x^2)$



$$\tilde{\mu} = \tilde{\mu}_1 + \tilde{\mu}_2 + \tilde{\mu}_3 + \tilde{\mu}_4 + \dots + 0$$

$$\varepsilon(n) \approx \frac{\sqrt{\mathbb{E}_D(x^2)}}{n} + \frac{T}{n^2} + \sqrt{\mathbb{E}_D(x^2) \cdot \Pr_D[x > T]}$$

**Assumptions:**  $\delta = 1/3$  + non-negative outcomes +  $\sigma^2$  replaced with  $\mathbb{E}_D(x^2)$




$$\tilde{\mu} = \tilde{\mu}_1 + \tilde{\mu}_2 + \tilde{\mu}_3 + \tilde{\mu}_4 + \dots + 0$$

$$\varepsilon(n) \approx \frac{\sqrt{\mathbb{E}_D(x^2)}}{n} + \frac{T}{n^2} + \sqrt{\mathbb{E}_D(x^2) \cdot \Pr_D[x > T]}$$

$\leq n\sqrt{\mathbb{E}_D(x^2)}$

**Assumptions:**  $\delta = 1/3$  + non-negative outcomes +  $\sigma^2$  replaced with  $\mathbb{E}_D(x^2)$



$$\tilde{\mu} = \tilde{\mu}_1 + \tilde{\mu}_2 + \tilde{\mu}_3 + \tilde{\mu}_4 + \dots + 0$$

$$\varepsilon(n) \approx \frac{\sqrt{\mathbb{E}_D(x^2)}}{n} + \frac{T}{n^2} + \sqrt{\mathbb{E}_D(x^2) \cdot \Pr_D[x > T]}$$

$\leq n\sqrt{\mathbb{E}_D(x^2)}$



- Estimating expected values of **observables**

- observable  $O$ , state  $|\psi\rangle$

- $\mu = \langle\psi|O|\psi\rangle$  and  $\sigma^2 = \langle\psi|(O - \mu)^2|\psi\rangle$

- Estimating expected values of **observables**

- observable  $O$ , state  $|\psi\rangle$

- $\mu = \langle\psi|O|\psi\rangle$  and  $\sigma^2 = \langle\psi|(O - \mu)^2|\psi\rangle$

- if  $|\psi\rangle = e^{-iHt}|0\rangle$  is obtained by Hamiltonian sim. then  $n = \#$  runs of Hs

- Estimating expected values of **observables**

- observable  $O$ , state  $|\psi\rangle$

- $\mu = \langle\psi|O|\psi\rangle$  and  $\sigma^2 = \langle\psi|(O - \mu)^2|\psi\rangle$

- if  $|\psi\rangle = e^{-iHt}|0\rangle$  is obtained by Hamiltonian sim. then  $n = \#$  runs of Hs

- **Monte Carlo** estimation

- integration

- partition functions

- mathematical finance

- Estimating expected values of **observables**

- observable  $O$ , state  $|\psi\rangle$

- $\mu = \langle \psi | O | \psi \rangle$  and  $\sigma^2 = \langle \psi | (O - \mu)^2 | \psi \rangle$

- if  $|\psi\rangle = e^{-iHt} |0\rangle$  is obtained by Hamiltonian sim. then  $n = \#$  runs of Hs

- **Monte Carlo** estimation

- integration

- partition functions

- mathematical finance

- ...

- Sub-Gaussian estimators

$$\Theta\left(\sqrt{\frac{\sigma^2 \log(1/\delta)}{n}}\right)$$

Classical

vs

$$\tilde{\Theta}\left(\frac{\sigma \log(1/\delta)}{n}\right)$$

Quantum

- Open questions

- Improve the log-factors
- Extend to the **multivariate** setting (recent work: [Cornelissen, Jerbi'21])
- Find other applications