Term Test 1 solutions

Calculus (University of Toronto)



messages.pdf_cover_qr_code_label

messages.downloaded_by
messages.studocu_not_sponsored_or_endorsed_by_college

University of Toronto
Faculty of Arts and Science

MAT137Y Test 1

X. Cui, I. Gaiur, T. Kojar, L. Salvetti Martinez, T. Xian, J. Zhang, X. Zhang

October 12, 2023

Duration: 110 minutes

No Aids Permitted

Sample Solutions

studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by



1. For each of the following questions, only your final answer will be graded.

No justification is necessary.

Evaluate the following limits. If a limit is ∞ or $-\infty$, clearly state this. If the limit does not exist and it's not ∞ or $-\infty$, put "DNE" in the box.

(1a) (2 points) Given the function $f(x) = \begin{cases} 7 - 4x & x < 1 \\ x^2 + 3 & x \geq 1 \end{cases}$.

- Find $\lim_{x \to -2} f(x)$.

Final Answer

- Find $\lim_{x \to 1} f(x)$.

Final Answer

(1b) (1 point) $\lim_{x \to \pi^-} \ln\left(\frac{1}{\sin x}\right)$

Final Answer

(1c) (1 point) $\lim_{x \to 4} \frac{\sqrt{x} - 2}{x - 4}$

Final Answer

(1d) (1 point) $\lim_{x \to +\infty} \frac{\cos(2x) + 2}{x^2 + x + 1}$

Final Answer

studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by



For each of the following questions, only your final answer will be graded.

No justification is necessary.

2. (2 points) Let $a, L \in \mathbb{R}$. Let f be a function defined on $\mathbb{R}$. Write the complete $\varepsilon - \delta$ formal definition for "$\lim_{x \to a^+} f(x) \neq L$".

Final Answer

[redacted]

3. (2 points) Let f be a function defined on $\mathbb{R}$. Write the complete $\varepsilon - \delta$ formal definition for "$f(x)$ is continuous at a".

Final Answer

[redacted]

4. (3 points) Classify the following statements as **TRUE**, **FALSE**.

   (a) $\forall x \in \mathbb{Z}, \exists y \in \mathbb{Q}$ such that $xy = 1$.

   ◯ True [redacted]

   (b) $5 \in \left\{x \in \mathbb{R} : \exists y \in \mathbb{Z} \text{ } s.t. \text{ } y^2 > 1 \text{ } and \text{ } y < x\right\}$.

   [redacted] ◯ False

   (c) $\forall x \in \mathbb{Q}, x > 0 \implies x \geq 0$

   [redacted] ◯ False

studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by



For each of the following questions, only your final answer will be graded.

No justification is necessary.

5. (5a) (1 point) Write down the set of rational numbers using set builder notation.

Final Answer

[redacted]

(5b) (1 point) Express the following set $\{x \in [0,1] : \exists y \in [-1,1],\ s.t.\ x > y^2\}$ in interval notation.

Final Answer

[redacted]

(5c) (1 point) Find a function f with domain $\mathbb{R}$ such that

- f is continuous on $\mathbb{R}$, except at 0.
- $\lim_{x \to 0^+} f(x) \neq \lim_{x \to 0^-} f(x)$ and $f(0) = 1$.

It is okay to use piece-wise-defined functions.

Final Answer

[redacted]

(5d) (1 point) Find a pair of two functions f and g defined on $\mathbb{R}$ such that $\lim_{x \to 1} f(x) = 0$ and $\lim_{x \to 1} f(x)g(x) \neq 0$. It is okay to use piece-wise-defined functions.

Final Answer

[redacted]


studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by



6. (6 points) Sketch the graph of a function f defined on $\mathbb{R}$ which satisfies the following properties at once.

- $\lim_{x\to 0} f(x) = -2$
- $\lim_{x\to 0} f(f(x)) = 1$
- $\lim_{x\to 0^-} f(\frac{1}{x}) = 3$
- $\lim_{x\to 2^+} f(x) = -\infty$
- $\lim_{x\to 2}[f(x)]^2 = \infty$
- $\lim_{x\to 3} f(x) = -1, f(3) = 2$



studocu
messages.pdf_cover_qr_code_label
messages.downloaded_by

7. Let $f(x)$ be a function defined on $\mathbb{R}$ such that $\lim_{x\to\infty} f(x) = 2$.

(7a) (2 points) Write the formal definition of $\lim_{x\to\infty} f(x) = 2$.

[redacted]

(7b) (5 points) Use the formal ε-δ definition of the limit to prove

$$\lim_{x\to\infty} 4f(3x+1) = 8.$$

Do not use any limit laws or any other theorems.

[redacted]

studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by



8. (5 points) Define $f(n) = 1^2 + 2^2 + 3^2 + \cdots + n^2$ where n is any positive integer. Prove the following statement by using math induction.

$$P(n) : f(n) \leq n^3$$

is true for all positive integer n.



studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by

9. Let A, B be any subsets of $\mathbb{R}$ and non-empty.
We say A is *calm* if

$$\exists a, b \in A, \ s.t. \ \forall x \in A, a \leq x \leq b$$

We say B is *happy* if

$$\exists a \in B, \ s.t. \ \forall x \in B, x \geq a$$

Determine if the following statements are true or false. If a claim is true, prove it. If a claim is false, provide a counterexample and a justification of how the counterexample shows the claim is false.

(9a) (3 points) If a non-empty set B is happy, then it is calm.

◯ True [redacted]

[redacted]

(9b) (3 points) If a non-empty set A is calm and a non-empty set B is happy, $A \neq B$, then $A \cup B$ is happy.

[redacted] ◯ False

[redacted]

studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by

