Term Test 1 Solutions 2021

Calculus (University of Toronto)



messages.pdf_cover_qr_code_label

messages.downloaded_by
messages.studocu_not_sponsored_or_endorsed_by_college

University of Toronto
Faculty of Arts and Science

MAT137Y Test 1

X. Cui, V. Dimitrov, A. Malusà, B. Khesin, Z.Qian

October 21, 2021

Duration: 110 minutes

No Aids Permitted

studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by



1. For each of the following questions, only your final answer will be graded.

No justification is necessary.

(1a) (2 points) Let $L \in \mathbb{R}$ and f be a function defined on an open interval of the form (p, ∞) for some $p \in \mathbb{R}$. Give the complete formal definition for $\lim_{x \to \infty} f(x) = L$.



(1b) (2 points) Let $a \in \mathbb{R}$ and f be a function defined on an open interval centered at a, except possibly at a. Give the complete formal definition for $\lim_{x \to a} f(x)$ does not exist.



For (1c) to (1f) below, evaluate the limit. If the limit does not exist and it's not $\pm\infty$, put "DNE" in the box.

(1c) (1 point) $\lim_{y \to 0} \left\{ \frac{3}{y} - \frac{3}{y^2 + y} \right\}$

Final Answer



(1d) (1 point) $\lim_{t \to 0} \frac{\sin(4t)\cos(t^2)}{\sin(7t)}$

Final Answer



(1e) (1 point) $\lim_{x \to -\infty} \frac{4x^2 + \sqrt{4x^6 + x}}{2x^3 - 2}$

Final Answer



(1f) (1 point) $\lim_{\theta \to \infty} \frac{\sin\theta}{\theta}$

Final Answer



studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by



2. For each of the following questions, only your final answer will be graded.

No justification is necessary.

(2a) (1 point) Let f be a function defined on $\mathbb{R}$. Assume we know that

$$\forall x \in \mathbb{R}, 0 < |x-4| < 0.2 \implies |f(x)-7| < 0.4$$

If the limit $L = \lim_{x \to 4} f(x)$ exists then what are the possible values of L?


Final Answer

[redacted]

(2b) (5 points) Sketch the graph of a function g defined on $(-5, 5)$which satisfies the following properties at once:

- $\lim_{x \to 2} g(x) = 3$
- $\lim_{x \to 2} g(g(x)) = -1$
- $\lim_{x \to -2} g(x)$ does not exist
- $\lim_{x \to -2} [g(x)]^2$ exists
- g has a removable discontinuity at $x = 0$.

[redacted]



studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by

3. For each of the following questions, only your final answer will be graded.

No justification is necessary.

(3a) (1 point) Write down the set of positive odd integers using set building notation.

Final Answer

[redacted]

(3b) (1 point) Write down the sentence which is the negation of the following statement:

On every page of the phone book, there is a phone number whose first digit is 2.

Final Answer

[redacted]

(3c) (6 points) For each statement, determine whether it is TRUE or FALSE. Circle the correct answer. Again, no justification is necessary. Here, we define $\mathbb{N} = \{0, 1, 2, 3, \cdots\}$.

i) $\forall x, y \in \mathbb{Z}, \exists z \in \mathbb{N}$ s.t. $x + y < z$ [redacted] FALSE

ii) $\exists x, y \in \mathbb{Z}$ s.t.$\forall z \in \mathbb{N}$, $x + y < z$ [redacted] FALSE

iii)IF $|f|$ is continuous at a THEN f is continuous at a. TRUE [redacted]

iv)IF there exists $x \in \mathbb{R}$ s.t.$x^2 < 0$ THEN $3 < 2$. [redacted] FALSE

v) Let f and g be functions defined on $\mathbb{R}$ and let $a \in \mathbb{R}$. TRUE [redacted]
IF $\lim_{x \to a} f(x) = 0$ THEN $\lim_{x \to a} f(x)g(x) = 0$

vi) Let f, g, and h be functions defined on $\mathbb{R}$ and let $a \in \mathbb{R}$. TRUE [redacted]
IF

- $\forall x \in \mathbb{R}$, $f(x) < g(x) < h(x)$
- $\lim_{x \to a} f(x)$ exists and $\lim_{x \to a} h(x) = \infty$

THEN $\lim_{x \to a} g(x)$ exists or is ∞.

studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by



4. (4a) (2 points) Let $a, L \in \mathbb{R}$. Let f be a function defined on $\mathbb{R}$. Write the formal definition of $\lim_{x \to a} f(x) = L$.

[redacted]

(4b) (4 points) Use the formal ϵ-δ definition of the limit to prove

$$\lim_{x \to 1} \frac{3x+2}{2x+3} = 1$$

Do not use any limit laws or any other theorems.

[redacted]

studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by



5. (6 points) Let f be a function defined on an interval centered at 4.

(a) (2 points) Write the formal ϵ-δ definition of " f is continuous at 4".



(b) (4 points) Assume f is continuous at 4 and $f(4) > 0$. Prove that there exists an open interval I, centered at 4, such that

$$\forall x \in I, f(x) > \frac{f(4)}{2}$$

Write a proof directly from the above formal ϵ-δ definition of continuity. Do not use any of the limit laws or any other theorems.



studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by



6. Let f and g be two strictly positive functions with domain $[1, \infty)$. We say g *controls* f if

$$\exists C > 0 \text{ s.t. } \forall a, b \in [1, \infty), |f(a) - f(b)| \leq C|g(a) - g(b)|$$

Define functions A and B by $A(x) = 3x$ and $B(x) = x^2$ for all $x \in [1, \infty)$.

(6a) (3 points) Show that B controls A.

[redacted]

(6b) (3 points) Show that A does NOT control B.

[redacted]

studocu

messages.pdf_cover_qr_code_label
messages.downloaded_by

