1. The probabili	ty of a leap ye	ar selected at r	andom contain 53			
Sunday is:						
(a) 53/ 366	(b) 1/7	(c) 2/7	(d) 53/365			
2. A bag contains	s 3 red and 2	blue marbles. A	marble is drawn at			
random. The prol	bability of dra	wing a black ba	ll is :			
(a) 3/5	(b) 2/5	(c) 0/5	(d) 1/5			
3. The probabilit	y that it will ra	ain tomorrow is	0.85. What is the			
probability that it	will not rain t	omorrow				
(a) 0.25	(b) 0.145	(c) 3/20	(d) none of these			
4. What is the pr	obability that	a number selec	ted from the numbers			
(1, 2, 3,,15	) is a multiple	of 4?				
(a) 1/5	(b) 4/5	(c) 2/15	(d) 1/3			
5. What are the	total outcome	es when we thro	w three coins?			
		(c) 8				
6. The probability	ty that a prime	e number select	ted at random from the			
numbers (1,2,3, .						
1 .			5 (d) none of these			
7. The sum of th						
		) 0 (d) no				
8. The following probabilities are given; choose the correct answer						
for that which is						
			(d) none of these.			
		nult <mark>ane</mark> ously, th	nan the probability of			
getting at least tw	wo heads, is:		4 -			
		(c) ½				
10. A letter is ch						
			e letter chosen has:			
(a) 6/13	(b) 7/13	(c) 1	(d) none of these.			
No. 100 No. 10			etting an even number.			
(A) 2/3	(B) 1	(C) 5/6	(D) 1/2			
12 Two soins are	ماه هم مرسواه م	o como timo. Fi	nd the muchability of			
12. Two coins are thrown at the same time. Find the probability of						
getting both head		(ים) ח				
(A) 3/4 (B) 1/4	(6) 1/2	(D) 0				
13. Two dice are thrown simultaneously. The probability of getting a						
sum of 9 is:		carreducty. The	probability or yetting a			

14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.						
	(B) 27/50	(C) 1	/4	(D) 29/100		
of drawing blue balls in	a blue ball is on a bag is:	double that	t of a red bal	alls .If the proba I, then the numb	•	
(A) 5	(B) 10	(C) 15	(D) 20			
	t random fron			lbs. One bulb is bability that it is		
(A) 143/150	0 (B) 14	7/150	(C) 1/25	(D) 1/50		
mixed thore the probabi		ard is draw umber on o	n from this b card is a peri	-		
<b>18. What is</b> (A) 1/7	_		_	<b>ys in a leap yea</b> D) 7/366	r?	
probability	of getting a k	ing of red	suit.	f 52 cards. Find	the	
equally like	ly to come to ,then the prol	rest pointi	ng to one of t it will point	nrrow which is the number to an odd numb ) 5/12	oer is:	
its outcome result i.e. the probability	e each time. 🛭	Aryan wins three tails Il lose the (	if all the toss and loses of	n 3 times and no ses give the sam therwise. Then t	ne	

2

(A) 1/10 (B) 3/10 (C) 1/9 (D) 4/9

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:							
(A) 364/365	(B) 31/36	51/365	(D) 1/133225				
	obability that x	<sup>2</sup> < 2 is?	e numbers -2, -1, 0 , 1,				
a marble is dra red is 2/3, then	awn at random	from the jar, the f white marbles	and others are white. If e probability that it is in the jar is:				
Then the proba		a multiple of 3	st 50 natural numbers. and 4 is:				
	owing up is pro s is?		at probability of a face The probability of face				
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	(d) $\frac{4}{21}$				
	_	_	atches are 50, 70, 82,				
	e standard dev b) 25.49	r <b>iation is</b> . c) 25.29					
	28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.						
		c) 18, 15	d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
a) $^{1}/_{2}$	b) $^{1}/_{3}$	$\frac{c}{1}$ <sub>4</sub>					
	ite between <b>0</b> a ) 7	and <b>3. The value</b> c) 27	of E(X <sup>2</sup> ) is				
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let $Z = 5X-2Y$ . The variance of Z is?							

a) 3	b) 4	c)	5	C	1) 7		
32.Out of the following values, which one is not possible in probability?  a) $P(x) = 1$ b) $\sum x P(x) = 3$ c) $P(x) = 0.5$ d) $P(x) = -0.5$							
c) $P(x) = 0.5$		$\frac{d}{d}$ P(x) =	_ 0.5				
33.lf E(x) = a) 2	<b>2 and l</b> b) 6	E(z) = 4, the	<b>en E(z – x</b> c) 0	) =?	d) Insu	fficient data	
34.The cova	ariance	of two ind	ependent	random	variable	is	
a) 1	b) 0		c) - 1		d) Und	efined	
<b>35.lf Σ P(x)</b> a) 0	= <b>k</b> <sup>2</sup> - b) 1		value of l	k is?	d) Insu	ıfficient data	
<b>36.If P(x) =</b> (a) 1		17.0	n E(x) = ? c) 4		d) 2		
37.In a disci is always?	rete pro	bability di	stribution	, the sun	ո of all p	probabilities	
a) 0	b) Infin	ite	c) 1		d) Unde	fined	
38.If the provariance.	obabilit	y of hitting	g the targe	et is 0.4,	find me	an and	
a) 0.4, 0.24		b) 0.6, 0.24	ļ	c) 0.4, 0	).16	d) 0.6, 0.16	
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance?  a) 0.6, 0.24  b) 6, 2.4  c) 0.4, 0.16  d) 4, 1.6							
<b>40.</b> Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1 <b>41.</b> What is the mean and variance for standard normal distribution?							
41. What is	me me	an and Val	iance for	Standard	i norma	i uistribution?	

a) Mean is 0 and variance is 1 b) Mean is 1 and variance is 0 c) Mean is 0 and variance is co d) Mean is co and variance is 0								
	42. Variance of a random variable X is given by a) $E(X)$ b) $E(X2)$ c) $E(X2)$ – $(E(X))2$ d) $(E(X))2$							
				_	iven by 2) - (E(X))		d) (E(X))2	
44.№ a) 0	44.Mean of a constant 'a' is a) 0							
<b>45.Variance of a constant 'a' is</b> .  a) 0 b) a c) a/2 d) 1								
46. Find the mean and variance of X?								
	Х	0	1	2	3	4		
	f(x)	1/9	2/9	3/9	2/9	1/9		
a) 2, 4/3 b) 3, 4/3 c) 2, 2/3 d) 3, 2/3								
47.Find the expectation of a random variable X?								

	х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) (	).5		b) 1.5	i	c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'g' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**
- a)  $\sqrt{np}$

- b) $\sqrt{pq}$  c) (np)2 d)  $\sqrt{npq}$