

Estimating error rates for bullet comparisons in forensic science

Yawei Ge¹, Heike Hofmann¹

^a*Department, Street, City, State, Zip*

^b*Department, Street, City, State, Zip*

Abstract

This is the abstract.

It consists of two paragraphs.

Text based on elsarticle sample manuscript, see <http://www.elsevier.com/author-schemas/latex-instructions#elsarticle>

Introduction and literature review

The firearm examiners are focusing on the same source and different source problems of bullets and cartridge cases which serve as important forensic evidence in the court/to the jurors. The subjectivity in the traditional forensic process is called to be reduced or complemented by more objective methods [The paper]. Some automatic matching algorithms are developed which usually return a similarity score (0-1) to quantify the similarity or the probability to be an actual match for a certain comparison. However, this raises questions about how to interpret the reported probabilities and how these probabilities are distributed. Thus, it is not all clear how to conduct inferences based on the similarity scores. [The paper] proposed binomial and beta-binomial for the number of matched cells of the CMC method. Thus, it provides a way to quantify the theoretical error rate of the algorithm. However, for the bullets LEA comparison, we haven't established similar distributional results. In this paper, we will evaluate the possible models/distributions for the LEA comparisons scores generated by the random forest proposed by Hare etc. And then, we will also evaluate the error rates based on the estimated distribution for the automatic matching algorithm.

In section 2, we will discuss proper distributional forms of the RF scores. In section 3, we will introduce the LAPD and the estimated distributions. In section 4, the theoretical error rates based on the distributions are discussed. In section 5, we evaluate the performance of the estimation, stability of the distribution within a changing sample size context. In section 6, we will conclude the discussion.

Email addresses: yaweige@iastate.edu (Yawei Ge), hofmann@iastate.edu (Heike Hofmann)

The distributional forms of the similarity scores

The quantitative methods used to objectively measure the similarity between LEAs reports various quantities, such as counts, correlations, distances, probabilities and more generally similarity scores references. To understand how those quantities reflect the strength of evidence and to study the underlying the error rates in making decisions based on those quantities, distributional forms are usually set up references. Particularly, we are focusing on the similarity scores which range from 0 to 1 and the probabilities reported as the likelihood of an actual match. The similarity scores reported in the forensic researches are usually categorized into two categories. One is that the compared LEAs or other forensic evidences are actual matches, the other is that the compared LEAs or other forensic evidences are not matches. We name the former as known matches (KM) and the latter as known non-matches (KNM), and the corresponding distributions are named as KM distributions and KNM distributions [figures]. However, this kind of classifications can be investigated only in the lab environment where the ground truth is known. When we make any decisions based on any quantitative measurement, we are actually making a distinguish between those two potential distributions. The strength of any identification process is also measured by the disparities of those distributions. However, in practice, we can hardly discriminate those two distributions totally, thus, we are never 100% sure which distribution the observed score comes from. This is where the identification error raises.

For the purpose of illustration, we have a look at the examples (could use Hamby and other sets, not necessarily LAPD) [figure(histogram?)]. Those are RF scores generated from the automatic matching algorithm references. The scores from the RF can be explained as probabilities that calculated through the algorithm based on the trained model to quantify the likelihood that a pair of LEAs are actual a match. Or we can think of the RF scores as general similarity scores that quantify the similarities. As the name indicated, the higher RF scores imply higher chances to be a match. For different combination of ammunition and firearms, the scores are distributed differently. It is expected that systematic differences exist there for different cases. So, it is necessary to study the scores under controlled conditions, thus, a threshold for the scores to do classifications varies based on the changes of the underlying distributions.

We can see from the figure [figure], which is a typical one we usually have for the scores, that the distributions are apart for the majorities. In the bullet LEA comparison, we usually have a well separated bullet scores but for the land scores, there are usually some overlaps [Susan]. We propose beta distributions for those scores. Because the beta distribution is a well-used one in statistics to describe a quantity from 0 to 1 which usually a probability for another distribution in Bayesian analysis. And it is very flexible to capture any unimodal shape in 0 to 1. However, it may not be adequate explain a very heavy tail or even a second mode. Thus, we also consider the beta mixture distribution which is a more complex distribution than the beta distribution. For the beta mixture distribution, we introduce a hierarchical structure with a prior probability to

combine two beta distributions as one. The distributions are stated as follows:
(formal definition with notations, assumptions, conditions for beta and beta-mix distribution)

LAPD data set and the estimated distributions

For the following sections of the paper, we will base our analysis on the LAPD data sets. This is a large data set of ... It is the first time a such large data set available to the researcher, which makes it possible for a statistical analysis for the distribution of the RF scores. (More!)

The estimation was done using EM algorithm implemented in “betareg” package in R [reference].

The Elsevier article class

Installation. If the document class *elsarticle* is not available on your computer, you can download and install the system package *texlive-publishers* (Linux) or install the LaTeX package *elsarticle* using the package manager of your TeX installation, which is typically TeX Live or MikTeX.

Usage. Once the package is properly installed, you can use the document class *elsarticle* to create a manuscript. Please make sure that your manuscript follows the guidelines in the Guide for Authors of the relevant journal. It is not necessary to typeset your manuscript in exactly the same way as an article, unless you are submitting to a camera-ready copy (CRC) journal.

Functionality. The Elsevier article class is based on the standard article class and supports almost all of the functionality of that class. In addition, it features commands and options to format the

- document style
- baselineskip
- front matter
- keywords and MSC codes
- theorems, definitions and proofs
- labes of enumerations
- citation style and labeling.

Front matter

The author names and affiliations could be formatted in two ways:

- (1) Group the authors per affiliation.
- (2) Use footnotes to indicate the affiliations.

See the front matter of this document for examples. You are recommended to conform your choice to the journal you are submitting to.

Bibliography styles

There are various bibliography styles available. You can select the style of your choice in the preamble of this document. These styles are Elsevier styles based on standard styles like Harvard and Vancouver. Please use BibTeX to generate your bibliography and include DOIs whenever available.

Here are two sample references: Feynman and Vernon Jr. (1963; Dirac, 1953).

References

Dirac, P.A.M., 1953. The lorentz transformation and absolute time. *Physica* 19, 888–896. doi:10.1016/S0031-8914(53)80099-6

Feynman, R.P., Vernon Jr., F.L., 1963. The theory of a general quantum system interacting with a linear dissipative system. *Annals of Physics* 24, 118–173. doi:10.1016/0003-4916(63)90068-X