1

Are formed by more cells, grasping in colonies and live in the water. They absorb and expel water by means of special pores Sponges are porifera. The name comes from the Ancient Greek for "pore bearer".

PORIFERA

2

Have a saclike body with a single opening (mouth) used for eating and eliminating the waste. Corals and sea anemones are also coelenterates.
The name coelenterate means hallow intestine in Latin. Cnidarian comes from the Latin word for nettle.

COELENTARATES or CNIDARIANS

3

Primate from the Latin *Primus* "prime, first rank" constituting the taxonomic order Primates. Primates arose 85–55 million years ago first from small terrestrial mammals. Many primate characteristics represent adaptations to life in this challenging environment, including large brains, color vision, and dextrous hands.

PRIMATES

4

The order Insectivora (from Latin insectum "insect" and vorare "to eat") is a now abandoned biological grouping within the class of mammals.

INSECTIVORA

5

Carnivora is an order of placental mammals that have specialized in primarily eating flesh. Its members are formally referred to as carnivorans, though some species are omnivorous, like raccoons and bears. The name comes from the Latin for "flesh devouring".

CARNIVORA

6

The annelids (Annelida, from Latin anellus, "little ring"), also known as the ringed worms or segmented worms, are a large phylum, with over 22,000 extant species including ragworms, earthworms, and leeches.

ANNELIDA

7

Mollusca is the second-largest phylum of invertebrate animals after the Arthropoda. The members are known as molluscs or mollusk. Mussels are part of the mollusca phylum.

MOLUSCA

8

Ganoids area a group of mostly extinct fishes characterized by hard, bony scales, the living species of which include the paddlefishes, sturgeons, and gars. The name means bright scales.

GANOIDS

9

Sharks are a group of fish characterized by a cartilaginous skeleton, five to seven gill slits on the sides of the head, and pectoral fins that are not fused to the head. The name comes from the old Ancient Greek *selakhos*: shark.

SELACHIANS

10

Rodents (from Latin rodere, "to gnaw") are mammals of the order Rodentia, which are characterized by a single pair of continuously growing incisors in each of the upper and lower jaws. About 40% of all mammal species are rodents (2,542 species); they are found in vast numbers on all continents except Antarctica.

RODENTIA

11

Protozoa is an informal term for a group of single-celled eukaryotes, either free-living or parasitic, which feed on organic matter such as other microorganisms or organic tissues and debris. The name comes from the Ancient Greek for *"primitve animal"*.

PROTOZOA

12

Ungulates are members of a diverse clade of primarily large mammals with hooves. These include odd-toed ungulates such as horses, rhinoceroses and tapirs, and even-toed ungulates such as cattle, pigs, giraffes, camels, sheep, deer, and hippopotamuses. The name means *hoofed animal.*

UNGULATES

13

Galliformes is an order of heavy-bodied ground-feeding birds that includes turkey, grouse and chicken. The name derives from "gallus", Latin for "cock" or "rooster".

GALLIFORMES

14

Cetaceans from Latin: cetus, lit. 'whale', from Ancient Greek: kētos, lit. 'huge fish' are aquatic mammals constituting the infraorder Cetacea. There are around 89 living species, including the dolphin , killer whales, porpoise, beluga whale, narwhal, sperm whale, and beaked whale.

CETACEA

15

Marsupials are any members of the mammalian infraclass Marsupialia. All extant marsupials are endemic to Australasia and the Americas. A distinctive characteristic common to most of these species is that the young are carried in a pouch. Well-known marsupials include kangaroos, wallabies, and koalas.

MARSUPIALIA

16

The name Pinnipedia means "fin footed". Pinnipeds, commonly known as seals, are a widely distributed and diverse clade of carnivorous, fin-footed, semiaquatic marine mammals. This clad includes the walrus, stellar sea lion, grey seal and the elephant seal.

PINNIPEDIA

17

Bats are mammals of the order Chiroptera. With their forelimbs adapted as wings, they are the only mammals capable of true and sustained flight. Bats are more manoeuvrable than birds, flying with their very long spread-out digits covered with a thin membrane or patagium. The name comes from the Greek for *"chiro + ptera "*- hand wings.

CHIROPTERA

18

An order of insects that comprises the beetles (including weevils), forming the largest order of animals on the earth. The name comes from the from Greek koleopteros, from koleos 'sheath' + pteron 'wing'. Beetles are included in this order.

COLEOPTERA

19

Teleostei (Greek: teleios, "complete" + osteon, "bone") are by far the largest infraclass in the class Actinopterygii, the ray-finned fishes,[a] and make up 96% of all extant species of fish. Tuna fish are included in this group.

TELEOSTEI

20

A passerine is any bird of the order Passeriformes, which includes more than half of all bird species. Sometimes known as perching birds or songbirds, passerines are distinguished from other orders of birds by the arrangement of their toes (three pointing forward and one back), which facilitates perching.

PASSERIFORMES

21

Are formed by more cells, grasping in colonies and live in the water. They absorb and expel water by means of special pores.

FALCONIFORMES

22

An order of tailless amphibians that comprises the frogs and toads. The name comes from the Latin an-'without' + Greek oura 'tail'.

ANURA

23

An order of amphibians that comprises the newts and salamanders, which retain the tail as adults. The name comes from the Latin uro-'tail' + Greek dēlos 'evident'.

URODELA

24

The name Echinodermata comes from Ancient Greek, echinos – "hedgehog" and derma – "skin" . The adults are recognizable by their (usually five-point) radial symmetry, and include starfish, sea urchins, sand dollars, and sea cucumbers, as well as the sea lilies or "stone lilies".

ECHINODERMATA

25

Snakes are elongated, legless, carnivorous reptiles of the suborder Serpentes. Living snakes are found on every continent except Antarctica. Additionally, sea snakes are widespread throughout the Indian and Pacific Oceans.

SERPENTES

26

Myriapoda is a subphylum of arthropods containing millipedes, centipedes, and others. The group contains over 16,000 species, most of which are terrestrial. Myriapods have a single pair of antennae and, in most cases, simple eyes.The name comes from the Latin for "thousand feet".

MYRIAPODA

27

A subclass of reptiles first appearing in the early Permian, comprising reptiles with two temporal openings and a scaly skin, and including modern lizards and tuataras. The name comes from the Greek for "scaled lizzards".

LEPIDOSAURIA

28

Arachnida is a class of joint-legged invertebrate animals (arthropods).Almost all adult arachnids have eight legs, although the front pair of legs in some species has converted to a sensory function, .The term is derived from the Greek word arachne from the myth of the hubristic human weaver Arachne, who was turned into a spider.

ARACHNIDA

29

The Hemiptera or true bugs are an order of insects comprising some 50,000 to 80,000 species of groups such as the cicadas, aphids, planthoppers, leafhoppers, and shield bugs. Most hemipterans feed on plants, using their sucking and piercing mouthparts to extract plant sap. The name comes from the Greek for half wings.

HEMIPTERA

30

Order or other division of Reptilia comprising the turtles and tortoises and being distinguished by a trunk more or less enclosed in a shell of bony dermal plates that is usually covered externally with horny shields and in nearly all cases firmly united with some of the vertebrae, ribs, and sternum, and by jaws that are toothless and sheathed.

TESTUDINES

31

Like other arthropods, crustaceans have an exoskeleton, which they moult to grow. The group has an extensive fossil record, reaching back to the Cambrian, and includes living fossils such as *Triops cancriformis*. More than 7.9 million tons of crustaceans per year are produced by fishery or farming for human consumption.

CRUSTACEA

32

Order of mostly large, predatory, semiaquatic reptiles, known as crocodilians. They first appeared 95 million years ago in the Late Cretaceous period (Cenomanian stage) and are the closest living relatives of birds, as the two groups are the only known survivors of the Archosauria.

CROCODILIA

33

Order two winged insects, the name being derived from the Greek di- "two", and pteron "wing". Insects of this order use only a single pair of wings to fly, the hindwings having evolved into advanced mechanosensory organs known as halteres, which allow dipterans to perform advanced aerobatics.

DIPTERA

34

Order of flying insects that includes the dragonflies and damselflies. Like most other flying insects they evolved in the early Mesozoic era. Fabricius coined the term Odonata from the Ancient Greek *odous* for 'tooth' apparently because they have teeth on their mandibles, even though most insects also have toothed mandibles.

ODONATA

35

Dictyoptera (from Greek diktyon "net" and pteron "wing") is an insect order that includes insects like the cockroaches.

DICTYOPTERA

36

Order of chiefly tropical fish-eating wading birds including the herons, storks, spoonbills, flamingos, and related forms all relatively large and usually with long legs and bills, upright carriage, and excepting the flamingos unwebbed feet. The name comes from the Latin for “stork like”.

CICONIIFORMES

37

Order of flightless birds, containing the extant ostriches and their extinct relatives. The name comes from the Latin for "ostrich like".

STRUTHIONIFORMES

38

Order of land birds with four unwebbed toes, short legs, small heads, and usually little visible difference between the sexes that includes the sandgrouse, the pigeons and doves. The name comes from the Latin for "pigeon like".

COLUMBIFORMES

39

Are formed by more cells, grasping in colonies and live in the water. They absorb and expel water by means of special pores.

MONOTREMATA

40

Order of birds comprising the ducks, geese, swans, and mergansers and the screamers. Most modern species in the order are highly adapted for an aquatic existence at the water surface. The name comes from the Latin for "goose like".

ANSERIFORMES

41

Highly specialized insects with complete metamorphosis that include the bees, wasps, ants, ichneumon flies, sawflies, gall wasps, and related forms, often associate in large colonies with complex social organization, and have usually four membranous wings. The name comes from the Greek *hymeno+ptera* for membrane-winged.

HYMENOPTERA

42

An order of nonpasserine birds formerly restricted to the woodpeckers but now usually including also the jacamars, puffbirds, barbets, honey guides, and toucans

PICIFORMES

43

Lepidoptera (from Ancient Greek lepís "scale" + pterón "wing") is an order of insects that includes butterflies and moths. Their bodies and wings are generally covered with scales.

LEPIDOPTERA

44

The Sirenia, commonly referred to as sea-cows or sirenians, are an order of fully aquatic, herbivorous mammals that inhabit swamps, rivers, estuaries, marine wetlands, and coastal marine waters. The name (Greek for "mermaid") comes from a legend about their discovery, involving lonely sailors mistaking them for mermaids.

SIRENIA

45

The order Pilosa, the word *pilosus* means "hairy" in Latin, is a group of placental mammals, found today only in the Central and South America. It includes the anteaters and sloths.

PILOSA

46

Dipnoi subclass (from the Greek for "two ways to breath") contains animals that have 2 ways of breathing. With the gills they take the oxygen from the water like the fish, and with the lung they take it from the air. If they end up in the mud they can breath the air until the water returns. These animals are mostly found in the Southern hemisphere.

DIPNOI

A

An arthropod (from the Greek for "joint" and "foot") is an invertebrate animal having an exoskeleton, a segmented body, and paired jointed appendages.

ARTHROPODA

B

Insects or Insecta (from Latin insectum for "cut body") are hexapod invertebrates and the largest group within the arthropod phylum. Nearlly all insects hatch from eggs and have the body divided in three parts: head, thorax and abdomen.

INSECTS

Animals with few exceptions consume organic material, breathe oxygen, are able to move and can reproduce sexually. 1.5 million animals have been described - of which around 1 million are insects - but it has been estimated that there are over 7 million species in total.

ANIMALIA

I

Invertebrates are animals that neither possess nor develop a vertebral column (commonly known as a backbone or spine), derived from the notochord.

INVERTEBRATES

II

Vertebrates are animals that possess or develop a vertebral column (commonly known as backbone or spine).

VERTEBRATES

A

FIsh are cold-blooded, most of them have a backbone and fins, breathe with gills and have scales that cover their bodies. It is currently believed that fish began to evolve about 480 million years ago. There are about 22,000 known species of fish.

FISHES

B

One thing that most amphibians do have in common is a moist skin, often coated with a slimy mucus. Adult amphibians breathe not only through their lungs but also through their skin, and the moisture is necessary for proper oxygen exchange. The life span of amphibians varies widely, from a few months to many years.

AMPHIBIA

C

Reptiles are air-breathing vertebrates covered in special skin made up of scales, bony plates, or a combination of both. Most of them are cold bloded. Reptile reproduction also depends on temperature. Only boas and pythons give birth to live young. The other species lay their eggs in a simple nest, and leave.

REPTILIA

D

Birds are a group of warm-blooded vertebrates constituting the class Aves, characterized by feathers, toothless beaked jaws, the laying of hard-shelled eggs, a high metabolic rate, a four-chambered heart, and a strong yet lightweight skeleton.

AVES

E

Mammal is any member of the group of vertebrate animals in which the young are nourished with milk from special mammary glands of the mother. Hair is a typical mammalian feature, although in many whales it has disappeared except in the fetal stage. Mammals are warm-blooded.

MAMMALIA