

THE EFFECT OF DANAZOL AND THE LHRH AGONIST ANALOGUE GOSERELIN (ZOLADEX) ON THE BIOLOGICAL ACTIVITY OF LUTEINIZING HORMONE IN WOMEN WITH ENDOMETRIOSIS

P. MAOURIS, M. DOWSETT, GILLIAN ROSE, D. K. EDMONDS,
CLAIRE ROTHWELL AND W. R. ROBERTSON

Queen Charlotte's and Chelsea Hospital, London, Department of Biochemical Endocrinology, Royal Marsden Hospital, Fulham Road, London and Department of Medicine (Clinical Biochemistry), University of Manchester, Hope Hospital, Salford M6 8HD, UK

(Received 3 January 1990; returned for revision 22 February 1990; finally revised 22 May 1990; accepted 12 June 1990)

SUMMARY

In an attempt to determine whether the suppression in oestradiol levels caused by danazol is due to an effect on the hypothalamic–pituitary axis, we compared the endocrine effects of danazol with those of the LHRH (GnRH) agonist analogue goserelin. Serum levels of immunoreactive LH (I-LH), FSH, 17β -oestradiol (E_2) and bioactive LH (B-LH) (using a mouse Leydig cell bioassay), were measured in ten and 20 women with endometriosis treated with danazol and goserelin, respectively. I-LH was measured both by radioimmunoassay (RIA) and immunoradiometric assay (IRMA). During 6 months of treatment with 600 mg of danazol daily, mean serum E_2 decreased ($P < 0.05$) to levels near the upper limit of the post-menopausal range (to a mean (and 95% confidence interval of the mean) of 117 (65–169) pmol/l) whereas FSH, I-LH (both by RIA and IRMA) and B-LH levels were not significantly altered. During 6 months of treatment with monthly depot injections of 3.6 mg goserelin, mean serum E_2 decreased ($P < 0.001$) to well within the post-menopausal range (to 23 (18–28) pmol/l). The mean FSH, I-LH and B-LH levels also decreased ($P < 0.05$) during therapy with goserelin (from 3.9 (3.1–4.7) to 2.0 (1.6–2.4) IU/l for FSH, from 5.3 (4.5–6.1) to 1.9 (1.7–2.1) IU/l for RIA-LH, from 2.9 (2.5–3.3) to < 0.5 (< 0.5) IU/l for IRMA-LH and from 9.1 (7.1–11.1) to 2.9 (2.6–3.2) IU/l for B-LH). The bioactive to immunoreactive (B:I) RIA LH ratio was altered during the first month of treatment with both drugs, increasing with danazol from 1.6 (1.3–1.9) to 2.0 (1.7–2.3) ($P < 0.05$) and decreasing with goserelin from 1.7 (1.5–1.9) to 1.4 (1.2–1.6) ($P < 0.05$). The data indicate that danazol causes a relatively hypo-oestrogenic state which cannot be attributed to a decrease in the immuno-

Correspondence: Dr P. Maouris, North Tees General Hospital, Stockton, Cleveland TS19 8PE, UK.

reactive gonadotrophin levels or to a decrease in the biological activity of LH and is associated with an increase in B:I LH ratio. In contrast, goserelin acts on the pituitary to produce suppression of gonadotrophin levels and effective down-regulation of the pituitary.

Danazol and luteinizing hormone-releasing hormone (LHRH) agonist analogues have similar efficacy in the treatment of endometriosis (*Lancet*, 1986; Henzl *et al.*, 1988; Fraser & Waxman, 1989). They both cause amenorrhoea, anovulation, and a hypo-oestrogenic state which makes them suitable agents for use in conditions which are oestrogen dependent or associated with menorrhagia and/or dysmenorrhoea (like endometriosis, fibromyomata, premenstrual syndrome and dysfunctional uterine bleeding).

Danazol, an isoxazole derivative of 17 α -ethinyltestosterone, was originally described as having strong antigonadotrophic properties (Dmowski, 1979), but current evidence suggests that whilst it can eliminate the midcycle LH and FSH surge, danazol does not alter basal gonadotrophin levels (Barbieri & Ryan, 1981). However, E₂ levels are suppressed. Although there is evidence that danazol suppresses ovarian steroidogenesis (Barbieri *et al.*, 1977; Nilsson *et al.*, 1984; Carlstrom *et al.*, 1984; Olsson *et al.*, 1986), exogenous LHRH stimulation in danazol-treated patients leads to normal secretion of radioimmunoassayed gonadotrophins and E₂ (Rannevik, 1979; Braun *et al.*, 1983).

LHRH agonist analogues cause an initial stimulation of gonadotrophin and gonadal sex steroid secretion followed by a desensitization of the pituitary. This leads to suppression of gonadotrophin secretion and a reduction in ovarian steroidogenesis with a resultant decrease in serum oestrogen concentrations to levels which are similar to those found in women after the menopause or after surgical oophorectomy (Fraser & Waxman, 1989). Thus decreased I-LH levels have been widely reported in patients on LHRH agonist therapy (Schriock *et al.*, 1985; Thomas *et al.*, 1986; West & Baird, 1987; Matta *et al.*, 1988; Nicholson *et al.*, 1987), but others have noted increased levels of I-LH which have been reported to last up to 24 weeks (Meldrum *et al.*, 1982; Lemay *et al.*, 1984). These increased levels are associated with an increased release of LH α -subunit or of LH fragments (Lemay *et al.*, 1987; Meldrum *et al.*, 1984) which may cross-react with LH radioimmunoassays giving an erroneously high I-LH estimation. Furthermore, characterization of the serum I-LH by Sephadex chromatography demonstrated a change in the elution profile of I-LH on treatment with LHRH agonists indicating that LHRH agonists may induce the pituitary to release an immunoreactive LH with diminished bioactivity (Meldrum *et al.*, 1984; Evans *et al.*, 1984).

In an attempt to clarify these findings, and in particular to determine whether the suppression in oestradiol levels caused by danazol is due to an effect on the hypothalamic-pituitary axis, we compared the endocrine effects of danazol with those of the LHRH agonist analogue goserelin which is known to cause a state of hypogonadotrophic hypogonadism consistent with complete ovarian ablation (Thomas *et al.*, 1986; Nicholson *et al.*, 1987; Matta *et al.*, 1988).

MATERIALS AND METHODS

Subjects

Thirty normally menstruating women aged between 25 and 40 years (mean, 31 years) with laparoscopically diagnosed endometriosis were randomly allocated to 6 months of

medical therapy with either danazol 600 mg daily taken orally in three divided doses, or monthly subcutaneous injections of 3.6 mg of goserelin (D-SER [tBU]⁶, Aza-GLY¹⁰-LHRH) in the ratio of 1:2, respectively (10 patients received danazol and 20 patients received goserelin). All patients gave written consent and the study was approved by the hospital ethical committee. A blood sample was obtained from each patient during the early follicular phase (days 1–7) of the pretreatment cycle, at months 1 and 3 of treatment and 2 months post-treatment (i.e. 2 months after the last danazol capsule and 3 months after the last injection of goserelin). Serum was stored at –20°C until analysis.

Immunoassays

Serum LH, FSH and E₂ were analysed by radioimmunoassay using methods previously described (Ferguson *et al.*, 1982 (LH and FSH); Dowsett *et al.*, 1987 (E₂)). The within and between assay coefficients of variation (CV) were: for LH 6 and 12% at 4.8 IU/l respectively, with a sensitivity of 0.8 IU/l; for FSH 5 and 8% at 1.6 IU/l respectively, with a sensitivity of 0.4 IU/l, and for E₂ 9 and 13% at 36 pmol/l respectively, with a sensitivity of 6 pmol/l.

LH was also measured using an immunoradiometric assay (Serono MAIAclone). The within and between assay CVs were 5 and 8% respectively, with a sensitivity of 0.5 IU/l.

LH Bioassay

The biological activity of LH was estimated using a dispersed mouse Leydig cell bioassay (Van Damme *et al.*, 1974) modified as previously described (Tsatsoulis *et al.*, 1987). The same pituitary LH reference standard (68/40) was used in this assay as for the RIA and IRMA LH assays. The within and between assay CVs were 7 and 12%, respectively, with a sensitivity of 2.5 IU/l.

Goserelin, danazol and its two major metabolites (ethisterone and 2-hydroxymethyl-ethisterone), did not interfere with the bioassay at concentrations of at least ten times greater than the expected maximum serum concentrations. Similarly, none of the drugs cross-reacted with the testosterone antibody employed in measuring testosterone secretion from Leydig cells.

Statistical analysis

Statistical analysis of E₂, FSH, I-LH (RIA and IRMA), B-LH, B:RIA LH ratio, B:IRMA LH ratio and RIA:IRMA LH ratio at each interval was performed using analysis of variance (ANOVA). Confirmatory *t*-tests were done using the mean squared error term from the ANOVA. Some of the B-LH and IRMA-LH assay values were less than the lower detection limit of the assay. In these cases the value allocated was 0.1 IU/l below the assay sensitivity (i.e. 2.4 IU/l for B-LH and 0.4 IU/l for IRMA-LH). This has the effect of underestimating any changes involving B-LH, IRMA-LH, B:RIA and B:IRMA LH ratios. In all analyses, *P* < 0.05 was considered significant.

RESULTS

The results are presented in Table 1.

Table 1. The effect of danazol and goserelin therapy on E₂, FSH, RIA LH, IRMA LH, B-LH, B:RIA LH and B:IRMA LH

Hormone	Danazol				Goserelin			
	Pre	M1	M3	Post	Pre	M1	M3	Post
E ₂	356	160	75	305	282	18	30	476
(pmol/l)	169; 543	55; 265	56; 94	188; 422	198; 366	14; 22	12; 48	326; 626
FSH	3.5	4.4	3.7	3.7	3.9	1.5	2.2	4.0
(IU/l)	2.3; 4.6	3.4; 5.4	2.9; 4.5	2.8; 4.5	3.1; 4.7	1.2; 1.8	1.7; 2.7	2.6; 5.4
RIA LH	6.8	7.3	6.5	7.1	5.3	2.2	1.7	10.2
(IU/l)	5.3; 8.3	5.6; 9.0	3.9; 9.1	4.2; 11.0	4.5; 6.1	2.0; 2.4	1.5; 1.9	4.6; 15.8
IRMA LH	3.9	4.6	3.7	3.8	2.9	0.5	.4	5.8
(IU/l)	3.0; 4.8	3.5; 5.7	3.1; 4.2	1.4; 6.2	2.5; 3.3	0.4; 0.6	0.4; 0.4	2.8; 8.8
B-LH	11.3	14.9	12.5	13.9	9.1	2.9	2.9	14.6
(IU/l)	7.2; 15.4	10.0; 19.8	6.6; 18.4	5.3; 22.5	7.1; 11.1	2.7; 3.1	2.6; 3.2	9.7; 19.5
B:RIA LH	1.6	2.0	1.9	1.8	1.7	1.4	1.7	1.9
	1.3; 1.9	1.7; 2.3	1.8; 2.0	1.7; 1.9	1.5; 1.9	1.2; 1.6	1.5; 2.0	1.6; 2.2
B:IRMA LH	2.9	3.3	3.4	3.7				
	2.3; 3.5	2.8; 3.8	2.9; 3.9	3.2; 4.2				

Pre, Pretreatment; M1, month 1; M3, month 3; Post, post-treatment. All values are given as mean and 95% confidence interval.

Danazol treatment led to a reduction ($P < 0.05$) in E₂ concentration at 1 and 3 months (to a mean (and 95% confidence interval of the mean) of 160 (55–265) and 75 (56–94) pmol/l, respectively), but hormone levels returned to within the pretreatment range (356 (169–543) pmol/l) 2 months after therapy was completed. Goserelin therapy led to a reduction ($P < 0.001$) of E₂ levels at 1 month to 18 (14–22) pmol/l. This level of E₂ was only 6% of the pretreatment concentrations and was far lower ($P = 0.01$) than that found with danazol. Secretion of E₂ remained severely suppressed after 3 months of therapy but in contrast to that found with danazol, increased ($P < 0.05$) to higher than pretreatment levels 2 months post-therapy.

The level of FSH was unchanged during danazol therapy whereas it decreased ($P < 0.01$) drastically during goserelin treatment, returning to pretreatment levels 2 months post-therapy.

The levels of RIA, IRMA or B-LH levels were unchanged during or after danazol treatment. The B:RIA LH ratio rose ($P < 0.05$) after 1 month of danazol treatment but returned to the upper limit of the pretreatment range at 3 months of treatment and remained within the pretreatment range 2 months post-treatment. The B:IRMA LH ratio also rose during danazol treatment but the increase was significant ($P < 0.05$) only at 3 months of treatment.

Mean RIA-LH and IRMA-LH levels fell ($P < 0.001$) during treatment with goserelin and rose after the end of therapy to above pretreatment levels ($P < 0.05$). B-LH levels also fell ($P < 0.001$) during goserelin therapy. The B:RIA LH ratio was lower ($P < 0.05$) during the first month of treatment compared to the pretreatment figure but rose to pretreatment levels by the third month of treatment with a further (not significant) rise by 2 months post-treatment. The B:IRMA LH ratio was not calculated in the case of goserelin therapy

because a large number of values in both the bioassay and the IRMA were below the lower detection limit of the assays.

For both groups, the RIA:IRMA LH ratio at pretreatment and post-treatment was similar (1.8 (1.5–2.1)). This was not altered by danazol therapy whereas goserelin therapy resulted in a greater decrease in LH IRMA compared to LH RIA values such that the RIA:IRMA LH ratio was increased to 4.4 (4.0–4.8) and 4.3 (3.9–4.7) at 1 and 3 months of treatment respectively ($P < 0.001$).

DISCUSSION

This is the first report relating bioactivity with radioimmunoactivity of LH in women treated with danazol. Comparisons were made with goserelin-treated patients since, like other LHRH agonists, this compound is known to be effective through suppression of gonadotrophin-stimulated ovarian steroidogenesis (West & Baird, 1987). We have examined differential changes in immunoreactivity and bioactivity as reflected in the B:I ratios since these ratios provide a sensitive index of qualitative changes in LH (Dufau & Veldhuis, 1987).

Endometriosis is an oestrogen-dependent condition not seen before puberty or after the menopause (in the absence of hormone replacement therapy). The desired pharmacological effect of any drug suitable for the treatment of endometriosis is, therefore, oestrogen deprivation. The marked suppression of E_2 levels seen with goserelin therapy was consistent with a complete ovarian ablation and was significantly greater than that seen with danazol. Despite this, danazol appears to have similar efficacy to LHRH agonists in the treatment of endometriosis (Lancet, 1986; Henzl *et al.*, 1988; Fraser & Waxman, 1989). The increase in free, biologically active testosterone seen with danazol therapy (Dowsett *et al.*, 1986; Forbes *et al.*, 1986) may contribute to the therapeutic efficacy of danazol. Direct inhibition of endometrial cell growth by danazol (Rose *et al.*, 1988), may also play a role.

The lack of suppression of the FSH and LH levels by danazol treatment is in agreement with Barbieri and Ryan (1981) and suggests that the observed decrease in E_2 levels is due to the inhibitory effect of danazol on ovarian steroidogenesis (Barbieri *et al.*, 1977; Nilsson *et al.*, 1984; Carlstrom *et al.*, 1984; Olsson *et al.*, 1986). This, however, does not exclude a direct effect of the drug on the hypothalamo-pituitary axis which prevents a measurable compensatory rise of gonadotrophin levels and also abolishes the mid-cycle gonadotrophin surge. The effect of danazol on LH pulsatility (Braun *et al.*, 1983; Dmowski *et al.*, 1983) may also result in reduced E_2 secretion as in women with hypothalamic dysfunction.

The marked suppression of FSH and RIA-LH levels during goserelin treatment is in agreement with most previous studies involving this agonist (Thomas *et al.*, 1986; West & Baird, 1987; Matta *et al.*, 1988; Nicholson *et al.*, 1984) and other LHRH agonists (Schriock *et al.*, 1985; Franssen *et al.*, 1989; Tummon *et al.*, 1989). The decrease in B-LH during goserelin therapy is in agreement with the findings of all the previous published studies on LH bioactivity using other LHRH agonist analogues (Meldrum *et al.*, 1984; Evans *et al.*, 1984; Chiang *et al.*, 1985;) and confirms that the decrease seen in the I-LH is of biological significance. This is reflected in the observed marked decrease in E_2 levels.

The increase in the E_2 , RIA-LH, IRMA-LH, and B-LH seen at 2 months after the end of goserelin treatment appears to be due to one-quarter of the patients being at mid-cycle at the time of the post-treatment blood sample.

The increase seen in the B:RIA LH ratio and the B:IRMA LH ratio at 1 and 3 months of danazol treatment may reflect an effect of low E₂ levels on the B:I-LH ratio similar to that seen in postmenopausal women and patients with Turner's syndrome (Dufau *et al.*, 1976), an androgenic effect of the drug as seen in studies in animals (Solano *et al.*, 1980) and in men (Beitins *et al.*, 1981; Carani *et al.*, 1987), or an effect on the LH pulsatility resulting in differential secretion of LH isohormones (Dufau & Veldhuis, 1987). In contrast, the decrease in the B:RIA LH ratio seen at 1 month on goserelin treatment is in agreement with previous studies on other LHRH agonists (Meldrum *et al.*, 1984; Chiang *et al.*, 1985; Lemay *et al.*, 1987). However, discordance between B-LH and RIA-LH was much greater in these last studies. This may be attributed to the incomplete suppression of the I-LH in those studies, the production of immunoreactive LH fragments or forms with reduced bioactivity (Meldrum *et al.*, 1984; Evans *et al.*, 1984), or differences in the polyclonal antibodies used in the RIAs.

Bischof and Herrmann (1988) demonstrated that LHRH agonist therapy in women with endometriosis produces a much greater decrease in LH as measured by an IRMA based on two different monoclonal anti-LH antibodies compared to LH measured by the RIA based on a rabbit polyclonal anti-LH antiserum.

Similarly, our data show a marked elevation in the RIA:IRMA LH ratio during goserelin therapy. This and other work (UK EQAS Report, 1988; Haavisto *et al.*, 1989) suggest that the RIA is unreliable at low levels (< 2.0 IU/l). For samples where LH levels are expected to be of a low order an IRMA rather than the RIA may be the assay of choice.

We conclude that the divergent effect of the two drugs on the B:RIA LH ratio reflects the fundamentally different mechanisms of action of these drugs on the pituitary. The data indicate that danazol causes a relatively hypo-oestrogenic state which cannot be attributed to a decrease in the immunoreactive gonadotrophin levels or to a decrease in the biological activity of LH and is associated with an increase in B:I LH ratio. In contrast, goserelin acts directly on the pituitary to produce suppression of gonadotrophin levels and effective down-regulation of the pituitary.

ACKNOWLEDGEMENTS

Winthrop Laboratories, ICI Pharmaceuticals, Serono Diagnostics and North-Western Regional Health Authority are acknowledged for financial support.

REFERENCES

- BARBIERI, R.L., CANICK, J.A., MAKIS, A., TODD, R.B., DAVIES, I.J. & RYAN, K.J. (1977) Danazol inhibits steroidogenesis. *Fertility and Sterility*, **28**, 809–813.
- BARBIERI, R.L. & RYAN, K.J. (1981) Danazol: endocrine pharmacology and therapeutic applications. *American Journal of Obstetrics and Gynecology*, **141**, 453–463.
- BEITINS, I.Z., AXELROD, L., OSTREA, T., LITTLE, R. & BADGER, T.M. (1981) Hypogonadism in a male with immunologically active, biologically inactive luteinizing hormone: characterisation of the abnormal hormone. *Journal of Clinical Endocrinology and Metabolism*, **52**, 1143–1149.

- BISCHOF, P. & HERMANN, W.L. (1988) Absence of immunoreactive luteinizing hormone following gonadotropin-releasing hormone agonist therapy in women with endometriosis. *Gynecologic and Obstetric Investigations*, **25**, 130–134.
- BRAUN, P., WILDT, L. & LEYENDECKER, G. (1983) The effect of danazol on gonadotropin secretion during the follicular phase of the menstrual cycle. *Fertility and Sterility*, **40**, 37–44.
- CARANI, C., ZINI, D., DELLA CASA, L., SYED, B., BALDINI, A. & MARRAMA, P. (1987) Chronic testosterone therapy increases biological activity of LH in men with mild hypogonadism. *First European Congress of Endocrinology, Copenhagen, 21–25 June 1987*. Abstracts (eds J. Jensen & C. Christiansen), p. 163.
- CARLSTROM, K., DOBERL, A., GERSHAGEN, S. & RANNEVIK, G. (1984) Peripheral levels of dehydroepiandrosterone sulfate, dehydroepiandrosterone, androstenedione, and testosterone following different doses of danazol. *Acta Obstetrica et Gynecologica Scandinavica*, (Suppl 123), 1265–129.
- CHIANG, R.S., BARNES, R.B., SHOUP, D. & LOBO, R.A. (1985) Dose-related changes in LH bioactivity with intranasal GnRH agonist administration. *Contraception*, **32**, 347–357.
- DMOWSKI, W.P. (1979) Endocrine properties and clinical application of danazol. *Fertility and Sterility*, **31**, 237–251.
- DMOWSKI, W.P., HEADLEY, S. & RADWANSKA, E. (1983) Effects of danazol on pulsatile gonadotropin patterns and on serum estradiol levels in normally cycling women. *Fertility and Sterility*, **39**, 49–55.
- DOWSETT, M., FORBES, K.L., ROSE, G.L., MUDGE, J.E. & JEFFCOATE, S.L. (1986) A comparison of the effects of danazol and gestrinone on testosterone binding to sex hormone binding globulin *in vitro* and *in vivo*. *Clinical Endocrinology*, **24**, 555–563.
- DOWSETT, M., GOSS, P.E., POWLES, T.J., HUTCHISON, G., BRODIE, A.M.H., JEFFCOATE, S.L. & COOMBES, R.C. (1987) Use of the aromatase inhibitor 4-hydroxyandrostenedione in postmenopausal breast cancer: optimization of therapeutic dose and route. *Cancer Research*, **47**, 1957–1961.
- DUFAU, M.L., BEITINS, I.Z., MCARTHUR, J.W. & CATT, K.J. (1976) Effects of luteinizing hormone releasing hormone (LHRH) upon bioactive and immunoreactive serum LH levels in normal subjects. *Journal of Clinical Endocrinology and Metabolism*, **43**, 658–667.
- DUFAU, M.L. & VELDHIJS, J.D. (1987) Pathophysiological relationships between the biological and immunological activities of luteinizing hormone. In *Baillière's Clinical Endocrinology and Metabolism*. Volume 1, no. 1 (ed. H.G. Burger), pp. 153–176. Baillière Tindall, London.
- EVANS, R.M., DOELLE, G.C., LINDNER, V., BRADLEY, V. & RABIN, D. (1984) A luteinizing-releasing hormone agonist decreases biological activity and modifies chromatographic behavior of luteinizing hormone in man. *Journal of Clinical Investigation*, **73**, 262–266.
- FERGUSON, K.M., HAYES, M. & JEFFCOATE, S.L. (1982) A standardised multicentre procedure for plasma gonadotrophin radioimmunoassay. *Annals of Clinical Biochemistry*, **19**, 358–361.
- FORBES, K.L., DOWSETT, M., ROSE, G., MUDGE, J.E. & JEFFCOATE, S.L. (1986) Dosage-related effects of danazol on sex hormone binding globulin and free total androgen levels. *Clinical Endocrinology*, **25**, 597–605.
- FRANSSSEN, A.M.H.W., KAUFER, F.M., CHADHA, D.R., ZIJLSTRA, J.A. & ROLLAND, R. (1989) Endometriosis: treatment with gonadotropin-releasing hormone agonist Buserelin. *Fertility and Sterility*, **51**, 401–408.
- FRASER, H.M. & WAXMAN, J. (1989) Gonadotrophin releasing hormone analogues for gynaecological disorders and infertility. *British Medical Journal*, **298**, 475–476.
- HAAVISTO, A.-M., DUNKEL, L., PETTERSON, K. & HUHTANIEMI, I. (1989) Measurement of serum LH in boys with delayed puberty by radioimmunoassay, highly sensitive immunofluorometric assay and *in vitro* bioassay. *Third International Conference on the Control of the Onset of Puberty, Amsterdam*. Abstract No. 24, p. 48.
- HENZL, M.R., CORSON, S.L., MOGHISSI, K., BUTTRAM, V.C., BERQVIST, C. & JACOBSON, J. (1988) Administration of nasal nafarelin as compared with oral danazol for endometriosis. *New England Journal of Medicine*, **318**, 485–489.
- LANCET (1986) LHRH Analogues in endometriosis. *Lancet*, **ii**, 1016–1017.
- LEMAY, A., FAURE, N. & MAHEUX, R. (1987) Loss of bioactivity and changes in immunoassayable subunits of serum LH during LHRH agonist treatment of endometriosis. Abstracts of the 43rd Annual Meeting of the American Fertility Society, September 28–30, 1987. *Fertility and Sterility (Program Supplement)*, p. 19.
- LEMAY, A., MAHEUX, R., FAURE, N., JEAN, C. & FAZEKAS, A.T.A. (1984) Reversible hypogonadism induced by a luteinizing hormone-releasing hormone (LH-RH) agonist (Buserelin) as a new therapeutic approach for endometriosis. *Fertility and Sterility*, **41**, 863–871.
- MATTA, W.H.M., SHAW, R.W. & BURFORD, G.D. (1988) Endocrinologic and clinical evaluation following a single administration of a gonadotrophin-releasing hormone agonist (Zoladex), in a depot formulation, to premenopausal women. *Fertility and Sterility*, **49**, 163–165.

- MELDRUM, D.R., CHANG, R.J., LU, J.K.H., VALE, W., RIVIER, J. & JUDD, H.L. (1982) 'Medical oophorectomy' using a long-acting GnRH agonist—A possible new approach to the treatment of endometriosis. *Journal of Clinical Endocrinology and Metabolism*, **54**, 1081–1083.
- MELDRUM, D.R., TSAO, Z., MONROE, S.E., BRAUNSTEIN, G.D., SLADEK, J., LU, J.K.H., VALE, W., RIVIER, J., JUDD, H.L. & CHANG, R.J. (1984) Stimulation of LH fragments with reduced bioactivity following GnRH agonist administration in women. *Journal of Clinical Endocrinology and Metabolism*, **58**, 755–757.
- NICHOLSON, R.I., WALKER, K.J., TURKES, A., DYAS, J.G., GOTTING, K.E., PLOWMAN, P.N., WILLIAMS, M., ELSTON, C.W. & BLAMEY, R.W. (1987) The British experience with the LH-RH agonist Zoladex (ICI 118630) in the treatment of breast cancer. In *Hormonal Manipulation of Cancer: Peptides, Growth Factors and new (anti) Steroidal Agents* (ed. J. G. M. Klijn), pp. 331–341. Raven Press, New York.
- NILSSON, L., DENNEFORS, B., OLSSON, J.-H., HILLENSJO, T. & RANNEVIK, G. (1984) In vitro effect of danazol on steroidogenesis in developing human follicles. *Acta Obstetrica et Gynecologica Scandinavica* (Suppl. 123), 99–102.
- OLSSON, J.-H., HILLENSJO, T. & NILSSON, L. (1986) Inhibitory effects of danazol on steroidogenesis in cultured human granulosa cells. *Fertility and Sterility*, **46**, 237–242.
- RANNEVIK, G. (1979) Hormonal, metabolic and clinical effects of danazol in the treatment of endometriosis. *Postgraduate Medical Journal*, **55** (Suppl. 5), 14–20.
- ROSE, G.L., DOWSETT, M., MUDGE, J.E., WHITE, J.O. & JEFFCOATE, S.L. (1988) The inhibitory effects of danazol, danazol metabolites, gestrinone, and testosterone on the growth of human endometrial cells *in vitro*. *Fertility and Sterility*, **49**, 224–228.
- SCHRIOCK, E., MONROE, S.E., HENZL, M. & JAFFE, R.B. (1985) Treatment of endometriosis with a potent agonist of gonadotropin-releasing hormone (nafarelin). *Fertility and Sterility*, **44**, 583–588.
- SOLANO, A.R., GARCIA-VELA, A., CATT, K.J. & DUFAU, M.L. (1980) Modulation of serum and pituitary luteinizing hormone bioactivity by androgen in the rat. *Endocrinology*, **106**, 1941–1948.
- THOMAS, E.J., JENKINS, J., LENTON, E.A. & COOKE, I.D. (1986) Endocrine effects of goserelin, a new depot luteinizing hormone releasing hormone agonist. *British Medical Journal*, **293**, 1407–1408.
- TSATSOULIS, A., WHITEHEAD, J., ST. JOHN, S., SHALET, M. & ROBERTSON, W.R. (1987) The pituitary–Leydig cell axis in men with severe damage to the germinal epithelium. *Clinical Endocrinology*, **27**, 683–689.
- TUMMON, I.S., PEPPING, M.E., BINOR, Z., RADWANSKA, E. & DMOWSKI, W.P. (1989) A randomized, prospective comparison of endocrine changes induced with intranasal leuprolide or danazol for treatment of endometriosis. *Fertility and Sterility*, **51**, 390–394.
- UNITED KINGDOM EXTERNAL QUALITY ASSESSMENT SCHEME FOR PEPTIDE HORMONES (1988) *Annual Review of Schemes*. Department of Clinical Chemistry, University of Edinburgh, p. 45.
- VAN DAMME, M.-P., ROBERTSON, D.M. & DICZFALUSY, E. (1974) An improved in-vitro bioassay method for measuring luteinizing hormone (LH) activity using mouse Leydig cell preparations. *Acta Endocrinologica*, **77**, 655–671.
- WEST, C.P. & BAIRD, D.T., (1987) Suppression of ovarian activity by Zoladex depot (ICI 118630), a long-acting luteinizing hormone releasing hormone agonist analogue. *Clinical Endocrinology*, **26**, 213–220.