

Poor performance of galactomannan and mannan sandwich enzyme-linked immunosorbent assays in the diagnosis of invasive fungal infection

The Platelia® *Aspergillus* and Platelia® *Candida* (BioRad Laboratories Ltd, Hertfordshire, UK) are commercially available sandwich enzyme-linked immunosorbent assay (ELISA) kits for the detection of galactomannan (GM) and mannan (M) respectively.

A number of prospective studies report excellent sensitivities (90–95%) and specificities using the Platelia® *Aspergillus* for the diagnosis of invasive aspergillosis (IA) (McLintock & Jones, 2004). The European Organisation for the Research and Treatment of Cancer and the Mycosis Study Group (EORTC/MSG) definitions of invasive fungal infection (IFI) include positive GM antigen testing as a microbiological criterion (Ascioglu *et al*, 2002). Other studies demonstrate poorer sensitivities (0–50%) (McLintock & Jones, 2004) and controversy exists over the most appropriate positive cut-off (Maertens *et al*, 2004).

There are no prospective studies which evaluate the Platelia® *Candida* and retrospective studies report modest sensitivities (30–69%) (McLintock & Jones, 2004). Positive M antigen testing is not included in EORTC/MSG definitions of IFI.

A prospective blinded study was performed to assess the value of twice-weekly screening of serum specimens, from haemato-oncology patients at high- and intermediate high-risk of IFI (Prentice *et al*, 2000), with Platelia® *Aspergillus* and Platelia® *Candida* ELISAs. Between December 2000 and December 2001, adult patients receiving allogeneic or autologous stem cell transplantation (SCT) or intensive chemotherapy were recruited. A treatment episode was defined as a single course of chemotherapy, an autologous SCT or an

allogeneic SCT until day 100 following the transplant, or a course of corticosteroid therapy for graft-versus-host disease. An investigator blinded to the patient data performed the ELISAs according to the manufacturer's instructions. Each sample was run in duplicate. Samples were re-tested if the coefficient of variation between duplicates was more than 20%. IFI was classified as proven, probable or possible according to the EORTC/MSG definitions (Ascioglu *et al*, 2002). As the Platelia® *Aspergillus* ELISA was being evaluated it was excluded from the EORTC/MSG definitions.

The Platelia® *Candida* ELISA assay was discontinued after analysis of 82 episodes as 61% of the kits failed the manufacturer's validation criteria and results were therefore uninterpretable. Additionally, approximately 30% of the available ELISA results were non-reproducible.

A total of 1625 Platelia® *Aspergillus* ELISA results, from 125 treatment episodes, involving 78 patients, aged 16–76 years (mean 42, median 44·5 years), were analysed. IFI was documented in 19/125 (15·2%) treatment episodes: five proven (all *Candida*); three probable (two *Candida*, one mixed infection with *Candida* and *Aspergillus*); and 11 possible (all *Aspergillus*). The Platelia® *Aspergillus* ELISA results were analysed at three different positive cut-offs: >1·5 (manufacturer's recommended positive cut-off); >1·0; and >0·5 (Table I).

Using a positive cut-off of >1·5, sequential positive (≥ 2 consecutive) results were recorded in only one of 125 treatment episodes, during which the patient had proven invasive candidiasis (IC) but no EORTC/MSG evidence of IA. In the 12 cases of EORTC/MSG-defined IA (one probable and

Table I. Sequential positive (Seq) (≥ 2 consecutive positive), non-sequential positive (Non-seq) (intermittent or single) and negative (Neg) GM ELISA results at different positive cut-offs (GMI) correlated with EORTC/MSG-defined invasive aspergillosis (IA).

IA/No IA (number of episodes)	GMI >1·5			GMI >1·0			GMI >0·5		
	Seq	Non-seq	Neg	Seq	Non-seq	Neg	Seq	Non-seq	Neg
EORTC/MSG IA (12)*	0	0	12	0	0	12	0	3†	9
No EORTC/MSG IA (113)	1‡	3	109	1‡	9§	103	11‡¶	26\$**	76

*11 possible/1 probable.

†Two cases IA (possible) and one case IA/IC (possible).

‡Case 1, IC (proven).

\$Case 2, IC (probable).

¶Case 3, IC (proven).

**Case 4, IC (proven).

11 possible) no sequential, intermittent or single positive results were recorded.

Despite reducing the positive cut-off to >0.5 , no sequential positive results were recorded in the 12 cases of EORTC/MSG-defined IA (one probable and 11 possible) and non-sequential results were recorded in only three cases of possible IA. In episodes with no EORTC/MSG evidence of IA (113), there were 11 sequential and 26 non-sequential positive results, including four cases of IC. As there was no evidence of EORTC/MSG IA by the time of neutrophil recovery, these positive ELISAs may represent genuine false positive results. False positive results have been associated with dietary GM, antibiotics and cross-reactivity with other fungi and bacteria (McLintock & Jones, 2004). However, it has been suggested that positive results that resolve with neutrophil recovery, or empirical antifungal therapy, may be the result of subclinical infection.

In summary, even with a reduction in the positive cut-off to >0.5 , and the inclusion of single positive results, the sensitivity of the Platelia® *Aspergillus* ELISA was poor when used to detect EORTC/MSG-defined possible or probable IA. This lower positive cut-off resulted in increased numbers of false-positive results, including positive results in patients with IC.

Acknowledgements

Support for this work was provided by grants from the Chief Scientists Office, Scotland, Wyeth Healthcare and Gilead Sciences.

Elaine K. Allan¹
 Niove E. Jordanides¹
 Lorna A. McIntock¹
 Mhairi Copland¹
 Maureen Devaney²
 Karen Stewart¹

Anne N. Parker³
 Peter R.E. Johnson²
 Tessa L. Holyoake¹
 Brian L. Jones⁴

¹Section of Experimental Haematology, Division of Cancer Science and Molecular Pathology, University of Glasgow, Glasgow, ²Departments of Haematology, Lothian University Hospitals NHS Trust, Edinburgh, ³North Glasgow University Hospitals NHS Trust, Glasgow, and ⁴Department of Medical Microbiology, North Glasgow, University Hospitals NHS Trust, Glasgow, UK.
 E-mail: b.l.jones@clinmed.gla.ac.uk

References

- Ascioglu, S., Rex, J.H., de Pauw, B., Bennett, J.E., Bille, J., Crokaert, F., Denning, D.W., Donnelly, J.P., Edwards, J.E., Erjavec, Z., Fiere, D., Lortholary, O., Maertens, J., Meis, J.F., Patterson, T.F., Ritter, J., Selleslag, D., Shah, P.M., Stevens, D.A. & Walsh, T.J. (2002) Defining opportunistic invasive fungal infections in immunocompromised patients with cancer and hematopoietic stem cell transplants: an international consensus. *Clinical Infectious Diseases*, **34**, 7–14.
- Maertens, J., Theunissen, K., Verbeken, E., Lagrou, K., Verhaegen, J., Boogaerts, M. & Eldere, J.V. (2004) Prospective clinical evaluation of lower cut-offs for galactomannan detection in adult neutropenic cancer patients and haematological stem cell transplant recipients. *British Journal of Haematology*, **126**, 852–860.
- McLintock, L.A. & Jones, B.L. (2004) Advances in the molecular and serological diagnosis of invasive fungal infection in haematology patients. *British Journal of Haematology*, **126**, 289–297.
- Prentice, H.G., Kibbler, C.C. & Prentice, A.G. (2000) Towards a targeted, risk-based, antifungal strategy in neutropenic patients. *British Journal of Haematology*, **110**, 273–284.

Keywords: fungal infection, enzyme-linked immunosorbent assay, haematological malignancy, *Candida*, *Aspergillus*.