

Serum p53 Antibody as Tumor Marker for Follow-Up of Colorectal Cancer After Curative Resection

Reiping Tang, MD¹, Chien Yuh Yeh, MD¹, Jeng-Yi Wang, MD¹, Chung Rong Changchien, MD¹, Jinn-Shiun Chen, MD¹, and Ling Ling Hsieh, DrPH²

¹Department of Surgery, Colorectal Section, Chang Gung Memorial Hospital, Tao-Yuan, Taiwan; ²Department of Public Health, Chang Gung University, Tao-Yuan, Taiwan

ABSTRACT

Background. No large-scale studies have examined the use of serial measurements of serum p53 antibodies (s-p53Abs) combined with carcinoembryonic antigen (CEA) measurements during the follow-up of colorectal cancer (CRC) patients after curative resection.

Methods. A highly specific enzyme-linked immunosorbent assay was used to analyze s-p53Abs levels in 305 CRC patients before and after curative resection at a single institution. Agreement between recurrence and serial s-p53Ab and CEA measurements was evaluated by diagnostic accuracy odds ratio (DOR), kappa, and area under receiver operating characteristic curve (AUC).

Results. Among 305 patients, 76 (25%) patients had disease recurrence during follow-up. None of the 168 s-p53Ab seronegative patients (s-p53Ab < 10 U/μL) without recurrence had an abnormal s-p53Ab test during follow-up. Among the remaining low-level (10 U/μL ≤ s-p53Ab ≤ 76 U/μL, *n* = 103) and high-level (s-p53Ab titer > 76 U/μL, *n* = 34) seropositive patients, recurrence defined by s-p53Ab tests resulted in a DOR of 4.3 and ∞, a kappa of 0.35 and 1.00, and an AUC of 0.633 [95% confidence interval (CI), 0.495 to 0.772; *P* = 0.047], and 1.0 (95% CI, 1.000 to 1.000; *P* < 0.0001), respectively. Recurrence defined by CEA tests had an AUC of 0.781 (95% CI, 0.654 to 0.909) for low-level and 0.796 (95% CI, 0.611 to 0.982) for high-level seropositive patients.

Conclusions. Agreement between clinical recurrence and serial s-p53Ab test was dependent upon preoperative s-p53Ab level. Serial s-p53Ab testing outperformed CEA

testing when predicting clinical recurrence in colorectal cancer patients with an abnormal preoperative s-p53Ab level.

Carcinoembryonic antigen (CEA) monitoring has been widely used for more than three decades for gastrointestinal cancers, especially colorectal cancer (CRC).^{1–4} However, the clinical usefulness of this marker has remained contentious owing to some studies arguing that it has limited sensitivity when predicting relapse.^{5,6} In recent years, although the use of various other serum markers for the monitoring of CRC has been described, few have proved to be superior or even comparable to CEA during the follow-up of CRC patients.^{2,3,7}

Circulating anti-p53 antibodies can be found in sera of patients with various human malignancies.^{8–12} Most studies of these antibodies note a strong relationship between p53 protein accumulation and gene mutation. The p53 humoral response is believed to result from self-immunization against p53 protein that has accumulated in tumor cells.¹³ Most studies have focused on measuring serum p53 antibodies (s-p53Abs) as a prognostic marker for a variety of cancers. Some studies have noted a correlation between preoperative s-p53Ab and a poor outcome in cancer patients.^{14–20} However, others have reported a different result.^{21–24} A previous study by this research group, as well as several studies from other groups, found that the adverse effects of a high s-p53Ab level on survival in CRC patients is due to an increased frequency of s-p53Ab positivity in patients with lymph node metastasis and advanced disease.^{25,26}

The value of serial measurements of s-p53Ab as a marker for early recurrence of disease has not been thoroughly elucidated. Some studies have reported that serial measurement of s-p53Ab provides useful clinical information during follow-up among cancer patients.^{27,28}

However, other studies have been inconclusive.²⁹ Other than one study, which reported a promising role for s-p53Ab when monitoring CRC follow-up in a small group of patients, there been no other large-scale surveillance studies of s-p53Abs in CRC patients.³⁰ This study analyzes serial s-p53Ab levels in CRC patients during a long-term follow-up in order to test the usefulness of the serial measurement of s-p53Ab alone or in conjunction with the simultaneous measurement of CEA, when used for the early detection of recurrence in curatively resected CRC patients.

MATERIALS AND METHODS

Patients

This prospective study was approved by the Institutional Review Board of Chang Gung Memorial Hospital (CGMH) and undertaken according to the ethical guidelines of human investigation. Informed consent was obtained from all the patients. Patients fulfilling the following criteria were enrolled in the study: (1) prior curative resection for histology-proven primary adenocarcinoma of the colorectum at CGMH between 1995 and 2002, (2) availability of serial serum samples from before the operation and from after the surgery, and (3) follow-up with a definitive clinical outcome. The exclusion criteria were as follows: (1) synchronous or metachronous extracolonic cancers, (2) having neoadjuvant therapy for rectal cancer, and (3) fewer than three follow-up samples available for s-p53Ab analysis. Serum samples were collected before surgery and at the time of regular outpatient follow-up after surgery and stored at -80°C until use. To increase serial serum sample numbers for the present study, some serum samples were retrieved from the hospital serum bank. All cases were followed up at the outpatient department every 3–6 months until death or until December 2007. All the patients were followed according to the hospital guidelines of care. Briefly, all patients underwent a follow-up protocol of an outpatient visits every 3–6 months. The follow-up included physical examination and carcinoembryonic antigen tests as well as chest X-ray, abdominal sonography or abdominal computer-assisted tomography scan, and colonoscopy every 1–3 years after operation. The primary end-point was clinical recurrence, which was defined as relapse confirmed by histology or by an imaging study. All the clinical data relevant to the patients were obtained from the Colorectal Cancer Registry Database.

Assay for S-p53 Antibodies and CEA

All the samples were analyzed within 1 year after collection without previous thawing. Serum anti-p53 antibodies

(s-p53Abs) were detected by enzyme-linked immunosorbent assay (ELISA) kit (p53-Autoantibody ELISA^{PLUS}; Calbiochem-Novabiochem, Darmstadt, Germany) in accordance with methods described previously.²⁶ Briefly, all samples were diluted 1:100 in reconstituted sample dilution buffer and assayed in duplicate. The control samples contained a negative control, control I (calibrator with defined p53 autoantibody titer), control II (calibrator: sample dilution buffer = 1:2.5), control III (calibrator: sample dilution buffer = 1:5), and control IV (calibrator: sample dilution buffer = 1:10), which were all included on each microtiter plate. If the absorptions of the samples were outside the calibration range, they were tested again at a higher dilution. The s-p53Ab titer was determined based on the calibration curve generated by serial dilution of the calibrator. Samples were designated as s-p53Ab seronegative if the titer was less than 10, which is equivalent to the cutoff value of the assay. According to a previous study, in which the sera from 611 normal control subjects were analyzed, the highest titer value detected for s-p53Abs was 76.²⁶ Thus, for the seropositive samples, a titer value of between 10 and 76 units per microliter (U/ μL) was designated as low and a value above 76 U/ μL was designated as high.²⁶ All samples were analyzed successively with blinding to clinical information by the same well-trained technician. The clinical dataset was decoded to analyze any association with the s-p53Abs only after completion of the assays. Serum CEA levels were determined by Abbott Architect 2000 (Abbott Laboratories, Abbott Park, IL; reference threshold <5 ng/mL) at Clinical Pathology, CGMH.³¹

Statistical Analysis

The differences in clinicopathological parameters between seronegative and seropositive groups were tested using Student's *t* test for continuous variables (age, CEA, and follow-up period) and Fisher's exact or chi-square test for categorical variables. A two-tailed *P* value of ≤ 0.05 was considered significant.

To assess seroconversion in seropositive patients after surgery, the lowest values of the serial follow-up s-p53Abs 1 year after surgery were identified for each patient. Four cutoffs of these values for positive classification were used: 10, 20, 30, and 40 U/ μL . Receiver operating characteristics (ROCs) were drawn for patients stratified by s-p53Ab serology. The power of postoperative seroconversion was evaluated using the area under the curve. The sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), diagnostic accuracy odds ratio (DOR), and kappa were calculated for positive s-p53Ab testing in patients stratified by level of s-p53Ab positivity.

The CEA test was defined as positive if two consecutive postoperative CEA values were greater than 5 ng/mL or the elevated preoperative CEA values had not returned to the normal level (<5 ng/mL) after surgery. Kappa values were calculated using the following equation: $2 \times (ad - bc) / [(a + b) \times (b + d) + (a + c) \times (c + d)]$, where the values of a , b , c , and d represents the number of true-negative, false-negative, false-positive, and true-positive patients, respectively. The SPSS software package (version 13.0; SPSS, Chicago, IL) was used for the statistically analysis.

RESULTS

A total of 305 patients were eligible for this study. There were 45 (14.8%), 130 (42.6%), and 130 (42.6%) cases in tumor-node-metastasis (TNM) stages I, II, and III, respectively. The preoperative titers of s-p53Ab varied widely from 0 to 60,000 U/ μ L. The titers were <10 U/ μ L (seronegative) in 168 patients, 10–76 U/ μ L (defined as low-level seropositivity) in 103 patients, and >76 U/ μ L (defined as high-level seropositivity) in 34 patients. The value of preoperative CEA widely ranged from 0.5 to 544 ng/mL [mean, 15.0; standard deviation (SD), 50.0 ng/mL]. One hundred and one patients had a preoperative value of 5.0 ng/mL or more. The distribution of CEA levels was similar among patients stratified as positive for s-p53Ab (<10 versus ≥ 10 U/ μ L) or by levels of positivity for s-p53Ab (<10 , 10–76, >76 U/ μ L). Table 1 shows characteristics of these patients stratified by presence of s-p53Ab (e.g. <10 versus ≥ 10 U/ μ L). The two groups did not show any significant differences in terms of gender, tumor stage, histological grade, preoperative CEA levels or frequency of clinical recurrence. However, mucinous adenocarcinomas were significantly associated with seronegativity, and seropositivity was associated with rectal cancers. In addition, seropositive patients were older, had shorter duration of follow-up (47 versus 55 months; $P = 0.020$) and more frequently received postoperative adjuvant therapy (50% versus 31%; $P = 0.001$).

Disease Status at End of Follow-Up and Pattern of Recurrence

The mean follow-up was 61.7 months with a range of 2–134 months. At end of follow-up, 76 (25%) patients had recurrent disease. The initial sites of recurrence included: 7 locoregional recurrences, 18 intra-abdominal or retroperitoneal recurrences, 29 hepatic recurrences, 17 pulmonary recurrences, and 9 recurrences in brain or bone. A total of 80 metastases were noted among 76 patients since 4 patients had simultaneous double metastases (2 with hepatic and pulmonary metastases, 1 with hepatic and

TABLE 1 Characteristics of patients with TNM stage I–III colorectal cancer versus status of preoperative s-p53Ab serology^a

| Characteristics | Seronegative patients ($n = 168$) | Seropositive patients ($n = 137$) | P value |
|-----------------------------------|--|--|-----------|
| Gender (%) | | | 0.198 |
| Female | 42 | 50 | |
| Male | 58 | 50 | |
| Age at diagnosis, years | | | 0.008 |
| Mean | 59 | 63 | |
| Range | 20–90 | 33–89 | |
| Tumor location (%) | | | 0.001 |
| Colon | 58 | 37 | |
| Rectum | 40 | 61 | |
| Colon and rectum | 2 | 2 | |
| Tumor TNM stage (%) | | | 0.528 |
| I | 16 | 13 | |
| II | 44 | 41 | |
| III | 40 | 46 | |
| Tumor morphology (%) | | | 0.469 |
| Polypoid | 17 | 23 | |
| Ulcerative | 77 | 71 | |
| Other type | 6 | 6 | |
| Histological grade (%) | | | 0.231 |
| Good differentiation | 20 | 15 | |
| Moderate differentiation | 74 | 82 | |
| Poor differentiation | 6 | 4 | |
| Histological type (%) | | | 0.001 |
| Adenocarcinoma | 91 | 99 | |
| Mucinous adenocarcinoma | 9 | 1 | |
| Preoperative CEA, ng/mL | | | 0.064 |
| Mean | 19.9 | 9.2 | |
| Range | 0.5–544.0 | 0.5–138.0 | |
| Preoperative CEA (%) | | | 0.690 |
| ≤ 5 ng/mL | 68 | 66 | |
| >5 ng/mL | 32 | 34 | |
| Adjuvant therapy ^b (%) | | | 0.001 |
| None | 69 | 50 | |
| Yes | 31 | 50 | |
| Duration of follow-up, months | | | 0.020 |
| Mean | 55 | 47 | |
| Range | 3–113 | 2–116 | |
| Relapse during follow-up (%) | | | 0.762 |
| Absence | 74 | 76 | |
| Presence | 26 | 24 | |

^a Seronegative when the value of preoperative s-p53Ab was less than 10 U/ μ L; seropositive when s-p53Ab ≥ 10 U/ μ L

^b 5-Fluorouracil-based chemotherapy for colon cancer and postoperative combined chemoradiation therapy for rectal cancer

retroperitoneal metastases, and 1 with bone and intra-abdominal metastases). The pattern of recurrence was not different between seronegative and seropositive group (data not shown). Briefly, liver was the most common site of disease failure among both groups (seronegative 34.9%, seropositive 33.3%). Intra-abdominal or retroperitoneal metastases were more frequent in the seronegative group (seronegative 27.9%, seropositive 12.1%) whereas lung (seronegative 16.3%, seropositive 24.2%) and bone/brain metastases (seronegative 7.0%, seropositive 15.2%) were seen more often in the seropositive group.

Determination of Cutoff Value of Postoperative Serial s-p53Ab

For s-p53Ab seronegative patients, the relationship between recurrence and cutoff value of postoperative serial s-p53Ab was not significant at any level; the areas under the ROC curve (AUC) were 0.508, 0.512, 0.500, and 0.500 for cutoff value of 10, 20, 30, and 40 U/ μ L, respectively (Fig. 1a). Among these 168 seronegative patients, 160 cases (95.2 %) remained seronegative and another 8 cases had at least a follow-up titer ranging from 10–23 U/ μ L, with a mean of 15.6 U/ μ L. For the low-level seropositive patients, the AUC values were 0.620 (95% CI, 0.488 to 0.752; $P = 0.73$), 0.633 (95% CI, 0.496 to 0.771, $P = 0.047$), 0.633 (95% CI, 0.495 to 0.772; $P = 0.047$), and 0.593 (95% CI, 0.455 to 0.732; $P = 0.164$) for cutoff value of 10, 20, 30, and 40 U/ μ L, respectively (Fig. 1b). For the high-level seropositive patients, the AUC values were 0.891 (95% CI, 0.779 to 1.004; $P < 0.001$), 0.978 (95% CI, 0.928 to 1.029; $P < 0.0001$), 1.0 (95% CI, 1.000 to 1.000; $P < 0.0001$), and 1.0 (95% CI, 1.000 to 1.000; $P < 0.0001$) for cutoff value of 10, 20, 30, and 40 U/ μ L, respectively (Fig. 1c). Taken together, the cutoff score of 30 U/ μ L performed best with regard to AUC among low-level and high-level seropositive patients combined. In addition, Table 2 shows that the cutoff value at 30 U/ μ L outperforms the other three cutoff values in terms of DOR, kappa, and AUC. Thus, an s-p53Ab of 30 U/ μ L was chosen as the cutoff value for the later analyses.

Seroconversion and Reconversion Among s-p53Abs-Positive Patients

Seroconversion was defined as postoperative serial s-p53Ab titers that were decreasing and with a lowest level less than 30 U/ μ L. Among 137 seropositive patients, 126 (92%) had complete seroconversion and 11 (8%) did not undergo seroconversion. The recurrence rates for those with seroconversion and for those without were 17% (22/126) and 100% (11/11), respectively ($P < 0.0001$). Among 126 patients with seroconversion, 14 (11%) reconverted to




FIG. 1 Receiver operating characteristic (ROC) curves for patients with **a** negative, **b** low-level, and **c** high-level preoperative s-p53Ab drawn with four cutoff values for the serial s-p53Ab levels (10 U/ μ L, blue line; 20 U/ μ L, green line; 30 U/ μ L, red line, and 40 U/ μ L, purple line)

seropositive (with a titer >30 U/ μ L). Compared with 12 out of 112 (11%) patients without reconversion, 10 out of these 14 (71%) patients with reconversion had a recurrence

TABLE 2 Results for serial s-p53Ab cutoff values for recurrence in patients with preoperative high-level and low-level s-p53Ab

| Preoperative level and postoperative cutoff value (U/μL) | Sensitivity (%) | Specificity (%) | PPV (%) | NPV (%) | DOR | Kappa | Area under ROC | |
|--|-----------------|-----------------|---------|---------|------|-------|----------------|-------------|
| | | | | | | | Area | 95% CI |
| High-level s-p53Ab (>76 U/μL) | | | | | | | | |
| ≥10 | 100.0 | 78.3 | 64.3 | 100.0 | 5.4 | 0.67 | 0.891 | 0.779–1.004 |
| ≥20 | 100.0 | 95.7 | 90.0 | 100.0 | 31.0 | 0.93 | 0.978 | 0.928–1.029 |
| ≥30 | 100.0 | 100 | 100.0 | 100.0 | ∞ | 1.00 | 1.000 | 1.000–1.000 |
| ≥40 | 100.0 | 100 | 100.0 | 100.0 | ∞ | 1.00 | 1.000 | 1.000–1.000 |
| Low-level s-p53Ab (10–76 U/μL) | | | | | | | | |
| ≥10 | 48.0 | 76.0 | 40.0 | 81.4 | 2.2 | 0.23 | 0.620 | 0.488–0.752 |
| ≥20 | 32.0 | 94.7 | 66.7 | 80.7 | 3.8 | 0.32 | 0.633 | 0.496–0.771 |
| ≥30 | 28.0 | 98.7 | 87.5 | 80.4 | 4.3 | 0.35 | 0.633 | 0.495–0.772 |
| ≥40 | 20.0 | 98.7 | 83.3 | 78.7 | 3.8 | 0.25 | 0.593 | 0.455–0.732 |

ROC receiver operating characteristics, PPV positive predictive value, NPV negative predictive value, DOR diagnostic accuracy odds ratio

($P < 0.0001$). Accordingly, patients were defined as having positive postoperative s-p53Ab tests if (1) they did not have seroconversion after surgery or (2) they had a seroconversion but with a later reconversion back to the initial state.

Agreement Between Clinical Recurrence and Serial s-p53Ab and CEA Testing

The overall agreement between clinical recurrence and serial CEA measurements during follow-up was fair, with a kappa of 0.70 (sensitivity, 74%; specificity, 94%; positive predictive value, 80%; negative predictive value, 92%). When stratified by preoperative CEA level, the sensitivity of the serial CEA test was 67% in patients with normal preoperative CEA levels and 82% in those with abnormal preoperative CEA levels.

Figure 2 shows the flow of patients from preoperative s-p53Ab level to clinical recurrence during follow-up and the testing results (serial s-p53Ab and CEA measurements) for each corresponding subgroup. A total of 76 patients had clinical recurrence. Time to recurrence ranged from 2 to 58 months (mean, 20.4 months). Positive s-p53Ab tests were observed in 21 patients prior to development of clinical recurrence (lag time ranging from 0.1 to 23.7 months with a mean of 6.8 months and an SD of 6.0 months). The remaining two patients had an abnormal test at the time of clinical recurrence. The pattern of recurrence was not related to their status in terms of serial s-p53Ab testing (data not shown).

Table 3 summarizes the diagnostic efficacy of the serial measurements of s-p53Ab, CEA, and the combination of both for monitoring CRC stratified by preoperative s-p53Ab levels. Firstly, these values showed a strong association between the diagnostic efficacy of serial s-p53Ab testing with the preoperative level of s-p53Ab.


The CEA test performed better than the s-p53Ab test (AUC, 0.879 versus 0.500; κ , 0.76 versus 0) in the seronegative group. Furthermore, it did as well as the s-p53Ab test in the low-level group, but worse than the s-p53Ab test (AUC, 0.796 versus 0.978; κ , 0.64 versus 0.93) in the high-level seropositive group. Secondly, the performance of serial CEA measurements was not related to the preoperative level of s-p53Ab. Thirdly, the results also show that adding s-p53Ab to CEA testing was not better than CEA alone when detecting recurrent disease among the s-p53Ab seronegative patients. Likewise, adding CEA to s-p53Ab testing did not improve the sensitivity, kappa or AUC of serial s-p53Ab testing alone among the s-p53Ab seropositive patients.

DISCUSSION

The analytical results indicated that serum anti-p53 antibody (s-p53Ab) levels may be a reliable tumor marker for recurrence during follow-up for s-p53Ab seropositive patients with curatively resected CRC. This may be particularly true for the group of patients with a high preoperative serum anti-p53 antibody level (>76 U/ μ L), where s-p53Ab monitoring outperforms CEA in terms of sensitivity, specificity, and diagnostic accuracy for detecting disease recurrence among CRC patients.

Of the numerous tumor markers suggested for CRC, CEA is the surveillance marker of choice and has been recommended for monitoring purposes on many occasions.^{1–3,31,32} However, as noted in several studies, its clinical use is limited since the sensitivity and specificity for serial CEA measurements when detecting recurrent disease are limited.^{5,6} In the current study, the sensitivity of CEA monitoring was low at 64% in the seropositive patients, compared with 81% in the seronegative patients. These analytical results also confirmed that increased CEA

FIG. 2 Consolidated chart of 305 patients with serial s-p53Ab and CEA testing results stratified by preoperative s-p53Ab level and clinical recurrence during follow-up. Patients were defined as having positive postoperative s-p53Ab tests if (1) they did not have seroconversion after surgery or (2) they had a seroconversion but with a later reconversion of s-p53Ab serology. The CEA test was defined as positive if two consecutive postoperative CEA values were greater than 5 ng/mL or the elevated preoperative CEA values did not returned to the normal level (<5 ng/mL) after surgery



in association with disease recurrence is not uncommon (67% in the current series) among patients with a normal preoperative CEA level.³³ Thus, CEA tests are needed for postoperative follow-up in all patients, both with and without an elevated CEA value.

The role that s-p53Abs could play in the follow-up of CRC patients after curative resection has not been adequately elucidated. Hammel et al., in a study of eight patients over 12–24 months of follow-up, reported that s-p53Ab monitoring might help physicians to follow up CRC patients.³⁰ Another small series that included 27 patients with early CRC treated by endoscopic resection suggested that p53 antibody seroconversion might indicate local curability after endoscopic treatment.³⁴ In the current study we observed that the sensitivity of s-p53Ab monitoring was highly correlated with s-p53Ab level before resection, being 0% for seronegative patients, 55% for low-level seropositive patients, and 100% for high-level seropositive patients. Likewise, the diagnostic efficacies were highly correlated with the preoperative s-p53Ab level with regard to DOR, κ , and AUC (Table 3). Abnormal s-p53Ab titers were noted only in the seropositive patients but not in seronegative patients. Therefore, postoperative serial s-p53Ab tests are needed only in patients having an elevated s-p53Ab titer before surgery.

This leaves open the question as to why the sensitivity of the s-p53Ab test is highly correlated with preoperative s-p53Ab level. It is possible that the anti-p53 antibodies are of the immunoglobulin G (IgG) isotype, which is dependent on helper T cells and therefore the majority of such patients display long-lasting anti-p53 T helper (Th)

immunity even in the absence of measurable s-p53Ab. This might easily last for several years after resection of the primary colorectal cancer.³⁵ Thus, the capacity of a patient to produce circulating anti-p53 antibodies in response to p53 protein from recurrent tumor cells will depend on whether the patient has been pre-immunized against the p53 protein. However, whether CRCs harboring p53 alterations (p53 mutations or accumulated proteins) are more immunogenic than wild-type p53 also requires further investigation.^{9,36,37} Other possible explanations include individual differences in major histocompatibility complex-based presentation of the p53 protein, the position of the p53 mutations, the presence of multiple steps in the carcinogenesis process, and a tissue-specific humoral response against the p53 antibody.^{9,26,36,38}

Alteration of the *P53* gene, the most frequent genetic alteration in human cancer, causes an accumulation of mutant p53 in the nucleus of tumor cells. It is noteworthy in the context of this paper that p53 antibodies have been found in the sera of patients with a variety of different forms of cancer as well as in those with inflammatory bowel disease.^{38,39} Although the frequency of abnormal s-p53Ab levels varies among different cancers, Lubin et al. demonstrated that the immune response of patients with p53 antibodies is restricted to a small subset of peptides regardless of the form of cancer.¹³ Thus, examination of s-p53Ab in addition to other specific tumor markers might also serve as a useful indicator of tumor recurrence in other malignancies that express p53 alterations.

In conclusion, serial s-p53Ab testing outperformed CEA testing in predicting clinical recurrence in colorectal cancer

TABLE 3 Statistics of serial measurements of s-p53Ab alone, CEA alone, and the combination of both among patients with three levels of preoperative s-p53Ab

| Level of preoperative s-p53Ab and serum markers | Sensitivity (%) | Specificity (%) | PPV (%) | NPV (%) | DOR | κ | Area under ROC | |
|---|-----------------|-----------------|---------|---------|------|----------|----------------|-------------|
| | | | | | | | Area | (95%CI) |
| Negative (<10 U/ μ L) | | | | | | | | |
| CEA | 81.4 | 94.4 | 83.3 | 93.7 | 10.2 | 0.76 | 0.879 | 0.807–0.951 |
| s-p53Ab | 0.0 | 100.0 | – | 74.4 | 2.9 | 0.00 | 0.500 | 0.400–0.600 |
| CEA + s-p53Ab | 81.4 | 94.4 | 83.3 | 93.7 | 10.2 | 0.76 | 0.879 | 0.807–0.951 |
| Low-level (10–76 U/ μ L) | | | | | | | | |
| CEA | 63.6 | 92.6 | 70.0 | 90.4 | 6.4 | 0.58 | 0.781 | 0.654–0.909 |
| s-p53Ab | 54.5 | 95.1 | 75.0 | 88.5 | 6.4 | 0.55 | 0.748 | 0.613–0.883 |
| CEA + s-p53Ab | 81.8 | 87.7 | 64.3 | 94.7 | 6.4 | 0.63 | 0.847 | 0.745–0.950 |
| High-level (>76 U/ μ L) | | | | | | | | |
| CEA | 63.6 | 95.7 | 87.5 | 84.6 | 5.8 | 0.64 | 0.796 | 0.611–0.982 |
| s-p53Ab | 100.0 | 95.7 | 91.7 | 100 | 33.0 | 0.93 | 0.978 | 0.928–1.028 |
| CEA + s-p53Ab | 100.0 | 91.3 | 84.6 | 100 | 16.0 | 0.87 | 0.957 | 0.886–1.027 |

CEA carcinoembryonic antigen, PPV positive predictive value, NPV negative predictive value, DOR diagnostic accuracy odds ratio, κ kappa. Dash indicates denominator = 0

patients with an abnormal preoperative s-p53Ab level. Such monitoring would seem to be a valuable and cost-effective component of the postoperative follow-up program for those with elevated preoperative s-p53Ab levels. This should take the form of postoperative CEA monitoring for s-p53Ab seronegative patients, with combined CEA and s-p53Ab tests for low-level seropositive patients and s-p53Ab monitoring for high-level seropositive patients. These findings require confirmation by further independent studies before the tumor marker (s-p53Ab) can be recommended for routine clinical use.

ACKNOWLEDGMENT We thank Miao-Chen Ko for assistance with the assay for s-p53 antibodies and Professor James Tsai-Yuan Wu for his valuable discussions. This study was supported by grant NSC90-2314-B-182A-11 from the National Science Council (R. Tang) and grants DOH85-HR-516, DOH86-HR-516 and DOH87-HR-516 from the National Health Research Institute, Department of Health, Executive Yuan, ROC (L.L. Hsieh).

REFERENCES

1. Chau I, Allen MJ, Cunningham D, et al. The value of routine serum carcino-embryonic antigen measurement and computed tomography in the surveillance of patients after adjuvant chemotherapy for colorectal cancer. *J Clin Oncol*. 2004;22:1420–9.
2. Locker GY, Hamilton S, Harris J, et al. ASCO 2006 update of recommendations for the use of tumor markers in gastrointestinal cancer. *J Clin Oncol*. 2006;24:33:5313–27.
3. Duffy MJ, van Dalen A, Haglund C, et al. Tumour markers in colorectal cancer: European Group on Tumour Markers (EGTM) guidelines for clinical use. *Eur J Cancer*. 2007;43:1348–60.
4. Borie F, Daures JP, Millat B, Tretarre B. Cost and effectiveness of follow-up examinations in patients with colorectal cancer resected for cure in a French population-based study. *J Gastrointest Surg*. 2004;8:552–8.
5. Korner H, Soreide K, Stokkeland PJ, Soreide JA. Diagnostic accuracy of serum-carcinoembryonic antigen in recurrent colorectal cancer: a receiver operating characteristic curve analysis. *Ann Surg Oncol*. 2007;14:417–23.
6. Hanke B, Riedel C, Lampert S, et al. CEA and CA 19-9 measurement as a monitoring parameter in metastatic colorectal cancer (CRC) under palliative first-line chemotherapy with weekly 24-hour infusion of high-dose 5-fluorouracil and folinic acid. *Ann Oncol*. 2001;12:221–6.
7. Wang JY, Lin SR, Wu DC, et al. Multiple molecular markers as predictors of colorectal cancer in patients with normal perioperative serum carcinoembryonic antigen levels. *Clin Cancer Res*. 2007;13:2406–13.
8. von Brevern MC, Hollstein MC, Cawley HM, et al. Circulating anti-p53 antibodies in esophageal cancer patients are found predominantly in individuals with p53 core domain mutations in their tumors. *Cancer Res*. 1996;56:4917–21.
9. Wu CM, Tang R, Chen JR, et al. Humoral response to p53 is associated with conserved domains II and IV mutations in human colorectal cancer: a case-control study in Taiwan. *Oncol Rep*. 2004;12:1045–51.
10. Shibata Y, Kotanagi H, Andoh H, Koyama K, Itoh H, Kudo S. Detection of circulating anti-p53 antibodies in patients with colorectal carcinoma and the antibody's relation to clinical factors. *Dis Colon Rectum*. 1996;39:1269–74.
11. Wunderlich H, Hindermann W, Huller M, et al. The correlation of p53 protein overexpression and p53 antibodies in serum of patients with transitional cell carcinoma of the urinary bladder. *Urol Int*. 2000;64:13–7.
12. Wild CP, Ridanpaa M, Anttila S, et al. p53 antibodies in the sera of lung cancer patients: comparison with p53 mutation in the tumour tissue. *Int J Cancer*. 1995;64:176–81.
13. Lubin R, Schlichtholz B, Bengoufa D, et al. Analysis of p53 antibodies in patients with various cancers define B-cell epitopes of human p53: distribution on primary structure and exposure on protein surface. *Cancer Res*. 1993;53:5872–6.
14. Peyrat JP, Bonnetterre J, Lubin R, Vanlemmens L, Fournier J, Soussi T. Prognostic significance of circulating P53 antibodies in patients undergoing surgery for locoregional breast cancer. *Lancet*. 1995;345:621–2.

15. Zalcmán G, Tredaniel J, Schlichtholz B, et al. Prognostic significance of serum p53 antibodies in patients with limited-stage small cell lung cancer. *Int J Cancer*. 2000;89:81–6.
16. Murray PV, Soussi T, O'Brien ME, et al. Serum p53 antibodies: predictors of survival in small-cell lung cancer?. *Br J Cancer*. 2000;83:1418–24.
17. Chow V, Yuen AP, Lam KY, Ho WK, Wei WI. Prognostic significance of serum p53 protein and p53 antibody in patients with surgical treatment for head and neck squamous cell carcinoma. *Head Neck*. 2001;23:286–91.
18. Kressner U, Glimelius B, Bergström R, Pahlman L, Larsson A, Lindmark G. Increased serum p53 antibody levels indicate poor prognosis in patients with colorectal cancer. *Br J Cancer*. 1998;77:1848–51.
19. Sainger RN, Shah MH, Desai AA, et al. Clinical significance of serum p53 antibodies in oral cancer. *Tumori*. 2006;92:134–9.
20. Cai HY, Wang XH, Tian Y, Gao LY, Zhang LJ, Zhang ZY. Changes of serum p53 antibodies and clinical significance of radiotherapy for esophageal squamous cell carcinoma. *World J Gastroenterol*. 2008;14:4082–6.
21. Rosenfeld MR, Malats N, Schramm L, et al. Serum anti-p53 antibodies and prognosis of patients with small-cell lung cancer. *J Natl Cancer Inst*. 1997;89:381–5.
22. Morita T, Tachikawa N, Kumamaru T, et al. Serum anti-p53 antibodies and p53 protein status in the sera and tumors from bladder cancer patients. *Eur Urol*. 2000;37:79–84.
23. Chang SC, Lin JK, Lin TC, Liang WY. Genetic alteration of p53, but not overexpression of intratumoral p53 protein, or serum p53 antibody is a prognostic factor in sporadic colorectal adenocarcinoma. *Int J Oncol*. 2005;26:65–75.
24. Soussi T. p53 Antibodies in the sera of patients with various types of cancer: a review. *Cancer Res*. 2000;60:1777–88.
25. Nozoe T, Yasuda M, Honda M, Inutsuka S, Korenaga D. Clinicopathologic significance in serum presence of anti-p53 antibody in patients with colorectal carcinoma. *Hepatogastroenterology*. 2007;54:1422–5.
26. Tang R, Ko MC, Wang JY, et al. Humoral response to p53 in human colorectal tumors: a prospective study of 1,209 patients. *Int J Cancer*. 2001;94:859–63.
27. Shimada H, Takeda A, Arima M, et al. Serum p53 antibody is a useful tumor marker in superficial esophageal squamous cell carcinoma. *Cancer*. 2000;89:1677–83.
28. Gumus E, Demirel G, Tanriverdi O, Horasanli K, Ozmen G, Miroglu C. Importance of serum p53 antibodies during follow-up after treatment of invasive bladder tumors. *Urol Int*. 2004;72:292–8.
29. Gadducci A, Ferdeghini M, Buttitta F, et al. Serum anti-p53 antibodies in the follow-up of patients with advanced ovarian carcinoma. *Anticancer Res*. 1998;18:3763–5.
30. Hammel P, Boissier B, Chaumette MT, et al. Detection and monitoring of serum p53 antibodies in patients with colorectal cancer. *Gut*. 1997;40:356–61.
31. Wang JY, Tang R, Chiang JM. Value of carcinoembryonic antigen in the management of colorectal cancer. *Dis Colon Rectum*. 1994;37:272–7.
32. Park JY, Lee KH. Carcinoembryonic antigen and patterns of recurrence after curative resection of the colorectal cancer. *Hepatogastroenterology*. 2007;54:1966–9.
33. Grossmann I, de Bock GH, Meershoek-Klein Kranenbarg WM, van de Velde CJ, Wiggers T. Carcinoembryonic antigen (CEA) measurement during follow-up for rectal carcinoma is useful even if normal levels exist before surgery. A retrospective study of CEA values in the TME trial. *Eur J Surg Oncol*. 2007;33:183–7.
34. Takeda A, Shimada H, Nakajima K, et al. Serum p53 antibody as a useful marker for monitoring of treatment of superficial colorectal adenocarcinoma after endoscopic resection. *Int J Clin Oncol*. 2001;6:45–9.
35. van der Burg SH, de Cock K, Menon AG, et al. Long lasting p53-specific T cell memory responses in the absence of anti-p53 antibodies in patients with resected primary colorectal cancer. *Eur J Immunol*. 2001;31:146–55.
36. Forslund A, Kressner U, Lindmark G, Inganas M, Lundholm K. Serum anti-p53 in relation to mutations across the entire translated p53 gene in colorectal carcinomas. *Int J Oncol*. 2001;19:501–6.
37. Lubin R, Zalcmán G, Bouchet L, et al. Serum p53 antibodies as early markers of lung cancer. *Nat Med*. 1995;1:701–2.
38. Lubin R, Schlichtholz B, Teillaud JL, Garay E, Bussel A, Wild CP. p53 antibodies in patients with various types of cancer: assay, identification, and characterization. *Clin Cancer Res*. 1995;1:1463–9.
39. Yoshizawa S, Matsuoka K, Inoue N, et al. Clinical significance of serum p53 antibodies in patients with ulcerative colitis and its carcinogenesis. *Inflamm Bowel Dis*. 2007;13:865–73.