

The use of a vascularised periosteal patch onlay graft in the management of nonunion of the proximal scaphoid

N. W. Thompson,
A. Kapoor,
J. Thomas,
M. J. Hayton

*From Wrightington
Hospital, Wigan,
England*

We describe the use of a vascularised periosteal patch onlay graft based on the 1,2 intercompartmental suprarotational artery in the management of 11 patients (ten men, one woman) with chronic nonunion involving the proximal third of the scaphoid. The mean age of the patients was 31 years (21 to 45) with the dominant hand affected in eight. Six of the patients were smokers and three had undergone previous surgery to the scaphoid. All of the proximal fragments were avascular. The presence of union was assessed using longitudinal axis CT.

Only three patients progressed to union of the scaphoid and four required a salvage operation for a symptomatic nonunion. The remaining four patients with a persistent nonunion are asymptomatic with low pain scores, good grip strength and a functional range of wrist movement.

Although this technique has potential technical advantages over vascularised pedicled bone grafting, the rate of union has been disappointing and we do not recommend it as a method of treatment.

The combined presence of nonunion of the proximal third of the scaphoid and avascular necrosis (AVN) adversely affects the chances of achieving union by fixation and conventional bone grafting.¹⁻⁵ Vascularised bone grafting has thus evolved as an option, with reported rates of union ranging between 27% and 100%.⁶⁻¹¹

Despite the potential advantages of this technique in achieving union, technical problems include difficulty in seating the graft, particularly when the proximal fragment is small, difficulty in achieving stable fixation, extrusion and fragmentation of the graft and dorsoradial impingement by the graft.^{6,12}

Many studies have shown that the periosteum regenerates both cartilage and bone from its progenitor cells.¹³⁻¹⁷ The osteogenic potential of free periosteum may be controversial,^{18,19} but the vascularised periosteal flap has been shown to have an excellent osteogenic capacity.²⁰⁻²³

We describe a surgical technique for the management of chronic nonunion of the proximal third of the scaphoid using a vascularised periosteal onlay patch graft based on the intercompartmental suprarotational artery.^{1,2} We present our results using this technique in a small series of patients.

Patients and Methods

We undertook bone grafting with the application of a vascularised periosteal onlay patch based on the 1,2 intercompartmental suprarotational artery in 12 patients with chronic nonunion of the proximal third of the scaphoid. All the operations were performed by or under supervision of the senior author (MJH). Approval of the Local Ethical Committee was obtained before commencement of the study.

One patient was lost to follow-up and was excluded leaving 11 patients (ten men and one woman) in the final analysis. Their mean age was 31 years (21 to 45) with the dominant hand being involved in eight. The mean duration of nonunion at the time of surgery was 24 months (10 to 55). Three had previous fixation of the scaphoid and non-vascularised bone grafting. One patient had early radioscaphoid arthritis on the pre-operative radiographs. Six patients were smokers.

Operative technique. Under tourniquet control, a dorsal longitudinal incision was made just ulnar to Lister's tubercle to expose the extensor retinaculum and the 1,2 intercompartmental suprarotational artery (Fig. 1). A periosteal patch was raised on a pedicle of sufficient length to reach the dorsal aspect of the nonunion at the proximal pole of the scaphoid (Fig. 2). The extensor retinaculum was reflected in an ulnar

■ N. W. Thompson, MPhil, FRCS(Trauma & Orth), Upper Limb Fellow
■ A. Kapoor, MRCS, Clinical Research Fellow
■ J. Thomas, FRACS, Upper Limb Fellow
■ M. J. Hayton, FRCS(Trauma & Orth), Consultant Hand Surgeon
Centre for Hand and Upper Limb Surgery
Wrightington Hospital, Hall Lane, Appley Bridge, Wigan WN6 9EP, UK.

Correspondence should be sent to Mr M. J. Hayton; e-mail: mikehayton@gmail.com

©2008 British Editorial Society of Bone and Joint Surgery
doi:10.1302/0301-620X.90B12.200808 \$2.00

J Bone Joint Surg [Br]
2008;90-B:1597-601.
Received 28 January 2008;
Accepted after revision 16 July 2008




Fig. 1

Photograph of the vascularised periosteal patch based on the 1,2 inter-compartmental supraretinacular artery (boundary of patch outlined with a marker pen; artery indicated by arrow).



Fig. 2

Photograph of the harvesting of the vascularised periosteal patch on its pedicle (arrow).

direction and the extensor tendons retracted to expose the dorsal capsule of the wrist.

A distally-based capsular flap was raised to expose the non-union which was carefully debrided and any metalwork removed. The proximal fragment was drilled using a fine Kirschner wire to determine the presence of punctate bleeding as a guide to its vascularity.^{6,24} Corticocancellous bone graft harvested from the iliac crest was packed into the defect to correct the length of the scaphoid and alignment followed by the insertion of a cannulated compression screw device (Kompressor compression screw system; Kinetikos Medical Inc., Carlsbad, California) under fluoroscopic control to stabilise the graft and to achieve compression (Fig. 3). The periosteal



Fig. 3

Fluoroscopic image showing the compression screw *in situ*.

patch was then secured over the dorsal aspect of the grafted nonunion using either transosseous sutures or mini-bone anchors (DePuy Mitek, Bracknell, United Kingdom). A below-elbow cast was applied for a minimum of six weeks.

Union was defined on plain radiographs according to the criteria described by Dias.²⁴ All patients had longitudinal axis CT of the scaphoid²⁵ approximately six months after surgery, which was evaluated by an experienced musculoskeletal radiologist (DT).

Patients who did not require subsequent surgery were recalled for clinical review at a mean of 19.4 months (13 to 27). This included the assessment of pain with palpation as evidence of clinical union, the assessment of pain during rest and on activity using a ten-point visual analogue score (VAS) in which zero indicated no pain, and ten the worst possible pain, and measurement of the range of movement and grip strength. Range of movement was measured using a hand-held goniometer and grip strength assessed by a Jamar Dynamometer (JA Preston, Jackson, Michigan).

Results

At operation the proximal pole of the scaphoid was deemed to be avascular in all patients with complete absence of punctate bleeding. Of the 11 patients, four required salvage operations for persistent, symptomatic nonunion by scaphoidectomy and four-corner fusion in three patients and proximal row carpectomy in the fourth. From the plain radiographs of all 11 patients, six were deemed to have



Fig. 4a



Fig. 4b

Radiographs a) six weeks after vascularised periosteal patch grafting and b) six months after operation showing union of the scaphoid.

united, but only three had definite evidence of bony union confirmed by CT.

Of the group of seven patients who did not require further surgery only three had evidence of bony union on CT. All were free from pain except for one with a pain score of 3 of 10 on heavy manual activity. In these seven patients, the mean grip strength was only 80% (75% to 85%) of the contralateral side and the mean wrist flexion and extension were 60° (50° to 70°) and 40° (30° to 50°) respectively. The mean ulnar deviation was 20° (10° to 30°) and the mean radial deviation 15° (10° to 20°).

Examination of the four patients with persistent nonunion but who have not required further surgery revealed pain scores ranging from 0 to 1 at rest, to 1 to 5 on activity. The mean grip strength was 89% (76% to 100%) of the contralateral side and the mean wrist flexion and extension 42° (30° to 55°) and 45° (35° to 60°) respectively. The mean ulnar deviation was 20° (10° to 30°) and the mean radial deviation 15° (10° to 20°).

One patient required evacuation of a haematoma from the donor site of the iliac-crest bone graft 48 hours after surgery. No other complications were encountered.

Discussion

Periosteum is a specialised connective tissue which forms a thin but tough fibrous membrane firmly anchored to the underlying bone. It is a well-vascularised tissue consisting of two discrete layers, an outer fibrous and an inner cambium

layer. The latter serves as a reservoir of undifferentiated pluripotent mesenchymal cells able to differentiate into chondrogenic and osteoblastic lineages and as a source of growth factors involved in bone healing and remodelling.²⁶⁻³¹ In adults, periosteum is inactive, but its bone-forming potential can be reactivated by stimuli such as trauma or infection.

Animal studies have demonstrated the osteogenic potential of periosteum elevated as a vascularised flap.³²⁻³⁴ In the clinical setting, vascularised periosteal grafts have been used in the treatment of fractures of the femoral neck,²³ AVN of the femoral head³⁵ and in head and neck surgery.^{36,37} Thus, it would seem that periosteum provides the potential for the reconstruction of small skeletal defects or defects with poorly vascularised tissue. The vascularised periosteal onlay patch has several technical advantages over the pedicled vascularised bone graft. First, it avoids the necessity of harvesting a block of bone on the vascular pedicle which may be difficult to bed particularly in the presence of a small proximal fragment. The graft may protrude from the dorsoradial border of the scaphoid and interfere with the function of the radiocarpal joint and the bone block may split or fragment during fixation and become devascularised due to disruption of its intra-osseous blood supply. Extrusion of the graft may also occur.¹² Secondly, it eliminates the need for accurate sizing of the bone block before raising the pedicle. With the patch technique, the scaphoid deformity can be corrected using non-vascularised bone graft with the vascularised periosteum providing a source of osteogenic cells and growth factors.

Thirdly, the vascularised pedicle may obscure the view of the graft within the nonunion bed and it can be difficult to ensure that there is good bone contact between the graft and the fragments of the fracture. In the periosteal patch method the defect can be packed under direct vision before application of the periosteal flap.

Despite these technical advantages only three of 11 scaphoid fractures in our series united (Fig. 4). This may have been due to a number of factors. First, the mean duration of nonunion was 24 months. Secondly, all the nonunions involved the proximal third of the scaphoid with avascularity of the proximal fragment in all cases. Thirdly, six patients were smokers. Chang et al¹² reported proximal pole AVN and tobacco use as univariate risk factors for failure of pedicled vascularised bone grafts in their series. Fourthly, we used the more revealing technique of longitudinal axis CT to confirm union rather than relying on the less sensitive method of plain radiography alone. Whereas six patients showed evidence of union on plain films only three did so on CT. Finally, since cell density and hence the osteogenic potential of periosteum may vary between different donor sites³⁸ and the osteogenic potential of periosteum may be reduced with ageing,³⁹ it may be that the periosteal flap raised on the 1,2 intercompartmental supraretrinacular artery in adult patients does not provide any significant osteogenic input.

We accept that in the three patients who proceeded to union we cannot be certain whether the presence of the screw fixation and non-vascularised grafting resulted in union as opposed to the vascularised periosteal patch. Because of the small size of the study we did not attempt to draw any statistical comparisons between those who achieved union and those who did not.

Finally, it is noteworthy that persistent nonunion of the scaphoid after surgery was relatively asymptomatic and provided satisfactory function as indicated by the pain score, grip strength and range of movement in four of our 11 patients. Partial denervation of the wrist and stabilisation of the fragments even in the absence of bony union may account for these findings. However, we acknowledge that, in the longer term, the screw may loosen and that this group of patients may become symptomatic.

In summary, the achievement of healing in chronic nonunion involving the proximal scaphoid remains a challenge. We found that the rate of union obtained by this method was disappointing. Therefore we do not recommend this technique for the treatment of nonunion of the proximal scaphoid.

We would like to thank Dr D. Timperley, Consultant Radiologist, Wrightington Hospital, for his help with the study.

No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article.

References

1. Herbert TJ, Fisher WE. Management of the fractured scaphoid using a new bone screw. *J Bone Joint Surg [Br]* 1984;66-B:114-23.
2. Green DP. The effect of avascular necrosis on Russe bone grafting for scaphoid nonunion. *J Hand Surg [Am]* 1985;10:597-605.

3. Inoue G, Sakuma M. The natural history of scaphoid non-union: radiographical and clinical analysis in 102 cases. *Arch Orthop Trauma Surg* 1996;115:1-4.
4. Shah J, Jones WA. Factors affecting the outcome in 50 cases of scaphoid non-union treated with Herbert screw fixation. *J Hand Surg [Br]* 1998;23:680-5.
5. Schuind F, Haentjens P, Van Innis F, et al. Prognostic factors in the treatment of carpal scaphoid non-unions. *J Hand Surg [Am]* 1999;24:761-76.
6. Straw RG, Davis TR, Dias JJ. Scaphoid nonunion: treatment with a pedicled vascularised bone graft based on the 1,2 intercompartmental supraretrinacular branch of the radial artery. *J Hand Surg [Br]* 2002;27:413-16.
7. Boyer MI, von Schroeder HP, Axelrod TS. Scaphoid non-union with avascular necrosis of the proximal pole: treatment with a vascularised bone graft from the dorsum of the distal radius. *J Hand Surg [Br]* 1998;23:686-90.
8. Gabl M, Reinhart C, Lutz M, et al. Vascularized bone graft from the iliac crest for the treatment of non-union of the proximal part of the scaphoid with an avascular fragment. *J Bone Joint Surg [Am]* 1999;81-A:1414-28.
9. Steinmann SP, Bishop AT, Berger RA. Use of the 1,2 intercompartmental supraretrinacular artery as a vascularised pedicled bone graft for difficult scaphoid non-union. *J Hand Surg [Am]* 2002;27:391-401.
10. Uerpaipakit C, Leechavengvongs S, Witoonchart K. Primary vascularised distal radius bone graft for non-union of the scaphoid. *J Hand Surg [Br]* 2000;25:266-70.
11. Malizos KN, Dailiana ZH, Zirou M, et al. Longstanding nonunions of scaphoid fractures with bone loss: successful reconstruction with vascularized bone grafts. *J Hand Surg [Br]* 2001;26:330-4.
12. Chang MA, Bishop AT, Moran SL, Shin AY. The outcomes and complications of 1,2 intercompartmental supraretrinacular artery pedicled vascularized bone grafting or scaphoid nonunions. *J Hand Surg [Am]* 2006;31:387-96.
13. O'Driscoll SW, Recklies AD, Poole AR. Chondrogenesis in periosteal explants: an organ culture model for in vitro study. *J Bone Joint Surg [Am]* 1994;76-A:1042-51.
14. O'Driscoll SW, Salter RB. The induction of neochondrogenesis in free intra-articular periosteal autografts under the influence of continuous passive motion: an experimental investigation in the rabbit. *J Bone Joint Surg [Am]* 1984;66-A:1248-57.
15. Rubak JM. Osteochondrogenesis of free periosteal grafts in the rabbit iliac crest. *Acta Orthop Scand* 1983;54:826-31.
16. Thorogood P. In vitro studies on skeletogenic potential of membrane bone periosteal cells. *J Embryol Exp Morphol* 1979;54:185-207.
17. Wakitani S, Goto T, Pineda SJ, et al. Mesenchymal cell-based repair of large, full-thickness defects of articular cartilage. *J Bone Joint Surg [Am]* 1994;76-A:579-92.
18. Kubota Y, Shirasuna K. The use of free-periosteum for secondary bone grafting to the maxillary alveolar clefts. *Ann Plast Surg* 2005;55:599-602.
19. Reynders P, Becker JH, Broos P. Osteogenic ability of free periosteal autografts in tibial fractures with severe soft tissue damage: an experimental study. *J Orthop Trauma* 1999;13:121-8.
20. Burstein FD, Canalis RF, Canalis EM, Ariyan M. Scanning electron microscopy and gel electrophoresis of vascularized periosteal autografts. *Plast Reconstr Surg* 1989;83:500-10.
21. Doi K, Sakai K. Vascularised periosteal bone graft from the supracondylar region of the femur. *Microsurgery* 1994;15:305-15.
22. Sakai K, Doi K, Kawai S. Free vascularised thin corticoperiosteal graft. *Plast Reconstr Surg* 1991;87:290-8.
23. Zhu SX, Shen YN, Lu SB, Zhang BX, Wang JF. Transposition of iliac periosteum with vascular pedicle and compression screw fixation for transcervical fracture of femur. *Chin Med J* 1989;102:34-40.
24. Dias JJ. Definition of union after acute fracture and surgery for fracture nonunion of the scaphoid. *J Hand Surg [Br]* 2001;26:321-25.
25. Bain GI, Bennett JD, Richards RS, Slethaug GP, Roth JH. Longitudinal computed tomography of the scaphoid: a new technique. *Skeletal Radiol* 1995;24:271-3.
26. Ito Y, Fitzsimmons JS, Sanyal A, et al. Localisation of chondrocyte precursors in periosteum. *Osteoarthritis Cartilage* 2001;9:215-23.
27. Li M, Amizuka N, Oda K, et al. Histological evidence of the initial chondrogenesis and osteogenesis in the periosteum of a rib fractured model: implications of osteocyte involvement in periosteal chondrogenesis. *Microsc Res Tech* 2004;1:330-42.
28. Nakahara H, Bruder SP, Goldberg VW, Caplan AI. In vivo osteochondrogenic potential of cultured cells derived from periosteum. *Clin Orthop* 1990;259:223-32.
29. Fukumoto T, Sanyal A, Fitzsimmons JS, O'Driscoll SW. Expression of beta 1 integrins during periosteal chondrogenesis. *Osteoarthritis Cartilage* 2002;10:223-32.
30. Stevens MM, Marini RP, Martin I, Langer R, Prasad Shastri V. FGF-2 enhances TGF-beta-1 induced periosteal chondrogenesis. *J Orthop Res* 2004;22:1114-19.
31. Ishida H, Tamai S, Yajima H, et al. Histologic and biochemical analyses of osteogenic capacity of vascularised periosteum. *Plast Reconstr Surg* 1996;97:512-18.

32. **Dailiana ZH, Shiamishis G, Niokou D, Ioachim E, Malizos KN.** Heterotopic neo-osteogenesis from vascularized periosteum and bone grafts. *J Trauma* 2002;53:934-8.
33. **Vogelin MDE, Jones NF, Lieberman JR, et al.** Prefabrication of bone by use of a vascularized periosteal flap and bone morphogenetic protein. *Plast Reconstr Surg* 2002;109:190-8.
34. **Vogelin E, Jones NF, Huang JI, Brekke JH, Lieberman JR.** Healing of a critical-sized defect in the rat femur with the use of a vascularized periosteal flap, a biodegradable matrix, and bone morphogenetic protein. *J Bone Joint Surg [Am]* 2005;87-A:1323-31.
35. **Zhu SX, Zhou MW.** Iliac periosteal graft with vascular pedicle in the treatment of vascular necrosis of the femoral head: experimental study and clinical application. *Clin Med J* 1992;105:849-55.
36. **Burstein FD, Canalis RF.** Studies on the osteogenic potential of vascularised periosteum: behavior of periosteal flaps transferred into soft tissues. *Otolaryngol Head Neck Surg* 1985;93:731-5.
37. **Canalis RF, Burstein FD, Dickman O, Berke G.** Early structural changes in vascularised periosteal flaps studies in situ. *Am J Otolaryngol* 1991;12:90-5.
38. **Gallay SH, Miura Y, Commisso CN, Fitzsimmons JJ, O'Driscoll SW.** Relationship of donor size to chondrogenic potential of periosteum in vitro. *J Orthop Res* 1994;12:515-25.
39. **O'Driscoll SW, Saris DB, Ito Y, Fitzsimmons JS.** The chondrogenic potential of periosteum decreases with age. *J Orthop Res* 2001;19:95-103.