

Topology of group actions, Morse theory and symplectic invariants

joint work with Laurent Côté

Yusuf Barış Kartal

University of Edinburgh

March 11, 2024

Morse theory: A brief introduction

Morse theory: study of the manifold topology using functions and their gradient flow

Morse theory: A brief introduction

Morse theory: study of the manifold topology using functions and their gradient flow

- functions can be complicated, restrict to a special class: Morse functions, such as $f = x^2 - y^2$ on \mathbb{R}^2

Morse theory: A brief introduction

Morse theory: study of the manifold topology using functions and their gradient flow

- functions can be complicated, restrict to a special class: Morse functions, such as $f = x^2 - y^2$ on \mathbb{R}^2
- a Morse function f has isolated critical points


Morse theory: A brief introduction

Morse theory: study of the manifold topology using functions and their gradient flow

- functions can be complicated, restrict to a special class: Morse functions, such as $f = x^2 - y^2$ on \mathbb{R}^2
- a Morse function f has isolated critical points

Example

The surface of revolution of $x^2 + (z - 1)^2 = 4$ around x -axis (torus), and the Morse function $f = z$ on it. Blue points are the critical points.



Floer homology: Invariants defined using Morse theory on “infinite dimensional spaces”.

From Morse to Floer

Floer homology: Invariants defined using Morse theory on “infinite dimensional spaces”.

Floer defined hugely successful invariants in symplectic/low dimensional topology.

From Morse to Floer

Floer homology: Invariants defined using Morse theory on “infinite dimensional spaces”.

Floer defined hugely successful invariants in symplectic/low dimensional topology.

- e.g. solution to a major conjecture by Vladimir Arnold

Floer homology: Invariants defined using Morse theory on “infinite dimensional spaces”.

Floer defined hugely successful invariants in symplectic/low dimensional topology.

- e.g. solution to a major conjecture by Vladimir Arnold
- gives rise to (half of) a mathematical formulation of mirror symmetry phenomenon in physics

Floer homology: Invariants defined using Morse theory on “infinite dimensional spaces”.

Floer defined hugely successful invariants in symplectic/low dimensional topology.

- e.g. solution to a major conjecture by Vladimir Arnold
- gives rise to (half of) a mathematical formulation of mirror symmetry phenomenon in physics
- applications to enumerative geometry (e.g. work of McLean)

Floer homology: Invariants defined using Morse theory on “infinite dimensional spaces”.

Floer defined hugely successful invariants in symplectic/low dimensional topology.

- e.g. solution to a major conjecture by Vladimir Arnold
- gives rise to (half of) a mathematical formulation of mirror symmetry phenomenon in physics
- applications to enumerative geometry (e.g. work of McLean)
- central focus of modern symplectic topology: shapes the field

Floer homology: Invariants defined using Morse theory on “infinite dimensional spaces”.

Floer defined hugely successful invariants in symplectic/low dimensional topology.

- e.g. solution to a major conjecture by Vladimir Arnold
- gives rise to (half of) a mathematical formulation of mirror symmetry phenomenon in physics
- applications to enumerative geometry (e.g. work of McLean)
- central focus of modern symplectic topology: shapes the field
- connects symplectic topology with other fields including algebraic geometry, algebraic topology (string topology), non-commutative geometry, low dimensional topology, ...

From Morse to Floer (cont'd)

Morse theory: Study of manifold topology using the data of critical points of Morse functions and their gradient flow lines. This gives a way to compute homology (classical).

From Morse to Floer (cont'd)

Morse theory: Study of manifold topology using the data of critical points of Morse functions and their gradient flow lines. This gives a way to compute homology (classical).

The data of critical points and gradient trajectories is all we need to define a “Floer theory”.

From Morse to Floer (cont'd)

Morse theory: Study of manifold topology using the data of critical points of Morse functions and their gradient flow lines. This gives a way to compute homology (classical).

The data of critical points and gradient trajectories is all we need to define a “Floer theory”.

- Let M be a nice open symplectic manifold (e.g. any open surface or $S^3 \times \mathbb{R}^3$).

From Morse to Floer (cont'd)

Morse theory: Study of manifold topology using the data of critical points of Morse functions and their gradient flow lines. This gives a way to compute homology (classical).

The data of critical points and gradient trajectories is all we need to define a “Floer theory”.


- Let M be a nice open symplectic manifold (e.g. any open surface or $S^3 \times \mathbb{R}^3$).
- Let $\mathcal{L}M$ be the infinite dimensional space of closed curves in M (more abstractly space of maps from S^1 to M).

From Morse to Floer (cont'd)

Morse theory: Study of manifold topology using the data of critical points of Morse functions and their gradient flow lines. This gives a way to compute homology (classical).

The data of critical points and gradient trajectories is all we need to define a “Floer theory”.

- Let M be a nice open symplectic manifold (e.g. any open surface or $S^3 \times \mathbb{R}^3$).
- Let $\mathcal{L}M$ be the infinite dimensional space of closed curves in M (more abstractly space of maps from S^1 to M).



From Morse to Floer (cont'd)

Definition

“Morse theory” on $\mathcal{L}M$ is called *symplectic cohomology*, and denoted by $SH(M; \mathbb{Z})$.

From Morse to Floer (cont'd)

Definition

“Morse theory” on $\mathcal{L}M$ is called *symplectic cohomology*, and denoted by $SH(M; \mathbb{Z})$.

Example

$$SH(\mathbb{C}^*; \mathbb{Z}) = \mathbb{Z}[z, z^{-1}].$$

From Morse to Floer (cont'd)

Definition

“Morse theory” on $\mathcal{L}M$ is called *symplectic cohomology*, and denoted by $SH(M; \mathbb{Z})$.

Example

$$SH(\mathbb{C}^*; \mathbb{Z}) = \mathbb{Z}[z, z^{-1}].$$

Remark

$SH(M; \mathbb{Z})$ is a generalization of quantum cohomology to non-compact manifolds.

From Morse to Floer (cont'd)

Definition

“Morse theory” on $\mathcal{L}M$ is called *symplectic cohomology*, and denoted by $SH(M; \mathbb{Z})$.

Example

$$SH(\mathbb{C}^*; \mathbb{Z}) = \mathbb{Z}[z, z^{-1}].$$

Remark

$SH(M; \mathbb{Z})$ is a generalization of quantum cohomology to non-compact manifolds.

Warning: The “Morse function” on $\mathcal{L}M$ depends on the symplectic geometry of M , and $SH(M; \mathbb{Z})$ is a symplectic invariant. It depends more than the topology of $\mathcal{L}M$.

Floer theory meets homotopy theory

Cohen-Jones-Segal ('95): introduce a framework to bridge Floer theory and homotopy theory

Floer theory meets homotopy theory

Cohen-Jones-Segal ('95): introduce a framework to bridge Floer theory and homotopy theory

Space \longrightarrow Homology groups

Floer theory meets homotopy theory

Cohen-Jones-Segal ('95): introduce a framework to bridge Floer theory and homotopy theory

Space \longrightarrow Homology groups

"Floer homotopy type"? \longrightarrow Floer homology groups
(a space)

Floer theory meets homotopy theory

Cohen-Jones-Segal ('95): introduce a framework to bridge Floer theory and homotopy theory

Space \longrightarrow Homology groups

"Floer homotopy type"? \longrightarrow Floer homology groups
(a space)

Outcome: Even more powerful invariants (e.g. by taking K -theory of this space or applying other generalized cohomology theories).

Floer theory meets homotopy theory

Cohen-Jones-Segal ('95): introduce a framework to bridge Floer theory and homotopy theory

Space \longrightarrow Homology groups

"Floer homotopy type"? \longrightarrow Floer homology groups
(a space)

Outcome: Even more powerful invariants (e.g. by taking K -theory of this space or applying other generalized cohomology theories). The field was dormant for a while, but

Floer theory meets homotopy theory

Cohen-Jones-Segal ('95): introduce a framework to bridge Floer theory and homotopy theory

Space \longrightarrow Homology groups

"Floer homotopy type"? \longrightarrow Floer homology groups
(a space)

Outcome: Even more powerful invariants (e.g. by taking K -theory of this space or applying other generalized cohomology theories). The field was dormant for a while, but

Recently: applications to dynamics (e.g. Abouzaid-Blumberg '21) (improvements to Arnold conjecture).

Floer theory meets homotopy theory

Cohen-Jones-Segal ('95): introduce a framework to bridge Floer theory and homotopy theory

Space \longrightarrow Homology groups

“Floer homotopy type”?
(a space) \longrightarrow Floer homology
groups

Outcome: Even more powerful invariants (e.g. by taking K -theory of this space or applying other generalized cohomology theories). The field was dormant for a while, but

Recently: applications to dynamics (e.g. Abouzaid-Blumberg '21) (improvements to Arnold conjecture).

Large '21: Defined a “space” $SH(M; \mathbb{S})$, which is a symplectic invariant.

Lie group actions?

- The group $SH(M; \mathbb{Z})$ is highly structured and “the space” $SH(M; \mathbb{S})$ is expected to be even more structured.

Lie group actions?

- The group $SH(M; \mathbb{Z})$ is highly structured and “the space” $SH(M; \mathbb{S})$ is expected to be even more structured.
- A crucial part of the expected structure is a circle action.

Lie group actions?

- The group $SH(M; \mathbb{Z})$ is highly structured and “the space” $SH(M; \mathbb{S})$ is expected to be even more structured.
- A crucial part of the expected structure is a circle action.
- More generally, it is frequently expected that Floer invariants carry the action of a compact Lie group (e.g. a circle action).

Lie group actions?

- The group $SH(M; \mathbb{Z})$ is highly structured and “the space” $SH(M; \mathbb{S})$ is expected to be even more structured.
- A crucial part of the expected structure is a circle action.
- More generally, it is frequently expected that Floer invariants carry the action of a compact Lie group (e.g. a circle action).

Observe: $\mathcal{L}M$ carries a circle action (given by changing the starting point of the closed curve, i.e. by rotating the closed loop). Can one induce an action on its “Morse theory” $SH(M; \mathbb{S})$?

Lie group actions?

- The group $SH(M; \mathbb{Z})$ is highly structured and “the space” $SH(M; \mathbb{S})$ is expected to be even more structured.
- A crucial part of the expected structure is a circle action.
- More generally, it is frequently expected that Floer invariants carry the action of a compact Lie group (e.g. a circle action).

Observe: \mathcal{LM} carries a circle action (given by changing the starting point of the closed curve, i.e. by rotating the closed loop). Can one induce an action on its “Morse theory” $SH(M; \mathbb{S})$?

Problem: The setup of Morse/Floer theory is not compatible with group actions.

Summary of main achievements

Our main achievements are:

Summary of main achievements

Our main achievements are:

- We introduce a setup that works equivariantly by going beyond Morse.

Summary of main achievements

Our main achievements are:

- We introduce a setup that works equivariantly by going beyond Morse.
- We define equivariant and “space level” versions of major invariants in symplectic topology.

Summary of main achievements

Our main achievements are:

- We introduce a setup that works equivariantly by going beyond Morse.
- We define equivariant and “space level” versions of major invariants in symplectic topology.
- We make a major upgrade to a well-known theorem (Viterbo isomorphism).

Summary of main achievements

Our main achievements are:

- We introduce a setup that works equivariantly by going beyond Morse.
- We define equivariant and “space level” versions of major invariants in symplectic topology.
- We make a major upgrade to a well-known theorem (Viterbo isomorphism).
- We give applications connecting Floer theory to the equivariant homotopy theory- a deep subfield of homotopy theory.

Lie group actions and Morse–Bott functions

Problem: Often, there are no equivariant Morse functions.

Lie group actions and Morse–Bott functions

Problem: Often, there are no equivariant Morse functions.

Solution: Extend the setup to work with *Morse–Bott* functions.

Lie group actions and Morse–Bott functions

Problem: Often, there are no equivariant Morse functions.

Solution: Extend the setup to work with *Morse–Bott* functions.

A *Morse–Bott function* is a special type of function whose critical set is a manifold. E.g. $f = y^2 - z^2$

Lie group actions and Morse–Bott functions

Problem: Often, there are no equivariant Morse functions.

Solution: Extend the setup to work with *Morse–Bott* functions.

A *Morse–Bott function* is a special type of function whose critical set is a manifold. E.g. $f = y^2 - z^2$

Good news: There are often plenty of equivariant Morse–Bott functions.

Lie group actions and Morse–Bott functions

Problem: Often, there are no equivariant Morse functions.


Solution: Extend the setup to work with *Morse–Bott* functions.

A *Morse–Bott function* is a special type of function whose critical set is a manifold. E.g. $f = y^2 - z^2$

Good news: There are often plenty of equivariant Morse–Bott functions.

Example

Let N be the torus obtained by revolving $(x - 1)^2 + z^2 = 4$ around z -axis. Then, $f = z$ is Morse–Bott and equivariant with respect to rotation. Critical set consists of two blue circles.



Main results

We package the data of $\text{crit}(f)$ and the gradient trajectories into an abstract object called a *framed Morse-Bott flow category*.

Main results

We package the data of $\text{crit}(f)$ and the gradient trajectories into an abstract object called a *framed Morse-Bott flow category*. We prove

Theorem (Côté-K.)

One can recover the “equivariant topology” of N from the associated framed Morse-Bott flow category, i.e. from the data of critical manifolds and the gradient trajectories between them.

Main results

We package the data of $\text{crit}(f)$ and the gradient trajectories into an abstract object called a *framed Morse-Bott flow category*. We prove

Theorem (Côté-K.)

One can recover the “equivariant topology” of N from the associated framed Morse-Bott flow category, i.e. from the data of critical manifolds and the gradient trajectories between them.

- One can now use Morse–Bott theory to study equivariant cohomology.

Main results

We package the data of $\text{crit}(f)$ and the gradient trajectories into an abstract object called a *framed Morse-Bott flow category*. We prove

Theorem (Côté-K.)

One can recover the “equivariant topology” of N from the associated framed Morse-Bott flow category, i.e. from the data of critical manifolds and the gradient trajectories between them.

- One can now use Morse–Bott theory to study equivariant cohomology.
- The proof involves construction of a “space” associated to such an abstract object and new tools to induce maps on this “space”.

Main results

We package the data of $\text{crit}(f)$ and the gradient trajectories into an abstract object called a *framed Morse-Bott flow category*. We prove

Theorem (Côté-K.)

One can recover the “equivariant topology” of N from the associated framed Morse-Bott flow category, i.e. from the data of critical manifolds and the gradient trajectories between them.

- One can now use Morse–Bott theory to study equivariant cohomology.
- The proof involves construction of a “space” associated to such an abstract object and new tools to induce maps on this “space”.
- **Most importantly:** the construction works for “infinite dimensional Morse theory”, aka Floer theory.

Main results (cont'd)

Let M be nice symplectic (e.g. an affine variety or cotangent bundle), and consider $SH(M; \mathbb{S})$ - the “space” obtained from the Morse theory of the infinite dimensional space \mathcal{LM} .

Main results (cont'd)

Let M be nice symplectic (e.g. an affine variety or cotangent bundle), and consider $SH(M; \mathbb{S})$ - the “space” obtained from the Morse theory of the infinite dimensional space $\mathcal{L}M$.

Theorem (Côté-K.)

- a. $SH(M; \mathbb{S})$ can be endowed with a circle action.

Main results (cont'd)

Let M be nice symplectic (e.g. an affine variety or cotangent bundle), and consider $SH(M; \mathbb{S})$ - the “space” obtained from the Morse theory of the infinite dimensional space $\mathcal{L}M$.

Theorem (Côté-K.)

- a. $SH(M; \mathbb{S})$ can be endowed with a circle action.
- b. (Viterbo isomorphism) if $M = T^*Q$, then $SH(M; \mathbb{S})$ is circle equivariantly equivalent to $\mathcal{L}Q$ (roughly).

Main results (cont'd)

Let M be nice symplectic (e.g. an affine variety or cotangent bundle), and consider $SH(M; \mathbb{S})$ - the “space” obtained from the Morse theory of the infinite dimensional space $\mathcal{L}M$.

Theorem (Côté-K.)

- a. $SH(M; \mathbb{S})$ can be endowed with a circle action.
 - b. (Viterbo isomorphism) if $M = T^*Q$, then $SH(M; \mathbb{S})$ is circle equivariantly equivalent to $\mathcal{L}Q$ (roughly).
- Circle actions are at the heart of the structures the “space” $SH(M; \mathbb{S})$ is expected to carry. It is crucial for future applications.

Main results (cont'd)

Let M be nice symplectic (e.g. an affine variety or cotangent bundle), and consider $SH(M; \mathbb{S})$ - the “space” obtained from the Morse theory of the infinite dimensional space $\mathcal{L}M$.

Theorem (Côté-K.)

- a. $SH(M; \mathbb{S})$ can be endowed with a circle action.
 - b. (Viterbo isomorphism) if $M = T^*Q$, then $SH(M; \mathbb{S})$ is circle equivariantly equivalent to $\mathcal{L}Q$ (roughly).
-
- Circle actions are at the heart of the structures the “space” $SH(M; \mathbb{S})$ is expected to carry. It is crucial for future applications.
 - Viterbo isomorphism connects symplectic topology to the topology of loop spaces (to a field called string topology).

An application

Problem: Symplectic cohomology is not sensitive to topology of M .

An application

Problem: Symplectic cohomology is not sensitive to topology of M .

E.g. $SH(M \times \mathbb{C}; \mathbb{Z}) = 0$ for all M . One can recover $H^*(M; \mathbb{Q})$ from a version of the equivariant symplectic cohomology, but there is no known way of recovering torsion part of $H^*(M; \mathbb{Z})$ from symplectic cohomology.

An application

Problem: Symplectic cohomology is not sensitive to topology of M .

E.g. $SH(M \times \mathbb{C}; \mathbb{Z}) = 0$ for all M . One can recover $H^*(M; \mathbb{Q})$ from a version of the equivariant symplectic cohomology, but there is no known way of recovering torsion part of $H^*(M; \mathbb{Z})$ from symplectic cohomology.

Theorem (Côté-K.)

One can recover the integral homology, complex K-theory (and more) from versions of the equivariant cohomology of the “space” $SH(M; \mathbb{S})$.

An application

Problem: Symplectic cohomology is not sensitive to topology of M .

E.g. $SH(M \times \mathbb{C}; \mathbb{Z}) = 0$ for all M . One can recover $H^*(M; \mathbb{Q})$ from a version of the equivariant symplectic cohomology, but there is no known way of recovering torsion part of $H^*(M; \mathbb{Z})$ from symplectic cohomology.

Theorem (Côté-K.)

One can recover the integral homology, complex K -theory (and more) from versions of the equivariant cohomology of the “space” $SH(M; \mathbb{S})$.

- This is surprising as symplectic cohomology seemingly loses all the information about the topology of M .

An application

Problem: Symplectic cohomology is not sensitive to topology of M .

E.g. $SH(M \times \mathbb{C}; \mathbb{Z}) = 0$ for all M . One can recover $H^*(M; \mathbb{Q})$ from a version of the equivariant symplectic cohomology, but there is no known way of recovering torsion part of $H^*(M; \mathbb{Z})$ from symplectic cohomology.

Theorem (Côté-K.)

One can recover the integral homology, complex K-theory (and more) from versions of the equivariant cohomology of the “space” $SH(M; \mathbb{S})$.

- This is surprising as symplectic cohomology seemingly loses all the information about the topology of M .
- It connects symplectic topology with equivariant (and chromatic) homotopy theory.

An application

Problem: Symplectic cohomology is not sensitive to topology of M .

E.g. $SH(M \times \mathbb{C}; \mathbb{Z}) = 0$ for all M . One can recover $H^*(M; \mathbb{Q})$ from a version of the equivariant symplectic cohomology, but there is no known way of recovering torsion part of $H^*(M; \mathbb{Z})$ from symplectic cohomology.

Theorem (Côté-K.)

One can recover the integral homology, complex K -theory (and more) from versions of the equivariant cohomology of the “space” $SH(M; \mathbb{S})$.

- This is surprising as symplectic cohomology seemingly loses all the information about the topology of M .
- It connects symplectic topology with equivariant (and chromatic) homotopy theory.
- Connections to mirror symmetry (e.g. conjectures about the complex K -theory of mirror pairs).

The vision I (joint work in progress with Côté):

This work is foundational for a long term project.

The vision I (joint work in progress with Côté):

This work is foundational for a long term project. *Fukaya categories* are abstract objects that are of significance in symplectic topology and mirror symmetry.

The vision I (joint work in progress with Côté):

This work is foundational for a long term project. *Fukaya categories* are abstract objects that are of significance in symplectic topology and mirror symmetry.

Our goal: To study a major invariant (THH) associated to Fukaya categories in terms of the symplectic/quantum cohomology. The structure involves a circle action and a Frobenius map. We plan to investigate further applications into

The vision I (joint work in progress with Côté):

This work is foundational for a long term project. *Fukaya categories* are abstract objects that are of significance in symplectic topology and mirror symmetry.

Our goal: To study a major invariant (THH) associated to Fukaya categories in terms of the symplectic/quantum cohomology. The structure involves a circle action and a Frobenius map. We plan to investigate further applications into

- arithmetic refinements of mirror symmetry: e.g. relate arithmetic invariants with the enumerative invariants of the mirror

The vision I (joint work in progress with Côté):

This work is foundational for a long term project. *Fukaya categories* are abstract objects that are of significance in symplectic topology and mirror symmetry.

Our goal: To study a major invariant (THH) associated to Fukaya categories in terms of the symplectic/quantum cohomology. The structure involves a circle action and a Frobenius map. We plan to investigate further applications into

- arithmetic refinements of mirror symmetry: e.g. relate arithmetic invariants with the enumerative invariants of the mirror
- understanding important homotopy theoretic invariants (algebraic K -theory) of Fukaya categories

The vision I (joint work in progress with Côté):

This work is foundational for a long term project. *Fukaya categories* are abstract objects that are of significance in symplectic topology and mirror symmetry.

Our goal: To study a major invariant (THH) associated to Fukaya categories in terms of the symplectic/quantum cohomology. The structure involves a circle action and a Frobenius map. We plan to investigate further applications into

- arithmetic refinements of mirror symmetry: e.g. relate arithmetic invariants with the enumerative invariants of the mirror
- understanding important homotopy theoretic invariants (algebraic K -theory) of Fukaya categories
- (ambitious) Fukaya categorical versions of major conjectures in number theory (e.g. non-commutative Weil conjectures)?

The vision I (joint work in progress with Côté):

This work is foundational for a long term project. *Fukaya categories* are abstract objects that are of significance in symplectic topology and mirror symmetry.

Our goal: To study a major invariant (THH) associated to Fukaya categories in terms of the symplectic/quantum cohomology. The structure involves a circle action and a Frobenius map. We plan to investigate further applications into

- arithmetic refinements of mirror symmetry: e.g. relate arithmetic invariants with the enumerative invariants of the mirror
- understanding important homotopy theoretic invariants (algebraic K -theory) of Fukaya categories
- (ambitious) Fukaya categorical versions of major conjectures in number theory (e.g. non-commutative Weil conjectures)?

The vision II (joint work in progress with Côté and Mak):

Expectations inspired from physics (Teleman et al): Let M be a symplectic manifold with a nice (Hamiltonian) G -action. Then it is expected:

The vision II (joint work in progress with Côté and Mak):

Expectations inspired from physics (Teleman et al): Let M be a symplectic manifold with a nice (Hamiltonian) G -action. Then it is expected:

- Algebras that occur naturally in representation theory (K -theoretic and elliptic Coulomb branch algebras) act on the equivariant symplectic/quantum cohomology.

The vision II (joint work in progress with Côté and Mak):

Expectations inspired from physics (Teleman et al): Let M be a symplectic manifold with a nice (Hamiltonian) G -action. Then it is expected:

- Algebras that occur naturally in representation theory (K -theoretic and elliptic Coulomb branch algebras) act on the equivariant symplectic/quantum cohomology.
- The action is compatible with the various structures.

The vision II (joint work in progress with Côté and Mak):

Expectations inspired from physics (Teleman et al): Let M be a symplectic manifold with a nice (Hamiltonian) G -action. Then it is expected:

- Algebras that occur naturally in representation theory (K -theoretic and elliptic Coulomb branch algebras) act on the equivariant symplectic/quantum cohomology.
- The action is compatible with the various structures.

We are working verification of these. This will further bridge symplectic topology to representation theory.

Thank you!