

# Base-extension semantics for Intuitionistic Linear Logic

Yli Buzoku

Department of Computer Science  
University College London

May 9, 2023

# Goals for this talk

- To briefly introduce Intuitionistic Linear Logic.
- To give an overview of the setup of Base-Extension Semantics in the substructural setting.
- Present a Base-extension semantics for Intuitionistic Linear Logic.

# Presentation root directory

- 1 Overview of Intuitionistic Linear Logic
- 2 Base-extension Semantics for ILL
- 3 Including the exponential

# Presentation root directory

- 1 Overview of Intuitionistic Linear Logic
- 2 Base-extension Semantics for ILL
- 3 Including the exponential

# Notation

- $\mathbb{A}$  represents a fixed, countably infinite set of propositional atoms.
- Lower case latin letters represent propositional atoms
- Upper case latin letters represent finite multisets of propositional atoms.
- The empty multiset is denoted  $\emptyset$ .
- The sum of two multisets  $P$  and  $Q$  is denoted  $P, Q$ .
- Lower case greek letters represent ILL formulas.
- Upper case greek letters represent ILL finite multisets thereof.
- Atomic multiset is taken to mean multiset of propositional atoms.

# A natural deduction system for ILL

$$\begin{array}{c}
 \frac{}{\varphi \vdash \varphi} \text{Ax} \\
 \\
 \frac{\Gamma, \varphi \vdash \psi}{\Gamma \vdash \varphi \multimap \psi} \multimap\text{-I} \qquad \frac{\Gamma \vdash \varphi \multimap \psi \quad \Delta \vdash \varphi}{\Gamma, \Delta \vdash \psi} \multimap\text{-E} \\
 \\
 \frac{\Gamma \vdash \varphi \quad \Delta \vdash \psi}{\Gamma, \Delta \vdash \varphi \otimes \psi} \otimes\text{-I} \qquad \frac{\Gamma \vdash \varphi \otimes \psi \quad \Delta, \varphi, \psi \vdash \chi}{\Gamma, \Delta \vdash \chi} \otimes\text{-E} \\
 \\
 \frac{}{\vdash 1} 1\text{-I} \qquad \frac{\Gamma \vdash \varphi \quad \Delta \vdash 1}{\Gamma, \Delta \vdash \varphi} 1\text{-E} \\
 \\
 \frac{\Gamma \vdash \varphi \quad \Gamma \vdash \psi}{\Gamma \vdash \varphi \& \psi} \&\text{-I} \qquad \frac{\Gamma \vdash \varphi_0 \& \varphi_1}{\Gamma \vdash \varphi_i} \&\text{-E}_i \\
 \\
 \frac{\Gamma \vdash \varphi_i}{\Gamma \vdash \varphi_0 \oplus \varphi_1} \oplus\text{-I}_i \qquad \frac{\Gamma \vdash \varphi \oplus \psi \quad \Delta, \varphi \vdash \chi \quad \Delta, \psi \vdash \chi}{\Gamma, \Delta \vdash \chi} \oplus\text{-E} \\
 \\
 \text{No 0 intro rule} \qquad \frac{\Gamma \vdash 0}{\Gamma \vdash \varphi} 0\text{-E}
 \end{array}$$

# A natural deduction system for ILL (cont.)

$$\frac{\Gamma_1 \vdash !\psi_1 \quad \dots \quad \Gamma_n \vdash !\psi_n \quad !\psi_1, \dots, !\psi_n \vdash \varphi}{\Gamma_1, \dots, \Gamma_n \vdash !\varphi} \text{!-Promotion}$$

$$\frac{\Gamma \vdash !\varphi \quad \Delta, \varphi \vdash \psi}{\Gamma, \Delta \vdash \psi} \text{!-Dereliction}$$

$$\frac{\Gamma \vdash !\varphi \quad \Delta \vdash \psi}{\Gamma, \Delta \vdash \psi} \text{!-Weakening}$$

$$\frac{\Gamma \vdash !\varphi \quad \Delta, !\varphi, !\varphi \vdash \psi}{\Gamma, \Delta \vdash \psi} \text{!-Contraction}$$

What is the general form of an inference figure?



What is the general form of an inference figure?

$$\frac{\Gamma \vdash \varphi \quad \Delta \vdash \psi}{\Gamma, \Delta \vdash \varphi \otimes \psi} \otimes\text{-I}$$

What is the general form of an inference figure?

$$\frac{\Gamma \vdash \varphi \quad \Gamma \vdash \psi}{\Gamma \vdash \varphi \ \& \ \psi} \&-I$$

What is the general form of an inference figure?

What is the general form of an inference figure?

$$\frac{\Gamma_1 \vdash \varphi_1 \quad \dots \quad \Gamma_n \vdash \varphi_n}{\Gamma_1 \dots \Gamma_n \vdash \psi}$$

$$\frac{\Gamma \vdash \varphi_1 \quad \dots \quad \Gamma \vdash \varphi_n}{\Gamma \vdash \psi}$$

What is the general form of an inference figure?

$$\frac{\Gamma_1 \vdash \varphi_1 \quad \dots \quad \Gamma_n \vdash \varphi_n}{\Gamma_1 \dots \Gamma_n \vdash \psi}$$

$$\frac{\Gamma \vdash \varphi_1 \quad \dots \quad \Gamma \vdash \varphi_n}{\Gamma \vdash \psi}$$

$$\frac{\Gamma \vdash \varphi \oplus \psi \quad \Delta, \varphi \vdash \chi \quad \Delta, \psi \vdash \chi}{\Gamma, \Delta \vdash \chi} \oplus\text{-E}$$

What is the general form of an inference figure?

$$\frac{\Gamma \vdash \varphi \oplus \psi \quad \Delta, \varphi \vdash \chi \quad \Delta, \psi \vdash \chi}{\Gamma, \Delta \vdash \chi} \oplus\text{-E}$$

What is the general form of an inference figure?

$$\frac{\Gamma \vdash \varphi \oplus \psi \quad \Delta \{ \varphi \vdash \chi \quad \psi \vdash \chi \}}{\Gamma, \Delta \vdash \chi} \oplus\text{-E}$$

What is the general form of an inference figure?

$$\frac{\Gamma\{\emptyset \vdash \varphi \oplus \psi\} \quad \Delta\{\varphi \vdash \chi \quad \psi \vdash \chi\}}{\Gamma, \Delta \vdash \chi} \oplus\text{-E}$$



What is the general form of an inference figure?

$$\frac{\Gamma\{\emptyset \vdash \varphi \oplus \psi\} \quad \Delta\{\varphi \vdash \chi \quad \psi \vdash \chi\}}{\Gamma, \Delta \vdash \chi} \oplus\text{-E}$$

What is the general form of an inference figure?

$$\frac{\Gamma\{\emptyset \vdash \varphi\} \quad \Delta\{\emptyset \vdash \psi\}}{\Gamma, \Delta \vdash \varphi \otimes \psi} \otimes\text{-I}$$

$$\frac{\Gamma\{\emptyset \vdash \varphi \quad \emptyset \vdash \psi\}}{\Gamma \vdash \varphi \& \psi} \&\text{-I}$$

What is the general form of an inference figure?

$$\frac{\dots \quad \Gamma_i \{ \Delta_{i_1} \vdash \varphi_{i_1} \quad \dots \quad \Delta_{i_{l_i}} \vdash \varphi_{i_{l_i}} \} \quad \dots}{\Gamma_1 \dots \Gamma_n \vdash \psi}$$

What is the general form of an inference figure?

# What is the general form of an inference figure?

Almost! What about:

$$\frac{\Gamma_1 \vdash !\psi_1 \quad \dots \quad \Gamma_n \vdash !\psi_n \quad !\psi_1, \dots, !\psi_n \vdash \varphi}{\Gamma_1, \dots, \Gamma_n \vdash !\varphi} \text{!-Promotion}$$

What is the general form of an inference figure?

$$\frac{\Gamma_1 \{\emptyset \vdash !\psi_1\} \quad \dots \quad \Gamma_n \{\emptyset \vdash !\psi_n\} \quad !\psi_1, \dots, !\psi_n \vdash \varphi}{\Gamma_1, \dots, \Gamma_n \vdash !\varphi} \text{!-Promotion}$$

# Presentation root directory

- 1 Overview of Intuitionistic Linear Logic
- 2 Base-extension Semantics for ILL
- 3 Including the exponential

# Substructural atomic derivability

## Definition

Basic rules take the following form:

$$\{(P_{1_i} \Rightarrow q_{1_i})\}_{i=1}^{l_1}, \dots, \{(P_{n_i} \Rightarrow q_{n_i})\}_{i=1}^{l_n} \Rightarrow r$$

where

Each  $P_i$  is an atomic multiset, called a premiss multiset.

Each  $q_i$  and  $r$  is an atomic proposition.

Each  $(P_i \Rightarrow q_i)$  is a pair  $(P_i, q_i)$  called an atomic sequent.

Each collection  $\{(P_{i_1} \Rightarrow q_{i_1}), \dots, (P_{i_{l_i}} \Rightarrow q_{i_{l_i}})\}$  is called an atomic box.



# Substructural atomic derivability

## Definition (Basic derivability relation)

The relation of derivability in a base  $\mathcal{B}$ , is defined inductively as so:

Ref  $p \vdash_{\mathcal{B}} p$

App Given that  $(\{(P_{1_i} \Rightarrow q_{1_i})\}_{i=1}^{l_1}, \dots, \{(P_{n_i} \Rightarrow q_{n_i})\}_{i=1}^{l_n} \Rightarrow r) \in \mathcal{B}$  and atomic multisets  $C_i$  such that the following hold:

$$C_i, P_{ij} \vdash_{\mathcal{B}} q_{ij} \text{ for all } i = 1, \dots, n \text{ and } j = 1, \dots, l_i$$

Then  $C_1, \dots, C_n \vdash_{\mathcal{B}} r$ .

# Base-extension Semantics for ILL

- (At)  $\Vdash_{\mathcal{B}}^L p$  iff  $L \vdash_{\mathcal{B}} p$ .
- ( $\otimes$ )  $\Vdash_{\mathcal{B}}^L \varphi \otimes \psi$  iff for every  $\mathcal{C} \supseteq \mathcal{B}$ , atomic multiset  $K$ , and atom  $p$ , if  $\varphi, \psi \Vdash_{\mathcal{C}}^K p$  then  $\Vdash_{\mathcal{C}}^{L,K} p$ .
- (1)  $\Vdash_{\mathcal{B}}^L 1$  iff for every  $\mathcal{C} \supseteq \mathcal{B}$ , atomic multiset  $K$ , and atom  $p$ , if  $\Vdash_{\mathcal{C}}^K p$ , then  $\Vdash_{\mathcal{C}}^{L,K} p$ .
- ( $\multimap$ )  $\Vdash_{\mathcal{B}}^L \varphi \multimap \psi$  iff  $\varphi \Vdash_{\mathcal{B}}^L \psi$ .

⋮

# Base-extension Semantics for ILL

$$\vdots$$

( $\&$ )  $\Vdash_{\mathcal{B}}^L \varphi \& \psi$  iff  $\Vdash_{\mathcal{B}}^L \varphi$  and  $\Vdash_{\mathcal{B}}^L \psi$ .

( $\oplus$ )  $\Vdash_{\mathcal{B}}^L \varphi \oplus \psi$  iff for every  $\mathcal{C} \supseteq \mathcal{B}$ , atomic multiset  $K$  and atom  $p \in \mathbb{A}$  such that  $\varphi \Vdash_{\mathcal{C}}^K p$  and  $\psi \Vdash_{\mathcal{C}}^K p$  hold, then  $\Vdash_{\mathcal{C}}^{L,K} p$ .

(0)  $\Vdash_{\mathcal{B}}^L 0$  iff  $\Vdash_{\mathcal{B}}^L p$ , for all  $p \in \mathbb{A}$ .

$$\vdots$$

# Base-extension Semantics for ILL

⋮

- ( $\circ$ )  $\Vdash_{\mathcal{B}}^L \Gamma, \Delta$  iff there are atomic multisets  $K$  and  $M$  such that  $L = K, M$  and that  $\Vdash_{\mathcal{B}}^K \Gamma$  and  $\Vdash_{\mathcal{B}}^M \Delta$ .
- ( $!$ )  $\Vdash_{\mathcal{B}}^L !\varphi$  iff for any  $\mathcal{C}$  such that  $\mathcal{C} \supseteq \mathcal{B}$ , atomic multisets  $K$  and any  $p \in \mathbb{A}$ , if for any  $\mathcal{D}$  such that  $\mathcal{D} \supseteq \mathcal{C}$ , (if  $\Vdash_{\mathcal{D}}^{\emptyset} \varphi$  then  $\Vdash_{\mathcal{D}}^L p$ ) then  $\Vdash_{\mathcal{C}}^{L,K} p$ .
- (Inf) For  $\Gamma = !\Delta$ ,  $\Theta$  being a nonempty multiset,  $\Gamma \Vdash_{\mathcal{B}}^L \varphi$  iff for all  $\mathcal{C} \supseteq \mathcal{B}$ , atomic multisets  $K$ , if  $\Vdash_{\mathcal{C}}^K \Theta$  and  $\Vdash_{\mathcal{C}}^{\emptyset} \Delta$  then  $\Vdash_{\mathcal{C}}^{L,K} \varphi$ .

# Base-extension Semantics for ILL

⋮

- ( $\circ$ )  $\Vdash_{\mathcal{B}}^L \Gamma, \Delta$  iff there are atomic multisets  $K$  and  $M$  such that  $L = K, M$  and that  $\Vdash_{\mathcal{B}}^K \Gamma$  and  $\Vdash_{\mathcal{B}}^M \Delta$ .
- ( $!$ )  $\Vdash_{\mathcal{B}}^L !\varphi$  iff for any  $\mathcal{C}$  such that  $\mathcal{C} \supseteq \mathcal{B}$ , atomic multisets  $K$  and any  $p \in \mathbb{A}$ , if for any  $\mathcal{D}$  such that  $\mathcal{D} \supseteq \mathcal{C}$ , (if  $\Vdash_{\mathcal{D}}^{\emptyset} \varphi$  then  $\Vdash_{\mathcal{D}}^L p$ ) then  $\Vdash_{\mathcal{C}}^{L,K} p$ .
- (Inf) For  $\Gamma = !\Delta$ ,  $\Theta$  being a nonempty multiset,  $\Gamma \Vdash_{\mathcal{B}}^L \varphi$  iff for all  $\mathcal{C} \supseteq \mathcal{B}$ , atomic multisets  $K$ , if  $\Vdash_{\mathcal{C}}^K \Theta$  and  $\Vdash_{\mathcal{C}}^{\emptyset} \Delta$  then  $\Vdash_{\mathcal{C}}^{L,K} \varphi$ .

Note the clause for ( $!$ ) could be written as:

$\Vdash_{\mathcal{B}}^L !\varphi$  iff for any  $\mathcal{C}$  such that  $\mathcal{C} \supseteq \mathcal{B}$ , atomic multisets  $K$  and any  $p \in \mathbb{A}$ , if  $!\varphi \Vdash_{\mathcal{C}}^K p$  then  $\Vdash_{\mathcal{C}}^{L,K} p$ .

# Notes on $\Vdash_{\mathcal{B}}^L$

The sequent  $\langle \Gamma, \varphi \rangle$  is said to be valid just in case when  $\Gamma \Vdash_{\emptyset}^{\emptyset} \varphi$ .

## Some facts about the semantics

If  $\Vdash_{\mathcal{B}}^L \varphi$  then for all  $\mathcal{C} \supseteq \mathcal{B}$  we have  $\Vdash_{\mathcal{C}}^L \varphi$ .

$\Vdash_{\mathcal{B}}^L \varphi$  iff  $\Vdash_{\mathcal{B}}^L \varphi \otimes 1$  iff  $\Vdash_{\mathcal{B}}^L \varphi, 1$

$\Vdash_{\mathcal{B}}^L \varphi$  iff for all  $\mathcal{C} \supseteq \mathcal{B}$ , atomic multisets  $K$  and atoms  $p$  such that  $\varphi \Vdash_{\mathcal{C}}^K p$  then  $\Vdash_{\mathcal{C}}^{L,K} p$ .

# Presentation root directory

- 1 Overview of Intuitionistic Linear Logic
- 2 Base-extension Semantics for ILL
- 3 Including the exponential



# What of the exponential?

*The exponential is hard...*

# What of the exponential?

*The exponential is hard...*

*But I finally got it!*

# Why was it so hard?

## Alternative approach

We want to re-introduce in a controlled manner the ability for multiple and vacuous discharge of hypotheses.

## Alternative approach

We want to re-introduce in a controlled manner the ability for multiple and vacuous discharge of hypotheses.

It is not absurd to distinguish basic sentences based on their content.

## Alternative approach

We want to re-introduce in a controlled manner the ability for multiple and vacuous discharge of hypotheses.

It is not absurd to distinguish basic sentences based on their content.

The idea is to define a consequence relation which allows for simultaneous inference from both sentences which are "persistent" and "consumable", thus extending the definition of an basic rule to allow derivations to occur from both collections of sentences.

Thus our basic rule might look something like the following:




$$\begin{array}{ccc}
 [P_1; C_1] & & [P_n; C_n] \\
 \vdots & & \vdots \\
 q_1 & \dots & q_n
 \end{array}
 \frac{}{r} \mathcal{R}$$

# Thank you!

Thank you for listening!




Questions? Comments? Observations? Please ask and/or feel free to email me at [y.buzoku@ucl.ac.uk](mailto:y.buzoku@ucl.ac.uk).

# References I

-  Alexander V. Gheorghiu, Tao Gu, and David J. Pym, *Proof-theoretic semantics for intuitionistic multiplicative linear logic*, Automated Reasoning with Analytic Tableaux and Related Methods (Cham) (Revantha Ramanayake and Josef Urban, eds.), Springer Nature Switzerland, 2023, pp. 367–385.
-  J.-Y. Girard, *Linear logic: its syntax and semantics*, London Mathematical Society Lecture Note Series, p. 1–42, Cambridge University Press, 1995.
-  Alexander V. Gheorghiu and David J. Pym, *From proof-theoretic validity to base-extension semantics for intuitionistic propositional logic*, 2022.



## References II

-  Sara Negri, *A normalizing system of natural deduction for intuitionistic linear logic*, *Archive for Mathematical Logic* **41** (2002), no. 8, 789–810.
-  Tor Sandqvist, *An inferentialist interpretation of classical logic*, Ph.D. thesis, Uppsala universitet, 2005.
-  ———, *Base-extension semantics for intuitionistic sentential logic*, *Log. J. IGPL* **23** (2015), 719–731.