

# Translations between bases in Base-extension Semantics

Yll Buzoku

Department of Computer Science  
University College London

December 11, 2024

# Goals for this talk

- Introduce a more generalised framework of atomic rules and bases.
- Introduce two notions of atomic derivability.
- Show how one may relate these notions of atomic derivability.
- Recall support relations for IPL and ILL.
- Show how one may thus relate these notions of support.

# Presentation root directory

- 1 Bases and atomic rules
- 2 Atomic derivability
- 3 Comparing relations
- 4 BeS for IPL and ILL
- 5 Translating between the semantics

# Notation

- $\mathbb{A}$  represents a fixed, countably infinite set of propositional atoms.
- Lower case latin letters represent propositional atoms.
- Upper case latin letters represent finite multisets of propositional atoms.
- The sum of two multisets  $P$  and  $Q$  is denoted  $P \uplus Q$ .

# Atomic rules

Atomic rules take the form:

$$(P_1 \Rightarrow p_1), \dots, (P_n \Rightarrow p_n) \Rightarrow p$$

# Atomic rules

Pictorially, we can represent this as:

$$\frac{\begin{array}{ccc} [P_1] & & [P_n] \\ p_1 & \dots & p_n \end{array}}{p}$$

# Atomic rules

Pictorially, we can represent this as:

$$\frac{\begin{array}{ccc} [P_1] & & [P_n] \\ p_1 & \dots & p_n \end{array}}{p}$$

or linearly, like this:

$$(P_1 \Rightarrow p_1), \dots, (P_n \Rightarrow p_n) \Rightarrow p$$

# Contextual atomic rules

Contextual atomic rules are rules which may have contextual brackets distributed across them.



## Contextual atomic rules

Pictorially, we can represent such rules as:

$$\frac{\left\{ \begin{array}{ccc} [P_{1_1}] & & [P_{1_{l_1}}] \\ p_{1_1} & \dots & p_{1_{l_1}} \end{array} \right\} \quad \dots \quad \left\{ \begin{array}{ccc} [P_{n_1}] & & [P_{n_{l_n}}] \\ p_{n_1} & \dots & p_{n_{l_n}} \end{array} \right\}}{p}$$

# Contextual atomic rules

Pictorially, we can represent such rules as:

$$\frac{\left\{ \begin{array}{ccc} [P_{1_1}] & & [P_{1_{l_1}}] \\ p_{1_1} & \dots & p_{1_{l_1}} \end{array} \right\} \quad \dots \quad \left\{ \begin{array}{ccc} [P_{n_1}] & & [P_{n_{l_n}}] \\ p_{n_1} & \dots & p_{n_{l_n}} \end{array} \right\}}{p}$$

or linearly, like this:

$$\{(P_{1_i} \Rightarrow p_{1_i})\}_{i=1}^{l_1}, \dots, \{(P_{n_i} \Rightarrow p_{n_i})\}_{i=1}^{l_n} \Rightarrow p$$

## Examples of atomic rules

- $\{\Rightarrow a\} \Rightarrow c$
- $\{\Rightarrow a\}, \{(c, d \Rightarrow e), (f \Rightarrow g)\} \Rightarrow q$
- $\{a \Rightarrow b\}, \{\Rightarrow e\}, (f \Rightarrow g) \Rightarrow q$
  
- $(\Rightarrow a) \Rightarrow c$
- $(\Rightarrow b) \Rightarrow c$
- $(\Rightarrow c), (a \Rightarrow p), (b \Rightarrow p) \Rightarrow p$

# Bases

## Definition (Base)

A base is a set of atomic rules.

# Bases

## Definition (Base)

A base is a set of atomic rules.

A base is said to be contextual if it contains contextual rules.

Else it is said to be context-free.

# Bases

## Definition (Base)

A base is a set of atomic rules.

A base is said to be contextual if it contains contextual rules.

Else it is said to be context-free.

- $\{((\Rightarrow a) \Rightarrow c), ((\Rightarrow b) \Rightarrow c), ((\Rightarrow c), (a \Rightarrow p), (b \Rightarrow p) \Rightarrow p)\}$
- $\{(\{ \Rightarrow a \} \Rightarrow c), (\{ a \Rightarrow c \} \Rightarrow b), (\{ \Rightarrow a \}, \{ \Rightarrow a \} \Rightarrow a)\}$
- $\{(\{ \Rightarrow a \} \Rightarrow c), (\{ a \Rightarrow c \} \Rightarrow b), (\{ \Rightarrow d \}, \{ \Rightarrow d \} \Rightarrow a), (\{ \Rightarrow a \} \Rightarrow d)\}$

# Presentation root directory

- 1 Bases and atomic rules
- 2 Atomic derivability**
- 3 Comparing relations
- 4 BeS for IPL and ILL
- 5 Translating between the semantics

## Definition (Context-free atomic derivability $\triangleright_{\mathcal{B}}$ )

The relation of derivability in a context-free base  $\mathcal{B}$ , is defined as so:

**Ref**  $S \triangleright_{\mathcal{B}} p$  if  $p \in S$ .

**App** For  $((P_1 \Rightarrow p_1), \dots, (P_n \Rightarrow p_n) \Rightarrow q) \in \mathcal{B}$  and  $S, P_i \triangleright_{\mathcal{B}} p_i$  for each  $i \in \{1, \dots, n\}$  then  $S \triangleright_{\mathcal{B}} q$ .



## Definition (Context-free atomic derivability $\triangleright_{\mathcal{B}}$ )

The relation of derivability in a context-free base  $\mathcal{B}$ , is defined as so:

**Ref**  $S \triangleright_{\mathcal{B}} p$  if  $p \in S$ .

**App** For  $((P_1 \Rightarrow p_1), \dots, (P_n \Rightarrow p_n) \Rightarrow q) \in \mathcal{B}$  and  $S, P_i \triangleright_{\mathcal{B}} p_i$  for each  $i \in \{1, \dots, n\}$  then  $S \triangleright_{\mathcal{B}} q$ .

## Definition (Contextual atomic derivability $\vdash_{\mathcal{B}}$ )

The relation of derivability in a contextual base  $\mathcal{B}$ , is defined as so:

**Ref**  $p \vdash_{\mathcal{B}} p$ .

**App** For  $(\{(P_{1_i} \Rightarrow p_{1_i})\}_{i=1}^{l_1}, \dots, \{(P_{n_i} \Rightarrow p_{n_i})\}_{i=1}^{l_n} \Rightarrow q) \in \mathcal{B}$  and  $C_i, P_{ij} \vdash_{\mathcal{B}} p_{ij}$  for each  $i \in \{1, \dots, n\}$  and  $j \in \{1, \dots, l_i\}$  then  $C_1, \dots, C_n \vdash_{\mathcal{B}} q$ .

# Example derivations

## Example

Let  $\mathcal{B} = \{(\Rightarrow a), ((\Rightarrow a), (\Rightarrow b) \Rightarrow c)\}$

$$\frac{\frac{}{b \triangleright_{\mathcal{B}} b} \text{Ref} \quad \frac{}{b \triangleright_{\mathcal{B}} a} \Rightarrow a}{b \triangleright_{\mathcal{B}} c} (\Rightarrow a), (\Rightarrow b) \Rightarrow c$$

# Example derivations

## Example

Let  $\mathcal{B} = \{(\Rightarrow a), (\{\Rightarrow a\}, \{\Rightarrow b\} \Rightarrow c)\}$

$$\frac{\frac{}{b \vdash_{\mathcal{B}} b} \text{Ref} \quad \frac{}{\vdash_{\mathcal{B}} a} \Rightarrow a}{b \vdash_{\mathcal{B}} c} \{\Rightarrow a\}, \{\Rightarrow b\} \Rightarrow c$$

## Example derivations

### Example

By the definition of  $\vdash_{\mathcal{B}}$ , deriving  $a$  from  $a, a$  in the empty base is not possible, i.e.  $a, a \vdash_{\emptyset} a$  is not possible. But  $a, a \triangleright_{\emptyset} a$  is possible.

## Example derivations

### Example

By the definition of  $\vdash_{\mathcal{B}}$ , deriving  $a$  from  $a, a$  in the empty base is not possible, i.e.  $a, a \vdash_{\emptyset} a$  is not possible. But  $a, a \triangleright_{\emptyset} a$  is possible.

### Example

Let  $\mathcal{B} = \{(\{ \Rightarrow a \} \Rightarrow c), (\{ a \Rightarrow c \} \Rightarrow b)\}$

$$\frac{\frac{\frac{\times}{a, a \vdash_{\mathcal{B}} a} \quad \{ \Rightarrow a \} \Rightarrow c}{a, a \vdash_{\mathcal{B}} c} \quad \{ a \Rightarrow c \} \Rightarrow b}{a \vdash_{\mathcal{B}} b}$$

We see that in this base,  $a$  is not derivable from  $a, a$ .

# Example derivations

## Example (A possible fix)

Let  $(\mathcal{B})_\star = \{((\Rightarrow a) \Rightarrow c), ((a \Rightarrow c) \Rightarrow b)\}$

$$\begin{array}{c}
 \frac{}{a, a \triangleright_{(\mathcal{B})_\star} a} \text{Ref} \\
 \frac{}{a, a \triangleright_{(\mathcal{B})_\star} c} (\Rightarrow a) \Rightarrow c \\
 \frac{}{a \triangleright_{(\mathcal{B})_\star} b} (a \Rightarrow c) \Rightarrow b
 \end{array}$$

# Example derivations

## Example (Another possible fix)

Let  $\mathcal{C} = \{(\{\Rightarrow a\} \Rightarrow c), (\{a \Rightarrow c\} \Rightarrow b), (\{\Rightarrow a\}, \{\Rightarrow a\} \Rightarrow a)\}$

$$\begin{array}{c}
 \frac{}{a \vdash_{\mathcal{C}} a} \text{Ref} \quad \frac{}{a \vdash_{\mathcal{C}} a} \text{Ref} \\
 \hline
 \frac{}{a, a \vdash_{\mathcal{C}} a} \{\Rightarrow a\}, \{\Rightarrow a\} \Rightarrow a \\
 \hline
 \frac{}{a, a \vdash_{\mathcal{C}} c} \{\Rightarrow a\} \Rightarrow c \\
 \hline
 \frac{}{a \vdash_{\mathcal{C}} b} \{a \Rightarrow c\} \Rightarrow b
 \end{array}$$

# Example derivations

## Example (A very interesting derivation)

Let  $\mathcal{B}$  be the following set of rules:

$\{(\{ \Rightarrow b \} \Rightarrow c), (\{ \Rightarrow a \}, \{ b \Rightarrow c \} \Rightarrow d), (\{ \Rightarrow d, \Rightarrow a \} \Rightarrow e), (\{ a \Rightarrow e \} \Rightarrow f)\}$

$$\begin{array}{c}
 \frac{}{a \vdash_{\mathcal{B}} a} \text{Ref} \quad \frac{\frac{}{b \vdash_{\mathcal{B}} b} \text{Ref}}{b \vdash_{\mathcal{B}} c} \{ \Rightarrow b \} \Rightarrow c}{\frac{}{a \vdash_{\mathcal{B}} d} \{ \Rightarrow a \}, \{ b \Rightarrow c \} \Rightarrow d} \text{Ref} \quad \frac{}{a \vdash_{\mathcal{B}} a} \text{Ref} \\
 \frac{}{a \vdash_{\mathcal{B}} d} \{ \Rightarrow d, \Rightarrow a \} \Rightarrow e \quad \frac{a \vdash_{\mathcal{B}} e}{\vdash_{\mathcal{B}} f} \{ a \Rightarrow e \} \Rightarrow f
 \end{array}$$



# Presentation root directory

- 1 Bases and atomic rules
- 2 Atomic derivability
- 3 Comparing relations**
- 4 BeS for IPL and ILL
- 5 Translating between the semantics

# Comparing relations

## Definition (Structural rules)

We define two rules:

- $\text{Wk}_q^p = (\{\Rightarrow p\}, \{\Rightarrow q\} \Rightarrow q)$
- $\text{Ctn}_q^p = (\{\Rightarrow p\}, \{p, p \Rightarrow q\} \Rightarrow q)$

# Mappings between bases

# Mappings between bases

## Definition

Let  $\mathcal{B}$  be a context-free base. We define structural contextualisation of that base  $(\mathcal{B})^*$  as follows:

$$(\mathcal{B})^* = \{ \{P_1 \Rightarrow p_1\}, \dots, \{P_n \Rightarrow p_n\} \Rightarrow q \mid ((P_1 \Rightarrow p_1), \dots, (P_n \Rightarrow p_n) \Rightarrow q) \in \mathcal{B} \} \\ \cup \{ \text{Wk}_q^p, \text{Ctn}_q^p \mid \forall p, q \in \mathbb{A} \}$$

# Mappings between bases

## Definition

Let  $\mathcal{B}$  be a context-free base. We define structural contextualisation of that base  $(\mathcal{B})^*$  as follows:

$$(\mathcal{B})^* = \{ \{P_1 \Rightarrow p_1\}, \dots, \{P_n \Rightarrow p_n\} \Rightarrow q \mid ((P_1 \Rightarrow p_1), \dots, (P_n \Rightarrow p_n) \Rightarrow q) \in \mathcal{B} \} \\ \cup \{ \text{Wk}_q^p, \text{Ctn}_q^p \mid \forall p, q \in \mathbb{A} \}$$

We define the decontextualisation of a contextual base as:

$$(\mathcal{B})_\star = \{ (P_{1_1} \Rightarrow p_{1_1}), \dots, (P_{n_{l_n}} \Rightarrow p_{n_{l_n}}) \Rightarrow q \\ \mid (\{P_{1_i} \Rightarrow p_{1_i}\}_{i=1}^{l_1}, \dots, \{P_{n_i} \Rightarrow p_{n_i}\}_{i=1}^{l_n} \Rightarrow q) \in \mathcal{B} \}$$

# Properties of these mappings

Let  $\mathcal{B}$  be a context-free base. Then the following hold:

- $((\mathcal{B})^*)_{\star} \supseteq \mathcal{B}$

# Properties of these mappings

Let  $\mathcal{B}$  be a context-free base. Then the following hold:

- $((\mathcal{B})^*)_{\star} \supseteq \mathcal{B}$
- $((\mathcal{B})^*)_{\star}^* = (\mathcal{B})^*$

# Properties of these mappings

Let  $\mathcal{B}$  be a context-free base. Then the following hold:

- $((\mathcal{B})^*)_{\star} \supseteq \mathcal{B}$
- $((\mathcal{B})^*)_{\star}^* = (\mathcal{B})^*$
- For all  $\mathcal{C} \supseteq \mathcal{B}$  we have that  $(\mathcal{C})^* \supseteq (\mathcal{B})^*$



# Properties of these mappings

Let  $\mathcal{B}$  be a context-free base. Then the following hold:

- $((\mathcal{B})^*)_{\star} \supseteq \mathcal{B}$
- $((\mathcal{B})^*)_{\star}^* = (\mathcal{B})^*$
- For all  $\mathcal{C} \supseteq \mathcal{B}$  we have that  $(\mathcal{C})^* \supseteq (\mathcal{B})^*$
- For all  $\mathcal{C} \supseteq (\mathcal{B})^*$  there exists an extension  $\mathcal{X} \supseteq \mathcal{B}$  such that  $(\mathcal{X})^* = \mathcal{C}$

# Properties of these mappings

Let  $\mathcal{B}$  be a context-free base. Then the following hold:

- $((\mathcal{B})^*)_{\star} \supseteq \mathcal{B}$
- $((\mathcal{B})^*)_{\star}^* = (\mathcal{B})^*$
- For all  $\mathcal{C} \supseteq \mathcal{B}$  we have that  $(\mathcal{C})^* \supseteq (\mathcal{B})^*$
- For all  $\mathcal{C} \supseteq (\mathcal{B})^*$  there exists an extension  $\mathcal{X} \supseteq \mathcal{B}$  such that  $(\mathcal{X})^* = \mathcal{C}$
- For all  $\mathcal{C} \supseteq (\mathcal{B})^*$ , it follows immediately that  $\mathcal{C} = ((\mathcal{C})_{\star})^*$ .

# Structural admissibility in a structurally contextualised base

## Lemma

*Suppose  $L \vdash_{(\mathcal{B})^*} p$  holds. Then  $R, L \vdash_{(\mathcal{B})^*} p$  also holds for any atomic multiset  $R$ .*

Proof.

Let  $R = \{r_1, \dots, r_n\}$  for some  $n$ . Then we have that we can effectively weaken  $S$  away as follows:

$$\begin{array}{c}
 \frac{\frac{\frac{}{r_1 \vdash_{(\mathcal{B})^*} r_1} \text{Ref}}{r_{n-1} \vdash_{(\mathcal{B})^*} r_{n-1}} \text{Ref} \quad \frac{\frac{\frac{}{r_n \vdash_{(\mathcal{B})^*} r_n} \text{Ref} \quad L \vdash_{(\mathcal{B})^*} p}{r_n, L \vdash_{(\mathcal{B})^*} p} \text{Wk}_p^{r_n}}{r_n, L \vdash_{(\mathcal{B})^*} p} \text{Wk}_p^{r_{n-1}}}{\vdots} \\
 \frac{\frac{\frac{}{r_1 \vdash_{(\mathcal{B})^*} r_1} \text{Ref} \quad r_2, \dots, r_n, L \vdash_{(\mathcal{B})^*} p}{R, L \vdash_{(\mathcal{B})^*} p} \text{Wk}_p^{r_1}}{}
 \end{array}$$

□

# Structural admissibility in a structurally contextualised base

## Lemma

*Suppose  $R, R \vdash_{(\mathcal{B})^*} p$  holds. Then  $R \vdash_{(\mathcal{B})^*} p$  also holds.*

# Structural admissibility in a structurally contextualised base

## Lemma

*Suppose  $R, R \vdash_{(\mathcal{B})^*} p$  holds. Then  $R \vdash_{(\mathcal{B})^*} p$  also holds.*

## Corollary

*For arbitrary  $m \geq 1$ , if  $R^m \vdash_{(\mathcal{B})^*} p$  then  $R \vdash_{(\mathcal{B})^*} p$ .*

Proof.

Let  $R = \{r_1, \dots, r_n\}$  for some  $n$ . Then we have that we can effectively contract on  $R$  as follows:

$$\begin{array}{c}
 \frac{\frac{\frac{}{r_1 \vdash_{(\mathcal{B})^*} r_1} \text{Ref} \quad \vdots \quad \frac{\frac{\frac{}{r_n \vdash_{(\mathcal{B})^*} r_n} \text{Ref} \quad \frac{R, R \vdash_{(\mathcal{B})^*} p}{r_1, \dots, r_{n-1}, R \vdash_{(\mathcal{B})^*} p} \text{Ctn}_p^{r_n}}{r_1, \dots, r_{n-1}, R \vdash_{(\mathcal{B})^*} p} \text{Ctn}_p^{r_{n-1}}}{R \vdash_{(\mathcal{B})^*} p} \text{Ctn}_p^{r_1}
 \end{array}
 \quad \square$$

## Key results under this base translation

- If  $R \triangleright_{\mathcal{B}} p$  then  $R \vdash_{(\mathcal{B})^*} p$ .



## Key results under this base translation

- If  $R \triangleright_{\mathcal{B}} p$  then  $R \vdash_{(\mathcal{B})^*} p$ .
- $R \vdash_{(\mathcal{B})^*} p$  iff for all bases  $\mathcal{X} \supseteq (\mathcal{B})^*$  where for each  $r \in R$  we have  $\vdash_{\mathcal{X}} r$  then it follows that  $\vdash_{\mathcal{X}} p$ .

## Key results under this base translation

- If  $R \triangleright_{\mathcal{B}} p$  then  $R \vdash_{(\mathcal{B})^*} p$ .
- $R \vdash_{(\mathcal{B})^*} p$  iff for all bases  $\mathcal{X} \supseteq (\mathcal{B})^*$  where for each  $r \in R$  we have  $\vdash_{\mathcal{X}} r$  then it follows that  $\vdash_{\mathcal{X}} p$ .
- $R \vdash_{(\mathcal{B})^*} p$  iff  $!R \Vdash_{(\mathcal{B})^*}^{\emptyset} p$

## Key results under this base translation

- If  $R \triangleright_{\mathcal{B}} p$  then  $R \vdash_{(\mathcal{B})^*} p$ .
- $R \vdash_{(\mathcal{B})^*} p$  iff for all bases  $\mathcal{X} \supseteq (\mathcal{B})^*$  where for each  $r \in R$  we have  $\vdash_{\mathcal{X}} r$  then it follows that  $\vdash_{\mathcal{X}} p$ .
- $R \vdash_{(\mathcal{B})^*} p$  iff  $!R \Vdash_{(\mathcal{B})^*}^{\emptyset} p$
- $\Vdash_{(\mathcal{B})^*}^R p$  iff  $!R \Vdash_{(\mathcal{B})^*}^{\emptyset} p$

# Presentation root directory

- 1 Bases and atomic rules
- 2 Atomic derivability
- 3 Comparing relations
- 4 BeS for IPL and ILL**
- 5 Translating between the semantics

## BeS for IPL: Summary

- $\Vdash_{\mathcal{B}} p$  iff  $\triangleright_{\mathcal{B}} p$ .
- $\Vdash_{\mathcal{B}} \varphi \wedge \psi$  iff  $\Vdash_{\mathcal{B}} \varphi$  and  $\Vdash_{\mathcal{B}} \psi$ .
- $\Vdash_{\mathcal{B}} \varphi \vee \psi$  iff for all  $\mathcal{C} \supseteq \mathcal{B}$ ,  $p$ , if  $\varphi \Vdash_{\mathcal{C}} p$  and  $\psi \Vdash_{\mathcal{C}} p$  then  $\Vdash_{\mathcal{C}} p$ .
- $\Vdash_{\mathcal{B}} \varphi \rightarrow \psi$  iff  $\varphi \Vdash_{\mathcal{B}} \psi$ .
- $\Vdash_{\mathcal{B}} \perp$  iff  $\Vdash_{\mathcal{B}} p$  for all  $p$ .
- $\Vdash_{\mathcal{B}} \top$  iff always.
- $\Gamma \Vdash_{\mathcal{B}} \varphi$  iff for all  $\mathcal{C} \supseteq \mathcal{B}$  if  $\Vdash_{\mathcal{C}} \gamma$  for all  $\gamma \in \Gamma$  then  $\Vdash_{\mathcal{C}} \varphi$ .
- $\Gamma \Vdash \varphi$  iff  $\Gamma \Vdash_{\mathcal{B}} \varphi$  for all  $\mathcal{B}$ .

# BeS for ILL: Structure and the Exponential

- $\Vdash_{\mathcal{B}}^L p$  iff  $L \vdash_{\mathcal{B}} p$ .
- $\Vdash_{\mathcal{B}}^L !\varphi$  iff  
 $\forall \mathcal{C} \supseteq \mathcal{B}, K, p$ , if  $(\forall \mathcal{D} \supseteq \mathcal{C} (\Vdash_{\mathcal{D}}^{\emptyset} \varphi \text{ implies } \Vdash_{\mathcal{D}}^K p))$  then  $\Vdash_{\mathcal{C}}^{L,K} p$ .
- $\Gamma \Vdash_{\mathcal{B}}^L \varphi$  iff  
 $\forall \mathcal{C} \supseteq \mathcal{B}, K$ , if  $\Vdash_{\mathcal{C}}^{\emptyset} \Delta$  and  $\Vdash_{\mathcal{C}}^K \Theta$  where  $!\Delta, \Theta = \Gamma$  then  $\Vdash_{\mathcal{C}}^{L,K} \varphi$ .
- $\Gamma \Vdash \varphi$  iff  $\Gamma \Vdash_{\mathcal{B}}^{\emptyset} \varphi$  for all  $\mathcal{B}$ .

## BeS for ILL: Multiplicatives

- $\Vdash_{\mathcal{B}}^L \varphi \otimes \psi$  iff for all  $\mathcal{C} \supseteq \mathcal{B}$ ,  $K$  and  $p$ , if  $\varphi, \psi \Vdash_{\mathcal{C}}^K p$  then  $\Vdash_{\mathcal{C}}^{L,K} p$ .
- $\Vdash_{\mathcal{B}}^L \varphi \multimap \psi$  iff  $\varphi \Vdash_{\mathcal{B}}^L \psi$ .
- $\Vdash_{\mathcal{B}}^L 1$  iff for all  $\mathcal{C} \supseteq \mathcal{B}$ ,  $K$  and  $p$ , if  $\Vdash_{\mathcal{C}}^K p$  then  $\Vdash_{\mathcal{C}}^{L,K} p$ .

## BeS for ILL: Additives

- $\Vdash_{\mathcal{B}}^L \varphi \& \psi$  iff  $\Vdash_{\mathcal{B}}^L \varphi$  and  $\Vdash_{\mathcal{B}}^L \psi$ .
- $\Vdash_{\mathcal{B}}^L \varphi \oplus \psi$  iff for all  $\mathcal{C} \supseteq \mathcal{B}$ ,  $K$  and  $p$ , if  $\varphi \Vdash_{\mathcal{C}}^K p$  and  $\psi \Vdash_{\mathcal{C}}^K p$  then  $\Vdash_{\mathcal{C}}^{L,K} p$ .
- $\Vdash_{\mathcal{B}}^L 0$  iff  $\Vdash_{\mathcal{B}}^L p$  for all  $p$ .
- $\Vdash_{\mathcal{B}}^L \top$  iff always.



## BeS for ILL: Summary

- $\Vdash_{\mathcal{B}}^L p$  iff  $L \vdash_{\mathcal{B}} p$ .
- $\Vdash_{\mathcal{B}}^L \varphi \otimes \psi$  iff  $\forall \mathcal{C} \supseteq \mathcal{B}, K, p (\varphi, \psi \Vdash_{\mathcal{C}}^K p \text{ implies } \Vdash_{\mathcal{C}}^{L,K} p)$ .
- $\Vdash_{\mathcal{B}}^L \varphi \multimap \psi$  iff  $\varphi \Vdash_{\mathcal{B}}^L \psi$ .
- $\Vdash_{\mathcal{B}}^L 1$  iff  $\forall \mathcal{C} \supseteq \mathcal{B}, K, p (\Vdash_{\mathcal{C}}^K p \text{ implies } \Vdash_{\mathcal{C}}^{L,K} p)$ .
- $\Vdash_{\mathcal{B}}^L \varphi \& \psi$  iff  $\Vdash_{\mathcal{B}}^L \varphi$  and  $\Vdash_{\mathcal{B}}^L \psi$ .
- $\Vdash_{\mathcal{B}}^L \varphi \oplus \psi$  iff  $\forall \mathcal{C} \supseteq \mathcal{B}, K, p (\varphi \Vdash_{\mathcal{C}}^K p \text{ and } \psi \Vdash_{\mathcal{C}}^K p \text{ implies } \Vdash_{\mathcal{C}}^{L,K} p)$ .
- $\Vdash_{\mathcal{B}}^L 0$  iff  $\Vdash_{\mathcal{B}}^L p$  for all  $p$ .
- $\Vdash_{\mathcal{B}}^L \top$  iff always.
- $\Vdash_{\mathcal{B}}^L !\varphi$  iff  
 $\forall \mathcal{C} \supseteq \mathcal{B}, K, p (\forall \mathcal{D} \supseteq \mathcal{C} (\Vdash_{\mathcal{D}}^{\emptyset} \varphi \text{ implies } \Vdash_{\mathcal{D}}^K p) \text{ implies } \Vdash_{\mathcal{C}}^{L,K} p)$ .
- $\Gamma \Vdash_{\mathcal{B}}^L \varphi$  iff  
 $\forall \mathcal{C} \supseteq \mathcal{B}, K, \Vdash_{\mathcal{C}}^{\emptyset} \delta \text{ and } \Vdash_{\mathcal{C}}^K \gamma \text{ for } !\delta \in \Gamma \text{ and } \gamma \in \Gamma \text{ then } \Vdash_{\mathcal{C}}^{L,K} \varphi$ .
- $\Gamma \vdash \varphi$  iff  $\Gamma \Vdash_{\mathcal{B}}^{\emptyset} \varphi$  for all  $\mathcal{B}$ .

# Presentation root directory

- 1 Bases and atomic rules
- 2 Atomic derivability
- 3 Comparing relations
- 4 BeS for IPL and ILL
- 5 Translating between the semantics**

## Comparing clauses

(IPL)  $\Vdash_{\mathcal{B}} p$  iff  $\triangleright_{\mathcal{B}} p$ .

(ILL)  $\Vdash_{\mathcal{B}}^L p$  iff  $L \vdash_{\mathcal{B}} p$ .

## Comparing clauses

(IPL)  $\Vdash_{\mathcal{B}} p$  iff  $\triangleright_{\mathcal{B}} p$ .

(ILL)  $\Vdash_{\mathcal{B}}^L p$  iff  $L \vdash_{\mathcal{B}} p$ .

---

(IPL)  $\Vdash_{\mathcal{B}} \varphi \wedge \psi$  iff  $\Vdash_{\mathcal{B}} \varphi$  and  $\Vdash_{\mathcal{B}} \psi$ .

(ILL)  $\Vdash_{\mathcal{B}}^L \varphi \& \psi$  iff  $\Vdash_{\mathcal{B}}^L \varphi$  and  $\Vdash_{\mathcal{B}}^L \psi$ .

## Comparing clauses

(IPL)  $\Vdash_{\mathcal{B}} p$  iff  $\triangleright_{\mathcal{B}} p$ .

(ILL)  $\Vdash_{\mathcal{B}}^L p$  iff  $L \vdash_{\mathcal{B}} p$ .

---

(IPL)  $\Vdash_{\mathcal{B}} \varphi \wedge \psi$  iff  $\Vdash_{\mathcal{B}} \varphi$  and  $\Vdash_{\mathcal{B}} \psi$ .

(ILL)  $\Vdash_{\mathcal{B}}^L \varphi \& \psi$  iff  $\Vdash_{\mathcal{B}}^L \varphi$  and  $\Vdash_{\mathcal{B}}^L \psi$ .

---

(IPL)  $\Vdash_{\mathcal{B}} \varphi \vee \psi$  iff for all  $\mathcal{C} \supseteq \mathcal{B}$ ,  $p$ , if  $\varphi \Vdash_{\mathcal{C}} p$  and  $\psi \Vdash_{\mathcal{C}} p$  then  $\Vdash_{\mathcal{C}} p$ .

(ILL)  $\Vdash_{\mathcal{B}}^L \varphi \oplus \psi$  iff for all  $\mathcal{C} \supseteq \mathcal{B}$ ,  $K$ ,  $p$ , if  $\varphi \Vdash_{\mathcal{C}}^K p$  and  $\psi \Vdash_{\mathcal{C}}^K p$  then  $\Vdash_{\mathcal{C}}^{L,K} p$ .

## Comparing clauses

(IPL)  $\Vdash_{\mathcal{B}} p$  iff  $\triangleright_{\mathcal{B}} p$ .

(ILL)  $\Vdash_{\mathcal{B}}^L p$  iff  $L \vdash_{\mathcal{B}} p$ .

---

(IPL)  $\Vdash_{\mathcal{B}} \varphi \wedge \psi$  iff  $\Vdash_{\mathcal{B}} \varphi$  and  $\Vdash_{\mathcal{B}} \psi$ .

(ILL)  $\Vdash_{\mathcal{B}}^L \varphi \& \psi$  iff  $\Vdash_{\mathcal{B}}^L \varphi$  and  $\Vdash_{\mathcal{B}}^L \psi$ .

---

(IPL)  $\Vdash_{\mathcal{B}} \varphi \vee \psi$  iff for all  $\mathcal{C} \supseteq \mathcal{B}$ ,  $p$ , if  $\varphi \Vdash_{\mathcal{C}} p$  and  $\psi \Vdash_{\mathcal{C}} p$  then  $\Vdash_{\mathcal{C}} p$ .

(ILL)  $\Vdash_{\mathcal{B}}^L \varphi \oplus \psi$  iff for all  $\mathcal{C} \supseteq \mathcal{B}$ ,  $K$ ,  $p$ , if  $\varphi \Vdash_{\mathcal{C}}^K p$  and  $\psi \Vdash_{\mathcal{C}}^K p$  then  $\Vdash_{\mathcal{C}}^{L,K} p$ .

---

(IPL)  $\Vdash_{\mathcal{B}} \varphi \rightarrow \psi$  iff  $\varphi \Vdash_{\mathcal{B}} \psi$ .

(ILL)  $\Vdash_{\mathcal{B}}^L \varphi \multimap \psi$  iff  $\varphi \Vdash_{\mathcal{B}}^L \psi$ .

## Comparing clauses

(IPL)  $\Vdash_{\mathcal{B}} p \text{ iff } \triangleright_{\mathcal{B}} p.$

(ILL)  $\Vdash_{\mathcal{B}}^L p \text{ iff } L \vdash_{\mathcal{B}} p.$

---

(IPL)  $\Vdash_{\mathcal{B}} \varphi \wedge \psi \text{ iff } \Vdash_{\mathcal{B}} \varphi \text{ and } \Vdash_{\mathcal{B}} \psi.$

(ILL)  $\Vdash_{\mathcal{B}}^L \varphi \& \psi \text{ iff } \Vdash_{\mathcal{B}}^L \varphi \text{ and } \Vdash_{\mathcal{B}}^L \psi.$

---

(IPL)  $\Vdash_{\mathcal{B}} \varphi \vee \psi \text{ iff for all } \mathcal{C} \supseteq \mathcal{B}, p, \text{ if } \varphi \Vdash_{\mathcal{C}} p \text{ and } \psi \Vdash_{\mathcal{C}} p \text{ then } \Vdash_{\mathcal{C}} p.$

(ILL)  $\Vdash_{\mathcal{B}}^L \varphi \oplus \psi \text{ iff for all } \mathcal{C} \supseteq \mathcal{B}, K, p, \text{ if } \varphi \Vdash_{\mathcal{C}}^K p \text{ and } \psi \Vdash_{\mathcal{C}}^K p \text{ then } \Vdash_{\mathcal{C}}^{L,K} p.$

---

(IPL)  $\Vdash_{\mathcal{B}} \varphi \rightarrow \psi \text{ iff } \varphi \Vdash_{\mathcal{B}} \psi.$

(ILL)  $\Vdash_{\mathcal{B}}^L \varphi \multimap \psi \text{ iff } \varphi \Vdash_{\mathcal{B}}^L \psi.$

---

(IPL)  $\Vdash_{\mathcal{B}} \perp \text{ iff } \Vdash_{\mathcal{B}} p \text{ for all } p.$

(ILL)  $\Vdash_{\mathcal{B}}^L 0 \text{ iff } \Vdash_{\mathcal{B}}^L p \text{ for all } p.$

---

# Translation of formulas

We seem to have a correspondence between the clauses for the following pairs of connectives:

- $\wedge$  and  $\&$
- $\vee$  and  $\oplus$
- $\rightarrow$  and  $\multimap$
- $\perp$  and  $0$



# Translation of formulas

One possible translation of formulas between IPL and ILL is:

- $\llbracket p \rrbracket \equiv p$ , where  $p$  is an atom.
- $\llbracket \varphi \wedge \psi \rrbracket \equiv \llbracket \varphi \rrbracket \& \llbracket \psi \rrbracket$
- $\llbracket \varphi \vee \psi \rrbracket \equiv !\llbracket \varphi \rrbracket \oplus !\llbracket \psi \rrbracket$
- $\llbracket \varphi \rightarrow \psi \rrbracket \equiv !\llbracket \varphi \rrbracket \multimap \llbracket \psi \rrbracket$
- $\llbracket \perp \rrbracket \equiv 0$

## Translation of formulas

(IPL)  $\Vdash_{\mathcal{B}} p \text{ iff } \triangleright_{\mathcal{B}} p.$

(ILL)  $\Vdash_{\mathcal{B}}^L p \text{ iff } L \vdash_{\mathcal{B}} p.$

---

(IPL)  $\Vdash_{\mathcal{B}} \varphi \wedge \psi \text{ iff } \Vdash_{\mathcal{B}} \varphi \text{ and } \Vdash_{\mathcal{B}} \psi.$

(ILL)  $\Vdash_{\mathcal{B}}^L \varphi \& \psi \text{ iff } \Vdash_{\mathcal{B}}^L \varphi \text{ and } \Vdash_{\mathcal{B}}^L \psi.$

---

(IPL)  $\Vdash_{\mathcal{B}} \varphi \vee \psi \text{ iff for all } \mathcal{C} \supseteq \mathcal{B}, p, \text{ if } \varphi \Vdash_{\mathcal{C}} p \text{ and } \psi \Vdash_{\mathcal{C}} p \text{ then } \Vdash_{\mathcal{C}} p.$

(ILL)  $\Vdash_{\mathcal{B}}^L \varphi \oplus \psi \text{ iff for all } \mathcal{C} \supseteq \mathcal{B}, K, p, \text{ if } \varphi \Vdash_{\mathcal{C}}^K p \text{ and } \psi \Vdash_{\mathcal{C}}^K p \text{ then } \Vdash_{\mathcal{C}}^{L,K} p.$

---

(IPL)  $\Vdash_{\mathcal{B}} \varphi \rightarrow \psi \text{ iff } \varphi \Vdash_{\mathcal{B}} \psi.$

(ILL)  $\Vdash_{\mathcal{B}}^L \varphi \multimap \psi \text{ iff } \varphi \Vdash_{\mathcal{B}}^L \psi.$

---

(IPL)  $\Vdash_{\mathcal{B}} \perp \text{ iff } \Vdash_{\mathcal{B}} p \text{ for all } p.$

(ILL)  $\Vdash_{\mathcal{B}}^L 0 \text{ iff } \Vdash_{\mathcal{B}}^L p \text{ for all } p.$

---

# Translation of formulas

One possible translation of formulas between IPL and ILL is:

- $\llbracket p \rrbracket \equiv p$ , where  $p$  is an atom.
- $\llbracket \varphi \wedge \psi \rrbracket \equiv \llbracket \varphi \rrbracket \& \llbracket \psi \rrbracket$
- $\llbracket \varphi \vee \psi \rrbracket \equiv !\llbracket \varphi \rrbracket \oplus !\llbracket \psi \rrbracket$
- $\llbracket \varphi \rightarrow \psi \rrbracket \equiv !\llbracket \varphi \rrbracket \multimap \llbracket \psi \rrbracket$
- $\llbracket \perp \rrbracket \equiv 0$

# Translation of formulas

One possible translation of formulas between IPL and ILL is:

- $\llbracket p \rrbracket \equiv p$ , where  $p$  is an atom.
- $\llbracket \varphi \wedge \psi \rrbracket \equiv \llbracket \varphi \rrbracket \& \llbracket \psi \rrbracket$
- $\llbracket \varphi \vee \psi \rrbracket \equiv !\llbracket \varphi \rrbracket \oplus !\llbracket \psi \rrbracket$
- $\llbracket \varphi \rightarrow \psi \rrbracket \equiv !\llbracket \varphi \rrbracket \multimap \llbracket \psi \rrbracket$
- $\llbracket \perp \rrbracket \equiv 0$

Sequents are translated as follows:

- If  $\Gamma \vdash_{\text{IPL}} \varphi$  then  $!\llbracket \Gamma \rrbracket \vdash_{\text{ILL}} \llbracket \varphi \rrbracket$

# Translation of formulas

One possible translation of formulas between IPL and ILL is:

- $\llbracket p \rrbracket \equiv p$ , where  $p$  is an atom.
- $\llbracket \varphi \wedge \psi \rrbracket \equiv \llbracket \varphi \rrbracket \& \llbracket \psi \rrbracket$
- $\llbracket \varphi \vee \psi \rrbracket \equiv !\llbracket \varphi \rrbracket \oplus !\llbracket \psi \rrbracket$
- $\llbracket \varphi \rightarrow \psi \rrbracket \equiv !\llbracket \varphi \rrbracket \multimap \llbracket \psi \rrbracket$
- $\llbracket \perp \rrbracket \equiv 0$

Sequents are translated as follows:

- If  $\Gamma \vdash_{\text{IPL}} \varphi$  then  $!\llbracket \Gamma \rrbracket \vdash_{\text{ILL}} \llbracket \varphi \rrbracket$

Semantically we should therefore have:

- If  $\Gamma \Vdash \varphi$  then  $!\llbracket \Gamma \rrbracket \Vdash \llbracket \varphi \rrbracket$

# Immediate correspondence between bases

$$\begin{array}{lll}
 \Gamma \vdash_{\text{IPL}} \varphi & \text{iff } \Gamma \Vdash \varphi & \text{iff for all } \mathcal{B}, p, \text{ if } \Vdash_{\mathcal{B}} \gamma \text{ then } \Vdash_{\mathcal{B}} \varphi \\
 ![\Gamma] \vdash_{\text{ILL}} [\varphi] & \text{iff } ![\Gamma] \Vdash [\varphi] & \text{iff for all } \mathcal{C}, p, \text{ if } \Vdash_{\mathcal{C}}^{\emptyset} [\gamma] \text{ then } \Vdash_{\mathcal{C}}^{\emptyset} [\varphi]
 \end{array}$$

# Immediate correspondence between bases

$$\begin{aligned} \Gamma \vdash_{\text{IPL}} \varphi & \quad \text{iff } \Gamma \Vdash \varphi & \quad \text{iff for all } \mathcal{B}, p, \text{ if } \Vdash_{\mathcal{B}} \gamma \text{ then } \Vdash_{\mathcal{B}} \varphi \\ ![\Gamma] \vdash_{\text{ILL}} [\varphi] & \quad \text{iff } ![\Gamma] \Vdash [\varphi] & \quad \text{iff for all } \mathcal{C}, p, \text{ if } \Vdash_{\mathcal{C}}^{\emptyset} [\gamma] \text{ then } \Vdash_{\mathcal{C}}^{\emptyset} [\varphi] \end{aligned}$$

Since we have that  $\Gamma \vdash_{\text{IPL}} \varphi$  implies  $![\Gamma] \vdash_{\text{ILL}} [\varphi]$ , then we have a mapping between bases  $\mathcal{B}$  and bases  $\mathcal{C}$ .

But what about if we have  $\Gamma \Vdash_{\mathcal{B}} \varphi$ ?



But what about if we have  $\Gamma \Vdash_{\mathcal{B}} \varphi$ ?




## Theorem

*If  $\Gamma \Vdash_{\mathcal{B}} \varphi$  then  $![[\Gamma]] \Vdash_{(\mathcal{B})^*}^{\emptyset} [[\varphi]]$*





Thank you!

**Thank you for listening!**  
**Comments? Observations? Please ask and/or feel free to**  
**email me at [y.buzoku@ucl.ac.uk](mailto:y.buzoku@ucl.ac.uk)**

# References I

-  G.M. Bierman, *On intuitionistic linear logic*, Tech. Report UCAM-CL-TR-346, University of Cambridge, Computer Laboratory, August 1994.
-  Yli Buzoku, *A proof-theoretic semantics for intuitionistic linear logic*, 2024.
-  Alexander V. Gheorghiu, Tao Gu, and David J. Pym, *Proof-theoretic semantics for intuitionistic multiplicative linear logic*, Automated Reasoning with Analytic Tableaux and Related Methods (Cham) (Revantha Ramanayake and Josef Urban, eds.), Springer Nature Switzerland, 2023, pp. 367–385.

## References II

-  J.-Y. Girard, *Linear logic: its syntax and semantics*, London Mathematical Society Lecture Note Series, p. 1–42, Cambridge University Press, 1995.
-  Alexander V. Gheorghiu and David J. Pym, *From proof-theoretic validity to base-extension semantics for intuitionistic propositional logic*, 2022.
-  Sara Negri, *A normalizing system of natural deduction for intuitionistic linear logic*, *Archive for Mathematical Logic* **41** (2002), no. 8, 789–810.
-  Tor Sandqvist, *An inferentialist interpretation of classical logic*, Ph.D. thesis, Uppsala universitet, 2005.

## References III



\_\_\_\_\_, *Base-extension semantics for intuitionistic sentential logic*,  
Log. J. IGPL **23** (2015), 719–731.