

Translations between bases in Base-extension Semantics

Yll Buzoku

Department of Computer Science
University College London

November 22, 2024

Goals for this talk

- Introduce the concept of bases and general inference figures
- Introduce two notions of atomic derivability
- Show how one may relate these notions of atomic derivability

Presentation root directory

- 1 Bases and inference figures
- 2 Atomic derivability
- 3 Comparing relations

Presentation root directory

1 Bases and inference figures

2 Atomic derivability

3 Comparing relations

Notation

- \mathbb{A} represents a fixed, countably infinite set of propositional atoms.
- Lower case latin letters represent propositional atoms.
- Upper case latin letters represent finite multisets of propositional atoms.
- Atomic multiset is taken to mean multiset of propositional atoms.
- The sum of two multisets P and Q is denoted $P \uplus Q$.

Atomic rules

Atomic rules take the form:

$$\langle \langle P_1, p_1 \rangle, \dots, \langle P_n, p_n \rangle, p \rangle$$

Atomic rules

Pictorially, we can represent this as:

$$\frac{[P_1] \quad \dots \quad [P_n]}{p}$$

Atomic rules

Pictorially, we can represent this as:

$$\frac{[P_1] \quad \dots \quad [P_n]}{p}$$

Any idea what these figures are supposed to look like?

Atomic rules

Pictorially, we can represent this as:

$$\frac{\begin{array}{ccc} [P_1] & & [P_n] \\ p_1 & \dots & p_n \end{array}}{p}$$

Any idea what these figures are supposed to look like?

Natural deduction!

Contextual atomic rules

Contextual atomic rules are rules which may have contextual brackets distributed across them.

Contextual atomic rules

Pictorially, we can represent this as:

$$\frac{\left\{ \begin{array}{ccc} [P_{1_1}] & & [P_{1_{l_n}}] \\ p_{1_1} & \dots & p_{1_{l_1}} \end{array} \right\} \quad \dots \quad \left\{ \begin{array}{ccc} [P_{n_1}] & & [P_{n_{l_n}}] \\ p_{n_1} & \dots & p_{n_{l_n}} \end{array} \right\}}{p}$$

Contextual atomic rules

Pictorially, we can represent this as:

$$\frac{\left\{ \begin{array}{ccc} [P_{1_1}] & & [P_{1_{l_1}}] \\ p_{1_1} & \dots & p_{1_{l_1}} \end{array} \right\} \quad \dots \quad \left\{ \begin{array}{ccc} [P_{n_1}] & & [P_{n_{l_n}}] \\ p_{n_1} & \dots & p_{n_{l_n}} \end{array} \right\}}{p}$$

or like this:

$$\langle \{ \langle P_{1_i}, p_{1_i} \rangle \}_{i=1}^{l_1}, \dots, \{ \langle P_{n_i}, p_{n_i} \rangle \}_{i=1}^{l_n}, p \rangle$$

Examples of atomic rules

- $\langle \{ \langle \emptyset, a \rangle \}, c \rangle$
- $\langle \{ \langle \emptyset, a \rangle \}, \{ \langle c, d, e \rangle, \langle f, g \rangle \}, q \rangle$
- $\langle \{ \langle a, b \rangle \}, \{ \langle \emptyset, e \rangle \}, \langle f, g \rangle, q \rangle$
- $\langle \langle \emptyset, a \rangle, c \rangle$
- $\langle \langle \emptyset, b \rangle, c \rangle$
- $\langle \langle \emptyset, c \rangle, \langle a, p \rangle, \langle b, p \rangle, p \rangle$

Presentation root directory

- 1 Bases and inference figures
- 2 Atomic derivability
- 3 Comparing relations

Context-free atomic derivability

Definition ($\triangleright_{\mathcal{B}}$)

The relation of derivability in a context-free base \mathcal{B} , is defined as so:

Ref $S \triangleright_{\mathcal{B}} p$ if $p \in S$.

App For $\langle \langle P_1, p_1 \rangle, \dots, \langle P_n, p_n \rangle, q \rangle \in \mathcal{B}$ and $S, P_i \triangleright_{\mathcal{B}} p_i$ for each $i \in \{1, \dots, n\}$ then $S \triangleright_{\mathcal{B}} q$.

Context-free atomic derivability

Definition ($\triangleright_{\mathcal{B}}$)

The relation of derivability in a context-free base \mathcal{B} , is defined as so:

Ref $S \triangleright_{\mathcal{B}} p$ if $p \in S$.

App For $\langle \langle P_1, p_1 \rangle, \dots, \langle P_n, p_n \rangle, q \rangle \in \mathcal{B}$ and $S, P_i \triangleright_{\mathcal{B}} p_i$ for each $i \in \{1, \dots, n\}$ then $S \triangleright_{\mathcal{B}} q$.

Definition ($\vdash_{\mathcal{B}}$)

The relation of derivability in a contextual base \mathcal{B} , is defined as so:

Ref $p \vdash_{\mathcal{B}} p$.

App For $\langle \{ \langle P_{1_i}, p_{1_i} \rangle \}_{i=1}^{l_1}, \dots, \{ \langle P_{n_i}, p_{n_i} \rangle \}_{i=1}^{l_n}, q \rangle \in \mathcal{B}$ and $C_i, P_{ij} \vdash_{\mathcal{B}} p_{ij}$ for each $i \in \{1, \dots, n\}$ and $j \in \{1, \dots, l_i\}$ then $C_1, \dots, C_n \vdash_{\mathcal{B}} q$.

Example derivations

Example (Derivation terminations)

Let $\mathcal{B} = \{(\Rightarrow a), (\{\Rightarrow a\}, \{\Rightarrow b\} \Rightarrow c)\}$

$$\frac{\frac{}{b \vdash_{\mathcal{B}} b} \text{Ref} \quad \frac{}{\vdash_{\mathcal{B}} a} \Rightarrow a}{b \vdash_{\mathcal{B}} c} \{\Rightarrow a\}, \{\Rightarrow b\} \Rightarrow c$$

Therefore, in \mathcal{B} , the atom c is derivable from b .

Example derivations

Example (Invalid derivation)

By the definition of $\vdash_{\mathcal{B}}$, deriving a from a, a in the empty base is not possible, i.e. $a, a \vdash_{\emptyset} a$ is not possible.

Example derivations

Example (Invalid derivation)

By the definition of $\vdash_{\mathcal{B}}$, deriving a from a, a in the empty base is not possible, i.e. $a, a \vdash_{\emptyset} a$ is not possible.

Example (Invalid derivation 2)

Let $\mathcal{B} = \{(\{ \Rightarrow a \} \Rightarrow c), (\{ a \Rightarrow c \} \Rightarrow b)\}$

$$\begin{array}{c}
 \times \\
 \hline
 \frac{a, a \vdash_{\mathcal{B}} a}{a, a \vdash_{\mathcal{B}} c} \{ \Rightarrow a \} \Rightarrow c \\
 \hline
 \frac{a \vdash_{\mathcal{B}} b}{a \vdash_{\mathcal{B}} b} \{ a \Rightarrow c \} \Rightarrow b
 \end{array}$$

We see that in this base, a is not derivable from a, a .

Example derivations

Example (Another possible fix for invalid derivation 2)

Let

$$\mathcal{D} = \{(\{\Rightarrow a\} \Rightarrow c), (\{a \Rightarrow c\} \Rightarrow b), (\{\Rightarrow d\}, \{\Rightarrow d\} \Rightarrow a), (\{\Rightarrow a\} \Rightarrow d)\}$$

$$\frac{\frac{\frac{}{a \vdash_{\mathcal{D}} a} \text{Ref}}{a \vdash_{\mathcal{D}} d} \{\Rightarrow a\} \Rightarrow d \quad \frac{\frac{}{a \vdash_{\mathcal{D}} a} \text{Ref}}{a \vdash_{\mathcal{D}} d} \{\Rightarrow a\} \Rightarrow d}{\frac{a, a \vdash_{\mathcal{D}} a}{a, a \vdash_{\mathcal{D}} c} \{\Rightarrow d\}, \{\Rightarrow d\} \Rightarrow a} \{\Rightarrow a\} \Rightarrow c$$

$$\frac{a, a \vdash_{\mathcal{D}} c}{a \vdash_{\mathcal{D}} b} \{a \Rightarrow c\} \Rightarrow b$$

Example derivations

Example (A live complicated derivation!)

Consider a base \mathcal{B} with only the following rules:

- $\{\Rightarrow c\}, \{\Rightarrow y\} \Rightarrow f$
- $\Rightarrow z$
- $\{\Rightarrow a\}, \{\Rightarrow b\}, \{\Rightarrow z\} \Rightarrow h$
- $\{\Rightarrow h\} \Rightarrow g$
- $\{\Rightarrow x\}, \{\Rightarrow g\} \Rightarrow e$
- $\{x \Rightarrow e\}, \{y \Rightarrow f\} \Rightarrow d$

is there a derivation of d from the multiset a, b, c in \mathcal{B} ?

Example derivations

Example (A live complicated derivation!)

Consider a base \mathcal{B} with only the following rules:

- $\{\Rightarrow c\}, \{\Rightarrow y\} \Rightarrow f$
- $\Rightarrow z$
- $\{\Rightarrow a\}, \{\Rightarrow b\}, \{\Rightarrow z\} \Rightarrow h$
- $\{\Rightarrow h\} \Rightarrow g$
- $\{\Rightarrow x\}, \{\Rightarrow g\} \Rightarrow e$
- $\{x \Rightarrow e\}, \{y \Rightarrow f\} \Rightarrow d$

is there a derivation of d from the multiset a, b, c in \mathcal{B} ?

Disclaimer: The fact I couldn't fit this derivation on the slide has nothing to do with why we are doing this derivation live :-)

Example derivations

Example (A very interesting derivation)

Let $\mathcal{B} = \{(\{ \Rightarrow b \} \Rightarrow c), (\{ \Rightarrow a \}, \{ b \Rightarrow c \} \Rightarrow d), (\{ \Rightarrow d, \Rightarrow a \} \Rightarrow e), (\{ a \Rightarrow e \} \Rightarrow f)\}$

$$\begin{array}{c}
 \frac{}{a \vdash_{\mathcal{B}} a} \text{Ref} \quad \frac{\frac{}{b \vdash_{\mathcal{B}} b} \text{Ref}}{b \vdash_{\mathcal{B}} c} \{ \Rightarrow b \} \Rightarrow c}{\frac{}{a \vdash_{\mathcal{B}} d} \{ \Rightarrow a \}, \{ b \Rightarrow c \} \Rightarrow d} \text{Ref} \quad \frac{}{a \vdash_{\mathcal{B}} a} \text{Ref} \\
 \frac{\frac{}{a \vdash_{\mathcal{B}} d} \{ \Rightarrow a \}, \{ b \Rightarrow c \} \Rightarrow d \quad \frac{}{a \vdash_{\mathcal{B}} a} \{ \Rightarrow d, \Rightarrow a \} \Rightarrow e}{\frac{a \vdash_{\mathcal{B}} e}{\vdash_{\mathcal{B}} f} \{ a \Rightarrow e \} \Rightarrow f} \text{Ref}
 \end{array}$$

Presentation root directory

- 1 Bases and inference figures
- 2 Atomic derivability
- 3 Comparing relations




Comparing relations

Thank you!





```
C:\>ver  
  
SCP/DOS Version 0.98  
  
C:\>          Thank you for listening!\  
              Comments? Observations? Please ask and/or feel free\  
              to email me at y.buzoku@ucl.ac.uk
```

Figure: Thank you from DOS! :D

References I

-  G.M. Bierman, *On intuitionistic linear logic*, Tech. Report UCAM-CL-TR-346, University of Cambridge, Computer Laboratory, August 1994.
-  Yli Buzoku, *A proof-theoretic semantics for intuitionistic linear logic*, 2024.
-  Alexander V. Gheorghiu, Tao Gu, and David J. Pym, *Proof-theoretic semantics for intuitionistic multiplicative linear logic*, Automated Reasoning with Analytic Tableaux and Related Methods (Cham) (Revantha Ramanayake and Josef Urban, eds.), Springer Nature Switzerland, 2023, pp. 367–385.

References II

-  J.-Y. Girard, *Linear logic: its syntax and semantics*, London Mathematical Society Lecture Note Series, p. 1–42, Cambridge University Press, 1995.
-  Alexander V. Gheorghiu and David J. Pym, *From proof-theoretic validity to base-extension semantics for intuitionistic propositional logic*, 2022.
-  Sara Negri, *A normalizing system of natural deduction for intuitionistic linear logic*, *Archive for Mathematical Logic* **41** (2002), no. 8, 789–810.
-  Tor Sandqvist, *An inferentialist interpretation of classical logic*, Ph.D. thesis, Uppsala universitet, 2005.

References III



_____, *Base-extension semantics for intuitionistic sentential logic*,
Log. J. IGPL **23** (2015), 719–731.