

MT 4113: Computing in Statistics

Computer intensive statistics

Lecture 6: Permutation, randomization, and Monte-Carlo tests

Len Thomas

Oct 10 2018



University
of
St Andrews

Outline

- 1 Introduction
 - Revision – Traditional parametric and nonparametric methods
 - Motivating example
- 2 Permutation methods
 - Permutation test
 - Permutation interval
 - Summary
- 3 Randomization methods
 - Overview
 - Summary
- 4 Monte-Carlo tests
 - Overview
 - Summary



Outline

- 1 Introduction
 - Revision – Traditional parametric and nonparametric methods
 - Motivating example
- 2 Permutation methods
 - Permutation test
 - Permutation interval
 - Summary
- 3 Randomization methods
 - Overview
 - Summary
- 4 Monte-Carlo tests
 - Overview
 - Summary

Problem statement – inference from one sample

- We have a sample of data $x = x_1, \dots, x_n$ that are realizations of iid random variables $X = X_1, \dots, X_n$ with pdf $f()$.
- We wish to make inferences about some population characteristic – say the mean μ of $f()$.



Problem statement – inference from one sample

- We have a sample of data $x = x_1, \dots, x_n$ that are realizations of iid random variables $X = X_1, \dots, X_n$ with pdf $f()$.
- We wish to make inferences about some population characteristic – say the mean μ of $f()$.



Parametric methods

- Assume $f()$ is a known function with parameters θ – e.g., $f(\theta) = N(\mu, \sigma^2)$.
- For a (two-sided) hypothesis test
 - Specify the null hypothesis, H_0 – e.g., $\mu = \mu_0$
 - Construct a test statistic $T(X)$ that has a known distribution under H_0
 - e.g.,

$$T(x) = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

which has a t -distribution with mean 0, variance 1 and d.f. = $n - 1$, if H_0 is true

- Calculate $P(|T(X)| \geq |T(x)|; \mu_0)$ – the p -value.
- If $p\text{-value} \leq \alpha$ the result is statistically significant.



Parametric methods

- Assume $f()$ is a known function with parameters θ – e.g., $f(\theta) = N(\mu, \sigma^2)$.
- For a (two-sided) hypothesis test
 - Specify the null hypothesis, H_0 – e.g., $\mu = \mu_0$
 - Construct a test statistic $T(X)$ that has a known distribution under H_0
 - e.g.,

$$T(x) = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

which has a t -distribution with mean 0, variance 1 and d.f. = $n - 1$, if H_0 is true

- Calculate $P(|T(X)| \geq |T(x)|; \mu_0)$ – the p -value.
- If $p\text{-value} \leq \alpha$ the result is statistically significant.



Parametric methods

- Assume $f()$ is a known function with parameters θ – e.g., $f(\theta) = N(\mu, \sigma^2)$.
- For a (two-sided) hypothesis test
 - Specify the null hypothesis, H_0 – e.g., $\mu = \mu_0$
 - Construct a test statistic $T(X)$ that has a known distribution under H_0
 - e.g.,

$$T(x) = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

which has a t -distribution with mean 0, variance 1 and d.f. = $n - 1$, if H_0 is true

- Calculate $P(|T(X)| \geq |T(x)|; \mu_0)$ – the p -value.
- If $p\text{-value} \leq \alpha$ the result is statistically significant.



Parametric methods

- Assume $f()$ is a known function with parameters θ – e.g., $f(\theta) = N(\mu, \sigma^2)$.
- For a (two-sided) hypothesis test
 - Specify the null hypothesis, H_0 – e.g., $\mu = \mu_0$
 - Construct a test statistic $T(X)$ that has a known distribution under H_0
 - e.g.,

$$T(x) = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

which has a t -distribution with mean 0, variance 1 and d.f. = $n - 1$, if H_0 is true

- Calculate $P(|T(X)| \geq |T(x)|; \mu_0)$ – the p -value.
- If $p\text{-value} \leq \alpha$ the result is statistically significant.



Parametric methods

- Assume $f()$ is a known function with parameters θ – e.g., $f(\theta) = N(\mu, \sigma^2)$.
- For a (two-sided) hypothesis test
 - Specify the null hypothesis, H_0 – e.g., $\mu = \mu_0$
 - Construct a test statistic $T(X)$ that has a known distribution under H_0
 - e.g.,

$$T(x) = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

which has a t -distribution with mean 0, variance 1 and d.f. = $n - 1$, if H_0 is true

- Calculate $P(|T(X)| \geq |T(x)|; \mu_0)$ – the p -value.
- If $p\text{-value} \leq \alpha$ the result is statistically significant.



Parametric methods

- Assume $f()$ is a known function with parameters θ – e.g., $f(\theta) = N(\mu, \sigma^2)$.
- For a (two-sided) hypothesis test
 - Specify the null hypothesis, H_0 – e.g., $\mu = \mu_0$
 - Construct a test statistic $T(X)$ that has a known distribution under H_0
 - e.g.,

$$T(x) = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

which has a t -distribution with mean 0, variance 1 and d.f. = $n - 1$, if H_0 is true

- Calculate $P(|T(X)| \geq |T(x)|; \mu_0)$ – the p -value.
- If $p\text{-value} \leq \alpha$ the result is statistically significant.



Parametric methods

- Assume $f()$ is a known function with parameters θ – e.g., $f(\theta) = N(\mu, \sigma^2)$.
- For a (two-sided) hypothesis test
 - Specify the null hypothesis, H_0 – e.g., $\mu = \mu_0$
 - Construct a test statistic $T(X)$ that has a known distribution under H_0
 - e.g.,

$$T(x) = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

which has a t -distribution with mean 0, variance 1 and d.f. = $n - 1$, if H_0 is true

- Calculate $P(|T(X)| \geq |T(x)|; \mu_0)$ – the p -value.
- If $p\text{-value} \leq \alpha$ the result is statistically significant.



Parametric methods (contd.)

- For (two-sided) confidence intervals on μ
 - For $((1 - \alpha) \times 100)\%$ limits, find the scalar values $L(X)$ and $U(X)$ such that

$$P(L(X) < \mu \text{ and } U(X) > \mu) = 1 - \alpha$$

- Many methods for finding the limits
- E.g., inverting the test statistic
 - limits are the smallest and largest values of μ_0 that are not rejected under $H_0 : \mu = \mu_0$
 - E.g., t -based intervals $\bar{x} \pm t_{\alpha/2, n-1} s / \sqrt{n}$
- So, for parametric methods, you need to specify two distributions:
 - The pdf generating the data: $f(\theta)$
 - The distribution of the test statistic: $T(X)$

Parametric methods (contd.)

- For (two-sided) confidence intervals on μ
 - For $((1 - \alpha) \times 100)\%$ limits, find the scalar values $L(X)$ and $U(X)$ such that

$$P(L(X) < \mu \text{ and } U(X) > \mu) = 1 - \alpha$$

- Many methods for finding the limits
- E.g., inverting the test statistic
 - limits are the smallest and largest values of μ_0 that are not rejected under $H_0 : \mu = \mu_0$
 - E.g., t -based intervals $\bar{x} \pm t_{\alpha/2, n-1} s / \sqrt{n}$
- So, for parametric methods, you need to specify two distributions:
 - The pdf generating the data: $f(\theta)$
 - The distribution of the test statistic: $T(X)$

Parametric methods (contd.)

- For (two-sided) confidence intervals on μ
 - For $((1 - \alpha) \times 100)\%$ limits, find the scalar values $L(X)$ and $U(X)$ such that

$$P(L(X) < \mu \text{ and } U(X) > \mu) = 1 - \alpha$$

- Many methods for finding the limits
 - E.g., inverting the test statistic
 - limits are the smallest and largest values of μ_0 that are not rejected under $H_0 : \mu = \mu_0$
 - E.g., t -based intervals $\bar{x} \pm t_{\alpha/2, n-1} s / \sqrt{n}$
- So, for parametric methods, you need to specify two distributions:
 - The pdf generating the data: $f(\theta)$
 - The distribution of the test statistic: $T(X)$



Parametric methods (contd.)

- For (two-sided) confidence intervals on μ
 - For $((1 - \alpha) \times 100)\%$ limits, find the scalar values $L(X)$ and $U(X)$ such that

$$P(L(X) < \mu \text{ and } U(X) > \mu) = 1 - \alpha$$

- Many methods for finding the limits
- E.g., inverting the test statistic
 - limits are the smallest and largest values of μ_0 that are not rejected under $H_0 : \mu = \mu_0$
 - E.g., t -based intervals $\bar{x} \pm t_{\alpha/2, n-1} s / \sqrt{n}$
- So, for parametric methods, you need to specify two distributions:
 - The pdf generating the data: $f(\theta)$
 - The distribution of the test statistic: $T(X)$

Parametric methods (contd.)

- For (two-sided) confidence intervals on μ
 - For $((1 - \alpha) \times 100)\%$ limits, find the scalar values $L(X)$ and $U(X)$ such that

$$P(L(X) < \mu \text{ and } U(X) > \mu) = 1 - \alpha$$

- Many methods for finding the limits
- E.g., inverting the test statistic
 - limits are the smallest and largest values of μ_0 that are not rejected under $H_0 : \mu = \mu_0$
 - E.g., t -based intervals $\bar{x} \pm t_{\alpha/2, n-1} s / \sqrt{n}$
- So, for parametric methods, you need to specify two distributions:

- The pdf generating the data: $f(\theta)$
- The distribution of the test statistic: $T(X)$



Parametric methods (contd.)

- For (two-sided) confidence intervals on μ
 - For $((1 - \alpha) \times 100)\%$ limits, find the scalar values $L(X)$ and $U(X)$ such that

$$P(L(X) < \mu \text{ and } U(X) > \mu) = 1 - \alpha$$

- Many methods for finding the limits
- E.g., inverting the test statistic
 - limits are the smallest and largest values of μ_0 that are not rejected under $H_0 : \mu = \mu_0$
 - E.g., t -based intervals $\bar{x} \pm t_{\alpha/2, n-1} s / \sqrt{n}$
- So, for parametric methods, you need to specify two distributions:

- The pdf generating the data: $f(\theta)$
- The distribution of the test statistic: $T(X)$



Parametric methods (contd.)

- For (two-sided) confidence intervals on μ
 - For $((1 - \alpha) \times 100)\%$ limits, find the scalar values $L(X)$ and $U(X)$ such that

$$P(L(X) < \mu \text{ and } U(X) > \mu) = 1 - \alpha$$

- Many methods for finding the limits
- E.g., inverting the test statistic
 - limits are the smallest and largest values of μ_0 that are not rejected under $H_0 : \mu = \mu_0$
 - E.g., t -based intervals $\bar{x} \pm t_{\alpha/2, n-1} s / \sqrt{n}$
- So, for parametric methods, you need to specify two distributions:
 - The pdf generating the data: $f(\theta)$
 - The distribution of the test statistic: $T(X)$

Parametric methods (contd.)

- For (two-sided) confidence intervals on μ
 - For $((1 - \alpha) \times 100)\%$ limits, find the scalar values $L(X)$ and $U(X)$ such that

$$P(L(X) < \mu \text{ and } U(X) > \mu) = 1 - \alpha$$

- Many methods for finding the limits
- E.g., inverting the test statistic
 - limits are the smallest and largest values of μ_0 that are not rejected under $H_0 : \mu = \mu_0$
 - E.g., t -based intervals $\bar{x} \pm t_{\alpha/2, n-1} s / \sqrt{n}$
- So, for parametric methods, you need to specify two distributions:
 - The pdf generating the data: $f(\theta)$
 - The distribution of the test statistic: $T(X)$

Parametric methods (contd.)

- For (two-sided) confidence intervals on μ
 - For $((1 - \alpha) \times 100)\%$ limits, find the scalar values $L(X)$ and $U(X)$ such that

$$P(L(X) < \mu \text{ and } U(X) > \mu) = 1 - \alpha$$

- Many methods for finding the limits
- E.g., inverting the test statistic
 - limits are the smallest and largest values of μ_0 that are not rejected under $H_0 : \mu = \mu_0$
 - E.g., t -based intervals $\bar{x} \pm t_{\alpha/2, n-1} s / \sqrt{n}$
- So, for parametric methods, you need to specify two distributions:
 - The pdf generating the data: $f(\theta)$
 - The distribution of the test statistic: $T(X)$

Traditional nonparametric methods

- Avoid fully specifying the distribution of $f()$.
- E.g., Wilcoxon signed ranks test
 - $f()$ is symmetric about some median m
 - $H_0 : m = m_0$
 - Test statistic

$$T(x) = \sum_{i: x_i \neq m_0} \text{rank}(|x_i - m_0|) \text{sign}(x_i - m_0)$$

where $\text{rank}(x)$ denotes the rank order of x (smallest has value 1, next smallest 2, etc.) and $\text{sign}(x - y)$ is a sign function, having value 1 if $x > y$ and -1 if $x < y$.

- Under H_0 , $T(X)$ has (approximately) a normal distribution
- For confidence limits on m , can invert the test statistic



Traditional nonparametric methods

- Avoid fully specifying the distribution of $f()$.
- E.g., Wilcoxon signed ranks test
 - $f()$ is symmetric about some median m
 - $H_0 : m = m_0$
 - Test statistic

$$T(x) = \sum_{i: x_i \neq m_0} \text{rank}(|x_i - m_0|) \text{sign}(x_i - m_0)$$

where $\text{rank}(x)$ denotes the rank order of x (smallest has value 1, next smallest 2, etc.) and $\text{sign}(x - y)$ is a sign function, having value 1 if $x > y$ and -1 if $x < y$.

- Under H_0 , $T(X)$ has (approximately) a normal distribution
- For confidence limits on m , can invert the test statistic



Traditional nonparametric methods

- Avoid fully specifying the distribution of $f()$.
- E.g., Wilcoxon signed ranks test
 - $f()$ is symmetric about some median m
 - $H_0 : m = m_0$
 - Test statistic

$$T(x) = \sum_{i: x_i \neq m_0} \text{rank}(|x_i - m_0|) \text{sign}(x_i - m_0)$$

where $\text{rank}(x)$ denotes the rank order of x (smallest has value 1, next smallest 2, etc.) and $\text{sign}(x - y)$ is a sign function, having value 1 if $x > y$ and -1 if $x < y$.

- Under H_0 , $T(X)$ has (approximately) a normal distribution
- For confidence limits on m , can invert the test statistic



Traditional nonparametric methods

- Avoid fully specifying the distribution of $f()$.
- E.g., Wilcoxon signed ranks test
 - $f()$ is symmetric about some median m
 - $H_0 : m = m_0$
 - Test statistic

$$T(x) = \sum_{i: x_i \neq m_0} \text{rank}(|x_i - m_0|) \text{sign}(x_i - m_0)$$

where $\text{rank}(x)$ denotes the rank order of x (smallest has value 1, next smallest 2, etc.) and $\text{sign}(x - y)$ is a sign function, having value 1 if $x > y$ and -1 if $x < y$.

- Under H_0 , $T(X)$ has (approximately) a normal distribution
- For confidence limits on m , can invert the test statistic



Traditional nonparametric methods

- Avoid fully specifying the distribution of $f()$.
- E.g., Wilcoxon signed ranks test
 - $f()$ is symmetric about some median m
 - $H_0 : m = m_0$
 - Test statistic

$$T(x) = \sum_{i: x_i \neq m_0} \text{rank}(|x_i - m_0|) \text{sign}(x_i - m_0)$$

where $\text{rank}(x)$ denotes the rank order of x (smallest has value 1, next smallest 2, etc.) and $\text{sign}(x - y)$ is a sign function, having value 1 if $x > y$ and -1 if $x < y$.

- Under H_0 , $T(X)$ has (approximately) a normal distribution
- For confidence limits on m , can invert the test statistic



Traditional nonparametric methods

- Avoid fully specifying the distribution of $f()$.
- E.g., Wilcoxon signed ranks test
 - $f()$ is symmetric about some median m
 - $H_0 : m = m_0$
 - Test statistic

$$T(x) = \sum_{i: x_i \neq m_0} \text{rank}(|x_i - m_0|) \text{sign}(x_i - m_0)$$

where $\text{rank}(x)$ denotes the rank order of x (smallest has value 1, next smallest 2, etc.) and $\text{sign}(x - y)$ is a sign function, having value 1 if $x > y$ and -1 if $x < y$.

- Under H_0 , $T(X)$ has (approximately) a normal distribution
- For confidence limits on m , can invert the test statistic



Traditional nonparametric methods

- Avoid fully specifying the distribution of $f()$.
- E.g., Wilcoxon signed ranks test
 - $f()$ is symmetric about some median m
 - $H_0 : m = m_0$
 - Test statistic

$$T(x) = \sum_{i: x_i \neq m_0} \text{rank}(|x_i - m_0|) \text{sign}(x_i - m_0)$$

where $\text{rank}(x)$ denotes the rank order of x (smallest has value 1, next smallest 2, etc.) and $\text{sign}(x - y)$ is a sign function, having value 1 if $x > y$ and -1 if $x < y$.

- Under H_0 , $T(X)$ has (approximately) a normal distribution
- For confidence limits on m , can invert the test statistic



Computer-intensive statistical methods

- With computer intensive methods, you can avoid the need to fully specify either the distribution of $f()$ and/or $T(X)$
- Examples:
 - Permutation-based approach to Wilcoxon test (avoids the need to specify distribution for $T(X)$; Wilcoxon already avoids $f()$)
 - Monte-Carlo-based version of the t -test (avoids specification of $T(X)$, but not $f(\theta)$)
 - Parametric and non-parametric bootstrap confidence intervals (again avoids $T(X)$; either specify or not $f(\theta)$)



Computer-intensive statistical methods

- With computer intensive methods, you can avoid the need to fully specify either the distribution of $f()$ and/or $T(X)$
- Examples:
 - Permutation-based approach to Wilcoxon test (avoids the need to specify distribution for $T(X)$; Wilcoxon already avoids $f()$)
 - Monte-Carlo-based version of the t -test (avoids specification of $T(X)$, but not $f(\theta)$)
 - Parametric and non-parametric bootstrap confidence intervals (again avoids $T(X)$; either specify or not $f(\theta)$)



Computer-intensive statistical methods

- With computer intensive methods, you can avoid the need to fully specify either the distribution of $f()$ and/or $T(X)$
- Examples:
 - Permutation-based approach to Wilcoxon test (avoids the need to specify distribution for $T(X)$; Wilcoxon already avoids $f()$)
 - Monte-Carlo-based version of the t -test (avoids specification of $T(X)$, but not $f(\theta)$)
 - Parametric and non-parametric bootstrap confidence intervals (again avoids $T(X)$; either specify or not $f(\theta)$)



Computer-intensive statistical methods

- With computer intensive methods, you can avoid the need to fully specify either the distribution of $f()$ and/or $T(X)$
- Examples:
 - Permutation-based approach to Wilcoxon test (avoids the need to specify distribution for $T(X)$; Wilcoxon already avoids $f()$)
 - Monte-Carlo-based version of the t -test (avoids specification of $T(X)$, but not $f(\theta)$)
 - Parametric and non-parametric bootstrap confidence intervals (again avoids $T(X)$; either specify or not $f(\theta)$)

Computer-intensive statistical methods

- With computer intensive methods, you can avoid the need to fully specify either the distribution of $f()$ and/or $T(X)$
- Examples:
 - Permutation-based approach to Wilcoxon test (avoids the need to specify distribution for $T(X)$; Wilcoxon already avoids $f()$)
 - Monte-Carlo-based version of the t -test (avoids specification of $T(X)$, but not $f(\theta)$)
 - Parametric and non-parametric bootstrap confidence intervals (again avoids $T(X)$; either specify or not $f(\theta)$)

Outline

1 Introduction

- Revision – Traditional parametric and nonparametric methods
- **Motivating example**

2 Permutation methods

- Permutation test
- Permutation interval
- Summary

3 Randomization methods

- Overview
- Summary


4 Monte-Carlo tests

- Overview
- Summary



Motivating example


Pine martin habitat preference data



- 0.13, -0.01, -0.01, 0.42, -0.02, 0.01, 0.09, 0.03, 0.04, 0.06, 0.12, 0.03
- Shapiro-Wilk test for normality $W = 0.70$, $p < 0.001$

Motivating example

Pine martin habitat preference data




- 0.13, -0.01, -0.01, 0.42, -0.02, 0.01, 0.09, 0.03, 0.04, 0.06, 0.12, 0.03
- Shapiro-Wilk test for normality $W = 0.70$, $p < 0.001$

Motivating example

Pine martin example: parametric test

- Assume data come from a normal distribution
 - $H_0 : \mu = 0$
 - t -test: $t = 2.147$, d.f.=11, $p = 0.055$
 - $\bar{x} = 0.074$, 95% t -based CI: $(-0.002, 0.150)$
- Could try transforming – e.g., $\log(x_i + 0.03)$:




- $H_0 : \log(\mu + 0.03) = \log(0.03)$
- t -test: $t = 2.484$, d.f.=11, $p = 0.030$
- 95% t -based CI (back-transformed): $(0.003, 0.095)$



Motivating example

Pine martin example: parametric test

- Assume data come from a normal distribution
 - $H_0 : \mu = 0$
 - t -test: $t = 2.147$, d.f.=11, $p = 0.055$
 - $\bar{x} = 0.074$, 95% t -based CI: (-0.002, 0.150)
 - Could try transforming – e.g., $\log(x_i + 0.03)$:




- $H_0 : \log(\mu + 0.03) = \log(0.03)$
- t -test: $t = 2.484$, d.f.=11, $p = 0.030$
- 95% t -based CI (back-transformed): (0.003, 0.095)



Motivating example

Pine martin example: parametric test

- Assume data come from a normal distribution
 - $H_0 : \mu = 0$
 - t -test: $t = 2.147$, d.f.=11, $p = 0.055$
 - $\bar{x} = 0.074$, 95% t -based CI: $(-0.002, 0.150)$
- Could try transforming – e.g., $\log(x_i + 0.03)$:




- $H_0 : \log(\mu + 0.03) = \log(0.03)$
- t -test: $t = 2.484$, d.f.=11, $p = 0.030$
- 95% t -based CI (back-transformed): $(0.003, 0.095)$



Motivating example

Pine martin example: parametric test

- Assume data come from a normal distribution
 - $H_0 : \mu = 0$
 - t -test: $t = 2.147$, d.f.=11, $p = 0.055$
 - $\bar{x} = 0.074$, 95% t -based CI: $(-0.002, 0.150)$
- Could try transforming – e.g., $\log(x_i + 0.03)$:




- $H_0 : \log(\mu + 0.03) = \log(0.03)$
- t -test: $t = 2.484$, d.f.=11, $p = 0.030$
- 95% t -based CI (back-transformed): $(0.003, 0.095)$



Motivating example

Pine martin example: parametric test

- Assume data come from a normal distribution
 - $H_0 : \mu = 0$
 - t -test: $t = 2.147$, d.f.=11, $p = 0.055$
 - $\bar{x} = 0.074$, 95% t -based CI: $(-0.002, 0.150)$
- Could try transforming – e.g., $\log(x_i + 0.03)$:




- $H_0 : \log(\mu + 0.03) = \log(0.03)$
- t -test: $t = 2.484$, d.f.=11, $p = 0.030$
- 95% t -based CI (back-transformed): $(0.003, 0.095)$

Motivating example

Pine martin example: parametric test

- Assume data come from a normal distribution
 - $H_0 : \mu = 0$
 - t -test: $t = 2.147$, d.f.=11, $p = 0.055$
 - $\bar{x} = 0.074$, 95% t -based CI: $(-0.002, 0.150)$
- Could try transforming – e.g., $\log(x_i + 0.03)$:




- $H_0 : \log(\mu + 0.03) = \log(0.03)$
- t -test: $t = 2.484$, d.f.=11, $p = 0.030$
- 95% t -based CI (back-transformed): $(0.003, 0.095)$



Motivating example

Pine martin example: parametric test

- Assume data come from a normal distribution
 - $H_0 : \mu = 0$
 - t -test: $t = 2.147$, d.f.=11, $p = 0.055$
 - $\bar{x} = 0.074$, 95% t -based CI: $(-0.002, 0.150)$
- Could try transforming – e.g., $\log(x_i + 0.03)$:




- $H_0 : \log(\mu + 0.03) = \log(0.03)$
- t -test: $t = 2.484$, d.f.=11, $p = 0.030$
- 95% t -based CI (back-transformed): $(0.003, 0.095)$



Motivating example

Pine martin example: parametric test

- Assume data come from a normal distribution
 - $H_0 : \mu = 0$
 - t -test: $t = 2.147$, d.f.=11, $p = 0.055$
 - $\bar{x} = 0.074$, 95% t -based CI: $(-0.002, 0.150)$
- Could try transforming – e.g., $\log(x_i + 0.03)$:



- $H_0 : \log(\mu + 0.03) = \log(0.03)$
- t -test: $t = 2.484$, d.f.=11, $p = 0.030$
- 95% t -based CI (back-transformed): $(0.003, 0.095)$



Motivating example

Pine martin example: parametric test (contd.)

- Problems:
 - Result not invariant to transformation: what transformation to use?
 - If we use log, then what about bias correction?

Motivating example

Pine martin example: parametric test (contd.)

- Problems:
 - Result not invariant to transformation: what transformation to use?
 - If we use log, then what about bias correction?



Motivating example

Pine martin example: parametric test (contd.)

- Problems:
 - Result not invariant to transformation: what transformation to use?
 - If we use log, then what about bias correction?



Motivating example

Pine martin example: traditional nonparametric test

- Wilcoxon signed-ranks test:
 - $H_0 : m = 0$
 - $p = 0.012$
 - 95% CI on m : (0.010, 0.120)
- Disadvantages:
 - Requires us to formulate H_0 in terms of the median
 - Less powerful than a t -test when parametric assumptions met



Motivating example

Pine martin example: traditional nonparametric test

- Wilcoxon signed-ranks test:
 - $H_0 : m = 0$
 - $p = 0.012$
 - 95% CI on m : (0.010, 0.120)
- Disadvantages:
 - Requires us to formulate H_0 in terms of the median
 - Less powerful than a t -test when parametric assumptions met



Motivating example

Pine martin example: traditional nonparametric test

- Wilcoxon signed-ranks test:
 - $H_0 : m = 0$
 - $p = 0.012$
 - 95% CI on m : (0.010, 0.120)
- Disadvantages:
 - Requires us to formulate H_0 in terms of the median
 - Less powerful than a t -test when parametric assumptions met



Motivating example

Pine martin example: traditional nonparametric test

- Wilcoxon signed-ranks test:
 - $H_0 : m = 0$
 - $p = 0.012$
 - 95% CI on m : (0.010, 0.120)
- Disadvantages:
 - Requires us to formulate H_0 in terms of the median
 - Less powerful than a t -test when parametric assumptions met



Motivating example

Pine martin example: traditional nonparametric test

- Wilcoxon signed-ranks test:
 - $H_0 : m = 0$
 - $p = 0.012$
 - 95% CI on m : (0.010, 0.120)
- Disadvantages:
 - Requires us to formulate H_0 in terms of the median
 - Less powerful than a t -test when parametric assumptions met



Pine martin example: traditional nonparametric test

- Wilcoxon signed-ranks test:
 - $H_0 : m = 0$
 - $p = 0.012$
 - 95% CI on m: (0.010, 0.120)
- Disadvantages:
 - Requires us to formulate H_0 in terms of the median
 - Less powerful than a t -test when parametric assumptions met



Motivating example

Pine martin example: traditional nonparametric test

- Wilcoxon signed-ranks test:
 - $H_0 : m = 0$
 - $p = 0.012$
 - 95% CI on m : (0.010, 0.120)
- Disadvantages:
 - Requires us to formulate H_0 in terms of the median
 - Less powerful than a t -test when parametric assumptions met



Outline

- 1 Introduction
 - Revision – Traditional parametric and nonparametric methods
 - Motivating example
- 2 **Permutation methods**
 - **Permutation test**
 - Permutation interval
 - Summary
- 3 Randomization methods
 - Overview
 - Summary
- 4 Monte-Carlo tests
 - Overview
 - Summary



Pine martin example

- Under $H_0 : \mu = \mu_0$, and assuming the distribution of x_i is symmetric about μ_0 , then
 - $+(x_i - \mu_0)$ is just as likely to occur in the data as $-(x_i - \mu_0)$
 - For $\mu_0 = 0$ then $+x_i$ is just as likely as $-x_i$
 - So, can produce alternative equally likely realizations of the data by swapping any x_i for $-x_i$



Pine martin example

- Under $H_0 : \mu = \mu_0$, and assuming the distribution of x_i is symmetric about μ_0 , then
- $+(x_i - \mu_0)$ is just as likely to occur in the data as $-(x_i - \mu_0)$
- For $\mu_0 = 0$ then $+x_i$ is just as likely as $-x_i$
- So, can produce alternative equally likely realizations of the data by swapping any x_i for $-x_i$



Pine martin example

- Under $H_0 : \mu = \mu_0$, and assuming the distribution of x_i is symmetric about μ_0 , then
- $+(x_i - \mu_0)$ is just as likely to occur in the data as $-(x_i - \mu_0)$
- For $\mu_0 = 0$ then $+x_i$ is just as likely as $-x_i$
- So, can produce alternative equally likely realizations of the data by swapping any x_i for $-x_i$



Pine martin example

- Under $H_0 : \mu = \mu_0$, and assuming the distribution of x_i is symmetric about μ_0 , then
- $+(x_i - \mu_0)$ is just as likely to occur in the data as $-(x_i - \mu_0)$
- For $\mu_0 = 0$ then $+x_i$ is just as likely as $-x_i$
- So, can produce alternative equally likely realizations of the data by swapping any x_i for $-x_i$



Permutations

- Given 12 data points, there are $2^{12} = 4096$ ways to do this:

- 0.13, 0.01, 0.01, 0.42, 0.02, 0.01, 0.09, 0.03, 0.04, 0.06, 0.12, 0.03
- 0.13, 0.01, 0.01, 0.42, 0.02, 0.01, 0.09, 0.03, 0.04, 0.06, 0.12, 0.03
- 0.13, -0.01, 0.01, 0.42, 0.02, 0.01, 0.09, 0.03, 0.04, 0.06, 0.12, 0.03
- 0.13, 0.01, -0.01, 0.42, 0.02, 0.01, 0.09, 0.03, 0.04, 0.06, 0.12, 0.03
- ⋮
- 0.13, -0.01, -0.01, -0.42, -0.02, -0.01, -0.09, -0.03, -0.04, -0.06, -0.12, -0.03



Permutations

- Given 12 data points, there are $2^{12} = 4096$ ways to do this:
 - 0.13, 0.01, 0.01, 0.42, 0.02, 0.01, 0.09, 0.03, 0.04, 0.06, 0.12, 0.03
 - 0.13, 0.01, 0.01, 0.42, 0.02, 0.01, 0.09, 0.03, 0.04, 0.06, 0.12, 0.03
 - 0.13, -0.01, 0.01, 0.42, 0.02, 0.01, 0.09, 0.03, 0.04, 0.06, 0.12, 0.03
 - 0.13, 0.01, -0.01, 0.42, 0.02, 0.01, 0.09, 0.03, 0.04, 0.06, 0.12, 0.03
 - ⋮
 - 0.13, -0.01, -0.01, -0.42, -0.02, -0.01, -0.09, -0.03, -0.04, -0.06, -0.12, -0.03



Permutation test


- Find what proportion of the permutations have $|\bar{X}|$ greater than the observed $|\bar{x}|$.

- 48 out of 4096 are as extreme - so $p = 48/4096 = 0.012$

Permutation test

Permutation test

- Find what proportion of the permutations have $|\bar{X}|$ greater than the observed $|\bar{x}|$.



- 48 out of 4096 are as extreme - so $p = 48/4096 = 0.012$

Permuatation tests - the basic idea

- Pick a null hypothesis H_0 and test statistic $T(X)$
- Under H_0 , all possible appropriate data permutations (re-orderings) are equally likely
- So, calculate $T(X)$ for each permutation and see how extreme $T(x)$ is compared with these
 - If $T(x)$ is extreme, reject H_0
 - If $T(x)$ is not extreme, don't reject H_0

No need to specify distributions for:

- $f()$ - no parametric definition needed
- $T(X)$ - obtain this via the randomization procedure



Permuatation tests - the basic idea

- Pick a null hypothesis H_0 and test statistic $T(X)$
- Under H_0 , all possible appropriate data permutations (re-orderings) are equally likely
- So, calculate $T(X)$ for each permutation and see how extreme $T(x)$ is compared with these
 - If $T(x)$ is extreme, reject H_0
 - If $T(x)$ is not extreme, don't reject H_0

No need to specify distributions for:

- $f()$ - no parametric definition needed
- $T(X)$ - obtain this via the randomization procedure



Permuatation tests - the basic idea

- Pick a null hypothesis H_0 and test statistic $T(X)$
- Under H_0 , all possible appropriate data permutations (re-orderings) are equally likely
- So, calculate $T(X)$ for each permutation and see how extreme $T(x)$ is compared with these
 - If $T(x)$ is extreme, reject H_0
 - If $T(x)$ is not extreme, don't reject H_0

No need to specify distributions for:

- $f()$ - no parametric definition needed
- $T(X)$ - obtain this via the randomization procedure



Permuatation tests - the basic idea

- Pick a null hypothesis H_0 and test statistic $T(X)$
- Under H_0 , all possible appropriate data permutations (re-orderings) are equally likely
- So, calculate $T(X)$ for each permutation and see how extreme $T(x)$ is compared with these
 - If $T(x)$ is extreme, reject H_0
 - If $T(x)$ is not extreme, don't reject H_0

No need to specify distributions for:

- $f()$ - no parametric definition needed
- $T(X)$ - obtain this via the randomization procedure



Permuatation tests - the basic idea

- Pick a null hypothesis H_0 and test statistic $T(X)$
- Under H_0 , all possible appropriate data permutations (re-orderings) are equally likely
- So, calculate $T(X)$ for each permutation and see how extreme $T(x)$ is compared with these
 - If $T(x)$ is extreme, reject H_0
 - If $T(x)$ is not extreme, don't reject H_0

No need to specify distributions for:

- $f()$ - no parametric definition needed
- $T(X)$ - obtain this via the randomization procedure



Permuatation tests - the basic idea

- Pick a null hypothesis H_0 and test statistic $T(X)$
- Under H_0 , all possible appropriate data permutations (re-orderings) are equally likely
- So, calculate $T(X)$ for each permutation and see how extreme $T(x)$ is compared with these
 - If $T(x)$ is extreme, reject H_0
 - If $T(x)$ is not extreme, don't reject H_0

No need to specify distributions for:

- $f()$ - no parametric definition needed
- $T(X)$ - obtain this via the randomization procedure

Permuatation tests - the basic idea

- Pick a null hypothesis H_0 and test statistic $T(X)$
- Under H_0 , all possible appropriate data permutations (re-orderings) are equally likely
- So, calculate $T(X)$ for each permutation and see how extreme $T(x)$ is compared with these
 - If $T(x)$ is extreme, reject H_0
 - If $T(x)$ is not extreme, don't reject H_0

No need to specify distributions for:

- $f()$ - no parametric definition needed
- $T(X)$ - obtain this via the randomization procedure



Choice of test statistic

- With permutation-based methods (and many other computer-intensive methods) you are free to choose whatever test statistic you like
- For example, we could use the median, trimmed mean, t -statistic, etc., etc.
- The method readily extends to other situations – e.g., two sample tests, ANOVA, etc., etc.



Choice of test statistic

- With permutation-based methods (and many other computer-intensive methods) you are free to choose whatever test statistic you like
- For example, we could use the median, trimmed mean, t -statistic, etc., etc.
- The method readily extends to other situations – e.g., two sample tests, ANOVA, etc., etc.



Choice of test statistic

- With permutation-based methods (and many other computer-intensive methods) you are free to choose whatever test statistic you like
- For example, we could use the median, trimmed mean, t -statistic, etc., etc.
- The method readily extends to other situations – e.g., two sample tests, ANOVA, etc., etc.



Outline

- 1 Introduction
 - Revision – Traditional parametric and nonparametric methods
 - Motivating example
- 2 **Permutation methods**
 - Permutation test
 - **Permutation interval**
 - Summary
- 3 Randomization methods
 - Overview
 - Summary
- 4 Monte-Carlo tests
 - Overview
 - Summary



Permutation interval

- Invert the test statistic:
Find the set of values of μ_0 such that $H_0 : \mu = \mu_0$ is not rejected
- For each potential value of μ_0 , follow the same procedure as before
 - calculating all permutations of $\pm(x_i - \mu_0)$
 - then seeing how extreme the observed \bar{x} is among the permutations
- Can use stochastic search methods¹ to find the confidence limits
- In our example, 95% CI is (0.013, 0.150) (c.f., t -based CI of (-0.002, 0.150)).

¹e.g., Robbins-Monroe search - see accompanying notes



Permutation interval

- Invert the test statistic:
Find the set of values of μ_0 such that $H_0 : \mu = \mu_0$ is not rejected
- For each potential value of μ_0 , follow the same procedure as before
 - calculating all permutations of $\pm(x_i - \mu_0)$
 - then seeing how extreme the observed \bar{x} is among the permutations
- Can use stochastic search methods¹ to find the confidence limits
- In our example, 95% CI is (0.013, 0.150) (c.f., t -based CI of (-0.002, 0.150)).

¹e.g., Robbins-Monroe search - see accompanying notes



Permutation interval

- Invert the test statistic:
Find the set of values of μ_0 such that $H_0 : \mu = \mu_0$ is not rejected
- For each potential value of μ_0 , follow the same procedure as before
 - calculating all permutations of $\pm(x_i - \mu_0)$
 - then seeing how extreme the observed \bar{x} is among the permutations
- Can use stochastic search methods¹ to find the confidence limits
- In our example, 95% CI is (0.013, 0.150) (c.f., t -based CI of (-0.002, 0.150)).

¹e.g., Robbins-Monroe search - see accompanying notes



Permutation interval

- Invert the test statistic:
Find the set of values of μ_0 such that $H_0 : \mu = \mu_0$ is not rejected
- For each potential value of μ_0 , follow the same procedure as before
 - calculating all permutations of $\pm(x_i - \mu_0)$
 - then seeing how extreme the observed \bar{x} is among the permutations
- Can use stochastic search methods¹ to find the confidence limits
- In our example, 95% CI is (0.013, 0.150) (c.f., t -based CI of (-0.002, 0.150)).

¹e.g., Robbins-Monroe search - see accompanying notes



Permutation interval

- Invert the test statistic:
Find the set of values of μ_0 such that $H_0 : \mu = \mu_0$ is not rejected
- For each potential value of μ_0 , follow the same procedure as before
 - calculating all permutations of $\pm(x_i - \mu_0)$
 - then seeing how extreme the observed \bar{x} is among the permutations
- Can use stochastic search methods¹ to find the confidence limits
- In our example, 95% CI is (0.013, 0.150) (c.f., t -based CI of (-0.002, 0.150)).

¹e.g., Robbins-Monroe search - see accompanying notes



Permutation interval

- Invert the test statistic:
Find the set of values of μ_0 such that $H_0 : \mu = \mu_0$ is not rejected
- For each potential value of μ_0 , follow the same procedure as before
 - calculating all permutations of $\pm(x_i - \mu_0)$
 - then seeing how extreme the observed \bar{x} is among the permutations
- Can use stochastic search methods¹ to find the confidence limits
- In our example, 95% CI is (0.013, 0.150) (c.f., t -based CI of (-0.002, 0.150)).

¹e.g., Robbins-Monroe search - see accompanying notes



Outline

- 1 Introduction
 - Revision – Traditional parametric and nonparametric methods
 - Motivating example
- 2 **Permutation methods**
 - Permutation test
 - Permutation interval
 - **Summary**
- 3 Randomization methods
 - Overview
 - Summary
- 4 Monte-Carlo tests
 - Overview
 - Summary



Summary

Pros

- Widely applicable
- Produces exact results
- No specific distribution assumed for the data
- Does not require analytic distribution of test statistic



Summary

Pros

- Widely applicable
- Produces exact results
- No specific distribution assumed for the data
- Does not require analytic distribution of test statistic



Summary

Pros

- Widely applicable
- Produces exact results
- No specific distribution assumed for the data
- Does not require analytic distribution of test statistic



Summary

Pros

- Widely applicable
- Produces exact results
- No specific distribution assumed for the data
- Does not require analytic distribution of test statistic



Cons

- Each permutation of the data must be equally likely (or probability of occurrence is known)
- To be generally applicable, requires thought, and ability to program a computer (permutation algorithms can be non-trivial)
- Can be prohibitively computer (and programmer) intensive in complex situations or for large sample sizes



Cons

- Each permutation of the data must be equally likely (or probability of occurrence is known)
- To be generally applicable, requires thought, and ability to program a computer (permutation algorithms can be non-trivial)
- Can be prohibitively computer (and programmer) intensive in complex situations or for large sample sizes



Cons

- Each permutation of the data must be equally likely (or probability of occurrence is known)
- To be generally applicable, requires thought, and ability to program a computer (permutation algorithms can be non-trivial)
- Can be prohibitively computer (and programmer) intensive in complex situations or for large sample sizes



Outline

- 1 Introduction
 - Revision – Traditional parametric and nonparametric methods
 - Motivating example
- 2 Permutation methods
 - Permutation test
 - Permutation interval
 - Summary
- 3 Randomization methods
 - Overview
 - Summary
- 4 Monte-Carlo tests
 - Overview
 - Summary



Introduction

- Randomization methods are like permutation methods, except that a only random subset of all the possible permutations are generated
- E.g., pine martin example:
 - To generate one randomization, list the set of values $|x_i - \mu_0|$ and randomly assign a sign to each value
 - From one run of 999 randomizations plus the original data value, I obtained $p = 0.015$, very similar to the $p = 0.012$ of the permutation method.
- Confidence intervals can be obtained in a similar way to permutation intervals

Introduction

- Randomization methods are like permutation methods, except that a only random subset of all the possible permutations are generated
- E.g., pine martin example:
 - To generate one randomization, list the set of values $|x_i - \mu_0|$ and randomly assign a sign to each value
 - From one run of 999 randomizations plus the original data value, I obtained $p = 0.015$, very similar to the $p = 0.012$ of the permutation method.
- Confidence intervals can be obtained in a similar way to permutation intervals

Introduction

- Randomization methods are like permutation methods, except that a only random subset of all the possible permutations are generated
- E.g., pine martin example:
 - To generate one randomization, list the set of values $|x_i - \mu_0|$ and randomly assign a sign to each value
 - From one run of 999 randomizations plus the original data value, I obtained $p = 0.015$, very similar to the $p = 0.012$ of the permutation method.
- Confidence intervals can be obtained in a similar way to permutation intervals

Introduction

- Randomization methods are like permutation methods, except that a only random subset of all the possible permutations are generated
- E.g., pine martin example:
 - To generate one randomization, list the set of values $|x_i - \mu_0|$ and randomly assign a sign to each value
 - From one run of 999 randomizations plus the original data value, I obtained $p = 0.015$, very similar to the $p = 0.012$ of the permutation method.
- Confidence intervals can be obtained in a similar way to permutation intervals

Introduction

- Randomization methods are like permutation methods, except that a only random subset of all the possible permutations are generated
- E.g., pine martin example:
 - To generate one randomization, list the set of values $|x_i - \mu_0|$ and randomly assign a sign to each value
 - From one run of 999 randomizations plus the original data value, I obtained $p = 0.015$, very similar to the $p = 0.012$ of the permutation method.
- Confidence intervals can be obtained in a similar way to permutation intervals

Outline

- 1 Introduction
 - Revision – Traditional parametric and nonparametric methods
 - Motivating example
- 2 Permutation methods
 - Permutation test
 - Permutation interval
 - Summary
- 3 Randomization methods
 - Overview
 - **Summary**
- 4 Monte-Carlo tests
 - Overview
 - Summary



Summary

Pros, compared with permutation methods

- All of the same advantages as permutation, plus
- Much easier to code
- Number of randomizations is fixed, rather than being a function of sample size



Summary

Pros, compared with permutation methods

- All of the same advantages as permutation, plus
- Much easier to code
- Number of randomizations is fixed, rather than being a function of sample size



Summary

Pros, compared with permutation methods

- All of the same advantages as permutation, plus
- Much easier to code
- Number of randomizations is fixed, rather than being a function of sample size



Summary

Cons, compared with permutation methods

- As with permutation tests, each randomization of the data must be equally likely (or probability of occurrence is known)
- Results will vary from run to run
 - Size of this "Monte-Carlo variation" depends on the number of randomizations



Summary

Cons, compared with permutation methods

- As with permutation tests, each randomization of the data must be equally likely (or probability of occurrence is known)
- Results will vary from run to run
 - Size of this “Monte-Carlo variation” depends on the number of randomizations

Summary

Cons, compared with permutation methods

- As with permutation tests, each randomization of the data must be equally likely (or probability of occurrence is known)
- Results will vary from run to run
 - Size of this “Monte-Carlo variation” depends on the number of randomizations



Outline

- 1 Introduction
 - Revision – Traditional parametric and nonparametric methods
 - Motivating example
- 2 Permutation methods
 - Permutation test
 - Permutation interval
 - Summary
- 3 Randomization methods
 - Overview
 - Summary
- 4 Monte-Carlo tests
 - Overview
 - Summary



Introduction

A generalization² of randomization tests to include parametric distributions for the data.

- Algorithm:

- 1 Set up some distribution for the data, $f()$ or $f(\theta)$, and some H_0
- 2 Repeat the following two steps many times:
 - 1 Simulate a data set according to H_0
 - 2 Calculate $T(X)$ using the simulated data
- 3 Add $T(x)$ evaluated from the sample data
- 4 Order all of the $T(X)$ s
- 5 p -value is the proportion of the $T(X)$ s as extreme or more extreme than the one from the sample data

²If the assumed model for $f()$ implies that all data orderings are equally likely then you get back to a randomization test

Introduction

A generalization² of randomization tests to include parametric distributions for the data.

- Algorithm:

- 1 Set up some distribution for the data, $f()$ or $f(\theta)$, and some H_0
- 2 Repeat the following two steps many times:
 - Simulate a data set according to H_0
 - Calculate $T(X)$ using the simulated data
- 3 Add $T(x)$ evaluated from the sample data
- 4 Order all of the $T(X)$ s
- 5 p -value is the proportion of the $T(X)$ s as extreme or more extreme than the one from the sample data

²If the assumed model for $f()$ implies that all data orderings are equally likely then you get back to a randomization test

Introduction

A generalization² of randomization tests to include parametric distributions for the data.

- Algorithm:

- 1 Set up some distribution for the data, $f()$ or $f(\theta)$, and some H_0
- 2 Repeat the following two steps many times:
 - 1 Simulate a data set according to H_0
 - 2 Calculate $T(X)$ using the simulated data
- 3 Add $T(x)$ evaluated from the sample data
- 4 Order all of the $T(X)$ s
- 5 p -value is the proportion of the $T(X)$ s as extreme or more extreme than the one from the sample data

²If the assumed model for $f()$ implies that all data orderings are equally likely then you get back to a randomization test

Introduction

A generalization² of randomization tests to include parametric distributions for the data.

- Algorithm:

- 1 Set up some distribution for the data, $f()$ or $f(\theta)$, and some H_0
- 2 Repeat the following two steps many times:
 - 1 Simulate a data set according to H_0
 - 2 Calculate $T(X)$ using the simulated data
- 3 Add $T(x)$ evaluated from the sample data
- 4 Order all of the $T(X)$ s
- 5 p -value is the proportion of the $T(X)$ s as extreme or more extreme than the one from the sample data

²If the assumed model for $f()$ implies that all data orderings are equally likely then you get back to a randomization test

Introduction

A generalization² of randomization tests to include parametric distributions for the data.

- Algorithm:

- 1 Set up some distribution for the data, $f()$ or $f(\theta)$, and some H_0
- 2 Repeat the following two steps many times:
 - 1 Simulate a data set according to H_0
 - 2 Calculate $T(X)$ using the simulated data
- 3 Add $T(x)$ evaluated from the sample data
- 4 Order all of the $T(X)$ s
- 5 p -value is the proportion of the $T(X)$ s as extreme or more extreme than the one from the sample data

²If the assumed model for $f()$ implies that all data orderings are equally likely then you get back to a randomization test

Introduction

A generalization² of randomization tests to include parametric distributions for the data.

- Algorithm:

- 1 Set up some distribution for the data, $f()$ or $f(\theta)$, and some H_0
- 2 Repeat the following two steps many times:
 - 1 Simulate a data set according to H_0
 - 2 Calculate $T(X)$ using the simulated data
- 3 Add $T(x)$ evaluated from the sample data
- 4 Order all of the $T(X)$ s
- 5 p -value is the proportion of the $T(X)$ s as extreme or more extreme than the one from the sample data

²If the assumed model for $f()$ implies that all data orderings are equally likely then you get back to a randomization test

Introduction

A generalization² of randomization tests to include parametric distributions for the data.

- Algorithm:

- 1 Set up some distribution for the data, $f()$ or $f(\theta)$, and some H_0
- 2 Repeat the following two steps many times:
 - 1 Simulate a data set according to H_0
 - 2 Calculate $T(X)$ using the simulated data
- 3 Add $T(x)$ evaluated from the sample data
- 4 Order all of the $T(X)$ s
- 5 p -value is the proportion of the $T(X)$ s as extreme or more extreme than the one from the sample data

²If the assumed model for $f()$ implies that all data orderings are equally likely then you get back to a randomization test

Introduction

A generalization² of randomization tests to include parametric distributions for the data.

- Algorithm:

- 1 Set up some distribution for the data, $f()$ or $f(\theta)$, and some H_0
- 2 Repeat the following two steps many times:
 - 1 Simulate a data set according to H_0
 - 2 Calculate $T(X)$ using the simulated data
- 3 Add $T(x)$ evaluated from the sample data
- 4 Order all of the $T(X)$ s
- 5 p -value is the proportion of the $T(X)$ s as extreme or more extreme than the one from the sample data

²If the assumed model for $f()$ implies that all data orderings are equally likely then you get back to a randomization test

Pine martin example - parametric simulation

- Assume observations are iid $N(\mu, \sigma^2)$, that $H_0 : \mu = 0$, and that $\sigma^2 = s^2$
- Simulate 999 data sets of 12 observations from $N(0, s^2)$.
- Calculate \bar{x} for each of these, and add in the mean from the sample data, to give 1000 values of \bar{x} .
- p -value is the proportion of these values that are more extreme (e.g., $p = 0.029$)



Pine martin example - parametric simulation

- Assume observations are iid $N(\mu, \sigma^2)$, that $H_0 : \mu = 0$, and that $\sigma^2 = s^2$
- Simulate 999 data sets of 12 observations from $N(0, s^2)$.
- Calculate \bar{x} for each of these, and add in the mean from the sample data, to give 1000 values of \bar{x} .
- p -value is the proportion of these values that are more extreme (e.g., $p = 0.029$)




Pine martin example - parametric simulation

- Assume observations are iid $N(\mu, \sigma^2)$, that $H_0 : \mu = 0$, and that $\sigma^2 = s^2$
- Simulate 999 data sets of 12 observations from $N(0, s^2)$.
- Calculate \bar{x} for each of these, and add in the mean from the sample data, to give 1000 values of \bar{x} .
- p -value is the proportion of these values that are more extreme (e.g., $p = 0.029$)



Pine martin example - parametric simulation

- Assume observations are iid $N(\mu, \sigma^2)$, that $H_0 : \mu = 0$, and that $\sigma^2 = s^2$
- Simulate 999 data sets of 12 observations from $N(0, s^2)$.
- Calculate \bar{x} for each of these, and add in the mean from the sample data, to give 1000 values of \bar{x} .
- p -value is the proportion of these values that are more extreme (e.g., $p = 0.029$)



Outline

- 1 Introduction
 - Revision – Traditional parametric and nonparametric methods
 - Motivating example
- 2 Permutation methods
 - Permutation test
 - Permutation interval
 - Summary
- 3 Randomization methods
 - Overview
 - Summary
- 4 Monte-Carlo tests
 - Overview
 - Summary



Summary

Pros

- Data can be assumed to follow any distribution - parametric or non-parametric (Randomization tests are a special case)
- The analytic distribution of the test statistic is not required



Summary

Pros

- Data can be assumed to follow any distribution - parametric or non-parametric (Randomization tests are a special case)
- The analytic distribution of the test statistic is not required



Summary

Cons, compared with permutation methods

- For parametric simulations, need to estimate nuisance parameters (e.g., σ^2 in the pine martin example)
- Results will vary from run to run



Summary

Cons, compared with permutation methods

- For parametric simulations, need to estimate nuisance parameters (e.g., σ^2 in the pine martin example)
- Results will vary from run to run



Summary

Monte Carlo confidence intervals

Confidence intervals can be constructed using similar methods to randomization intervals – but there is a better way – see next lecture...

