



The fiber-full scheme

Yairon Cid-Ruiz
Ghent University

(joint work with Ritvik Ramkumar)

Fellowship of the Ring seminar
October 12th, 2021

arXiv: <https://arxiv.org/abs/2108.13986>

slides: <https://ycid.github.io/SlidesFellowshipRing.pdf>

Goals & Plan

- 1 Introduce and motivate the construction of a new parameter space: “*the fiber-full scheme*”. Guiding goals:

Goals & Plan

- 1 Introduce and motivate the construction of a new parameter space: “*the fiber-full scheme*”. Guiding goals:
 - We want to parametrize closed subschemes $Z \subset \mathbb{P}_{\mathbb{k}}^r$ by fixing the “*entire cohomological data*” and not just the Hilbert polynomial.

Goals & Plan

- ① Introduce and motivate the construction of a new parameter space: “*the fiber-full scheme*”. Guiding goals:
 - We want to parametrize closed subschemes $Z \subset \mathbb{P}_{\mathbb{k}}^r$ by fixing the “*entire cohomological data*” and not just the Hilbert polynomial.
 - We want a stratification of the Hilbert schemes in terms of these new fiber-full schemes.

Goals & Plan

- ① Introduce and motivate the construction of a new parameter space: “*the fiber-full scheme*”. Guiding goals:
 - We want to parametrize closed subschemes $Z \subset \mathbb{P}_{\mathbb{k}}^r$ by fixing the “*entire cohomological data*” and not just the Hilbert polynomial.
 - We want a stratification of the Hilbert schemes in terms of these new fiber-full schemes.
- ② Review some applications that stem from the existence of the full fiber scheme.

Goals & Plan

- ➊ Introduce and motivate the construction of a new parameter space: “*the fiber-full scheme*”. Guiding goals:
 - We want to parametrize closed subschemes $Z \subset \mathbb{P}_{\mathbb{k}}^r$ by fixing the “*entire cohomological data*” and not just the Hilbert polynomial.
 - We want a stratification of the Hilbert schemes in terms of these new fiber-full schemes.
- ➋ Review some applications that stem from the existence of the full fiber scheme.
- ➌ From a technical side: results on sheaf/local cohomology and a flattening stratification theorem.

A a Noetherian commutative ring and $S = \operatorname{Spec}(A)$. $R = A[x_0, \dots, x_r]$ a standard graded commutative ring and $\mathbb{P}_A^r = \operatorname{Proj}(R)$. $P \in \mathbb{Q}[m]$ a numerical polynomial.

Hilbert scheme (Grothendieck, 1961)

$\operatorname{Hilb}_{\mathbb{P}_A^r}^P$ parametrizes closed subschemes $Z \subset \mathbb{P}_A^r$ with Hilbert polynomial P :

$\{\text{closed } Z \subset \mathbb{P}_A^r \mid Z \text{ is flat over } S \text{ and } Z_{\mathfrak{p}} \text{ has Hilbert polynomial } P \text{ for all } \mathfrak{p} \in S\}$,

which is the same as

$$\left\{ Z = \operatorname{Proj}(R/I) \subset \mathbb{P}_A^r \left| \begin{array}{l} I \subset R \text{ homogeneous ideal,} \\ I = I : (x_0, \dots, x_r)^\infty, \\ [R/I]_\nu \text{ is a locally free } A\text{-module} \\ \text{of constant rank } P(\nu) \text{ for all } \nu \gg 0 \end{array} \right. \right\}.$$

A a Noetherian commutative ring and $S = \operatorname{Spec}(A)$. $R = A[x_0, \dots, x_r]$ a standard graded commutative ring and $\mathbb{P}_A^r = \operatorname{Proj}(R)$. $P \in \mathbb{Q}[m]$ a numerical polynomial.

Hilbert scheme (Grothendieck, 1961)

$\operatorname{Hilb}_{\mathbb{P}_A^r}^P$ parametrizes closed subschemes $Z \subset \mathbb{P}_A^r$ with Hilbert polynomial P :

$\{\text{closed } Z \subset \mathbb{P}_A^r \mid Z \text{ is flat over } S \text{ and } Z_{\mathfrak{p}} \text{ has Hilbert polynomial } P \text{ for all } \mathfrak{p} \in S\}$,

which is the same as

$$\left\{ Z = \operatorname{Proj}(R/I) \subset \mathbb{P}_A^r \left| \begin{array}{l} I \subset R \text{ homogeneous ideal,} \\ I = I : (x_0, \dots, x_r)^\infty, \\ [R/I]_\nu \text{ is a locally free } A\text{-module} \\ \text{of constant rank } P(\nu) \text{ for all } \nu \gg 0 \end{array} \right. \right\}.$$

So, if $[Z] = [\operatorname{Proj}(R/I)] \in \operatorname{Hilb}_{\mathbb{P}_A^r}^P$, then $R/I \otimes_A \kappa(\mathfrak{p})$ has Hilbert polynomial P for all $\mathfrak{p} \in S$, where $\kappa(\mathfrak{p}) = A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}}$.

Example: Twisted cubics (Piene - Schlessinger, 1985)

A *twisted cubic* $C \subset \mathbb{P}_{\mathbb{k}}^3$ (\mathbb{k} alg. closed and $\text{char}(\mathbb{k}) = 0$) is a rational smooth curve of degree 3. Any such C is projectively equivalent to $C_0 = \phi(\mathbb{P}_{\mathbb{k}}^1)$ where $\phi : \mathbb{P}_{\mathbb{k}}^1 \rightarrow \mathbb{P}_{\mathbb{k}}^3$, $(u : v) \mapsto (u^3 : u^2v : uv^2 : v^3)$.

Theorem (Piene - Schlessinger, 1985)

$\text{Hilb}_{\mathbb{P}_{\mathbb{k}}^3}^{3m+1} = H \cup H'$ (two smooth irreducible components), where H parametrizes twisted cubics and H' parametrizes a plane cubic union an isolated point.

Remark

With the Hilbert scheme compactification of the space of twisted cubics we obtain the “extraneous” component H' .

$$\text{If } [Z] \in H - H \cap H', \text{ then } h^0(Z, \mathcal{O}_Z(\nu)) = \begin{cases} 3\nu + 1 & \text{if } \nu \geq 0 \\ 0 & \text{if } \nu \leq -1 \end{cases}$$

$$\text{and } h^1(Z, \mathcal{O}_Z(j)) = h^0(Z, \mathcal{O}_Z(\nu)) - (3\nu + 1).$$

$$\text{If } [Z] \in H', \text{ then } h^0(Z, \mathcal{O}_Z(\nu)) = \begin{cases} 3\nu + 1 & \text{if } \nu \geq 1 \\ 2 & \text{if } \nu = 0 \\ 1 & \text{if } \nu \leq -1 \end{cases}$$

$$\text{and } h^1(Z, \mathcal{O}_Z(j)) = h^0(Z, \mathcal{O}_Z(\nu)) - (3\nu + 1).$$

Stratification of in cohomological terms

$\text{Hilb}_{\mathbb{P}^r_{\mathbb{k}}}^{3m+1} = (H - H \cap H') \cup H'$. Thus, $h_Z^i : \mathbb{Z} \rightarrow \mathbb{N}$, $h_Z^i(\nu) := \dim_{\mathbb{k}} (H^i(Z, \mathcal{O}_Z(\nu)))$ is the same for any $[Z] \in H - H \cap H'$ and is the same for any $[Z] \in H'$.

Guiding goals

(1) Can we do this kind of cohomological stratification for any Hilbert scheme in terms of **locally closed subschemes**? (2) If so, we want to provide a unified and systematic treatment.

Formal definition of Hilbert and Quot schemes

$S = \operatorname{Spec}(A)$ with A Noetherian, $X \subset \mathbb{P}_A^r$ closed subscheme, \mathcal{F} coherent sheaf on X . For any S -scheme $T = \operatorname{Spec}(B)$, let $\mathcal{F}_T = \mathcal{F} \otimes_A B$. $P \in \mathbb{Q}[m]$ numerical polynomial.

Quot functor $\underline{\operatorname{Quot}}_{\mathcal{F}/X/S}^P$

$$\underline{\operatorname{Quot}}_{\mathcal{F}/X/S}^P(T) = \left\{ \text{coherent } \mathcal{F}_T \twoheadrightarrow \mathcal{G} \mid \begin{array}{l} \mathcal{G} \text{ if flat over } T \text{ and } G_t \text{ has} \\ \text{Hilbert polynomial } P \text{ for all } t \in T \end{array} \right\}.$$

Theorem (Grothendieck, 1961)

$\underline{\operatorname{Quot}}_{\mathcal{F}/X/S}^P$ is represented by a projective S -scheme $\operatorname{Quot}_{\mathcal{F}/X/S}^P$.

Formal definition of Hilbert and Quot schemes

$S = \operatorname{Spec}(A)$ with A Noetherian, $X \subset \mathbb{P}_A^r$ closed subscheme, \mathcal{F} coherent sheaf on X . For any S -scheme $T = \operatorname{Spec}(B)$, let $\mathcal{F}_T = \mathcal{F} \otimes_A B$. $P \in \mathbb{Q}[m]$ numerical polynomial.

Quot functor $\underline{\operatorname{Quot}}_{\mathcal{F}/X/S}^P$

$$\underline{\operatorname{Quot}}_{\mathcal{F}/X/S}^P(T) = \left\{ \text{coherent } \mathcal{F}_T \twoheadrightarrow \mathcal{G} \mid \begin{array}{l} \mathcal{G} \text{ if flat over } T \text{ and } G_t \text{ has} \\ \text{Hilbert polynomial } P \text{ for all } t \in T \end{array} \right\}.$$

Theorem (Grothendieck, 1961)

$\underline{\operatorname{Quot}}_{\mathcal{F}/X/S}^P$ is represented by a projective S -scheme $\operatorname{Quot}_{\mathcal{F}/X/S}^P$.

There is a **universal sheaf** $\mathcal{W}_{\mathcal{F}/X/S}^P \in \underline{\operatorname{Quot}}_{\mathcal{F}/X/S}^P(\operatorname{Quot}_{\mathcal{F}/X/S}^P)$ such that for any $\mathcal{G} \in \underline{\operatorname{Quot}}_{\mathcal{F}/X/S}^P(T)$ there is a **unique classifying S -morphism**

$$g_{\mathcal{G}} : T \rightarrow \operatorname{Quot}_{\mathcal{F}/X/S}^P \text{ such that } \mathcal{G} = \left(\mathcal{W}_{\mathcal{F}/X/S}^P \right)_T = (1_X \times_S g_{\mathcal{G}})^* \mathcal{W}_{\mathcal{F}/X/S}^P.$$

The functor we need to study

$S = \operatorname{Spec}(A)$ with A Noetherian, $X \subset \mathbb{P}_A^r$ closed subscheme, \mathcal{F} coherent sheaf on X . For any S -scheme $T = \operatorname{Spec}(B)$, let $\mathcal{F}_T = \mathcal{F} \otimes_A B$.

Let $\mathbf{h} = (h_0, \dots, h_r) : \mathbb{Z}^{r+1} \rightarrow \mathbb{N}^{r+1}$ be a tuple of functions.

Defintion: the fiber-full functor $\underline{\mathcal{F}ib}_{\mathcal{F}/X/S}^{\mathbf{h}}$

$$\underline{\mathcal{F}ib}_{\mathcal{F}/X/S}^{\mathbf{h}}(T) = \left\{ \begin{array}{l} \text{coherent } \mathcal{F}_T \twoheadrightarrow \mathcal{G} \quad \left| \quad \begin{array}{l} H^i(X_T, \mathcal{G}(\nu)) \text{ is a locally free } B\text{-module} \\ \text{of constant rank equal to } h_i(\nu) \\ \text{for all } 0 \leq i \leq r, \nu \in \mathbb{Z} \end{array} \right. \end{array} \right\}$$

The functor we need to study

$S = \operatorname{Spec}(A)$ with A Noetherian, $X \subset \mathbb{P}_A^r$ closed subscheme, \mathcal{F} coherent sheaf on X . For any S -scheme $T = \operatorname{Spec}(B)$, let $\mathcal{F}_T = \mathcal{F} \otimes_A B$.

Let $\mathbf{h} = (h_0, \dots, h_r) : \mathbb{Z}^{r+1} \rightarrow \mathbb{N}^{r+1}$ be a tuple of functions.

Defintion: the fiber-full functor $\underline{\operatorname{Fib}}_{\mathcal{F}/X/S}^{\mathbf{h}}$

$$\underline{\operatorname{Fib}}_{\mathcal{F}/X/S}^{\mathbf{h}}(T) = \left\{ \begin{array}{l} \text{coherent } \mathcal{F}_T \twoheadrightarrow \mathcal{G} \quad \left| \quad \begin{array}{l} H^i(X_T, \mathcal{G}(\nu)) \text{ is a locally free } B\text{-module} \\ \text{of constant rank equal to } h_i(\nu) \\ \text{for all } 0 \leq i \leq r, \nu \in \mathbb{Z} \end{array} \right. \end{array} \right\}$$

It is a functor because of the following base change result.

Lemma

Assume \mathcal{F} is S -flat and $H^i(X, \mathcal{F})$ are A -flat for all $0 \leq i \leq r$. Then

$$H^i(X, \mathcal{F}) \otimes_A B \xrightarrow{\cong} H^i(X_T, \mathcal{F}_T) \text{ for all } 0 \leq i \leq r.$$

In particular, all $H^i(X_T, \mathcal{F}_T)$ are B -flat.

- \mathcal{F} is S -flat $\iff H^0(X, \mathcal{F}(\nu))$ is A -flat for all $\nu \gg 0$.
- The Hilbert polynomial coincides with the Euler characteristic.

Relation between $\underline{\mathcal{F}ib}_{\mathcal{F}/X/S}^P$ and $\underline{Quot}_{\mathcal{F}/X/S}^P$

Let $P_{\mathbf{h}} = \sum_{i=0}^r (-1)^i h_i$. For any S -scheme $T = \text{Spec}(B)$, we have the inclusion

$$\underline{\mathcal{F}ib}_{\mathcal{F}/X/S}^P(T) \subset \underline{Quot}_{\mathcal{F}/X/S}^P(T).$$

Therefore, $\underline{\mathcal{F}ib}_{\mathcal{F}/X/S}^P$ is a **subfunctor** of $\underline{Quot}_{\mathcal{F}/X/S}^P$.

- \mathcal{F} is S -flat $\iff H^0(X, \mathcal{F}(\nu))$ is A -flat for all $\nu \gg 0$.
- The Hilbert polynomial coincides with the Euler characteristic.

Relation between $\underline{\text{Fib}}^P_{\mathcal{F}/X/S}$ and $\underline{\text{Quot}}^P_{\mathcal{F}/X/S}$

Let $P_h = \sum_{i=0}^r (-1)^i h_i$. For any S -scheme $T = \text{Spec}(B)$, we have the inclusion

$$\underline{\text{Fib}}^P_{\mathcal{F}/X/S}(T) \subset \underline{\text{Quot}}^P_{\mathcal{F}/X/S}(T).$$

Therefore, $\underline{\text{Fib}}^P_{\mathcal{F}/X/S}$ is a **subfunctor** of $\underline{\text{Quot}}^P_{\mathcal{F}/X/S}$.

Our main question!

- Is the fiber-full functor $\underline{\text{Fib}}^P_{\mathcal{F}/X/S}$ representable?
- If so, its representing scheme would grant us all our objectives. This scheme would control the entire cohomological data.

First, let us address the following question:

When is $H^i(X, \mathcal{F})$ a flat A -module?

First, let us address the following question:

When is $H^i(X, \mathcal{F})$ a flat A -module?

$S = \operatorname{Spec}(A)$, $X = \operatorname{Proj}(R) \subset \mathbb{P}_A^r$ closed subscheme, \mathcal{F} coherent sheaf on X , $\mathfrak{m} = [R]_+ \subset R$, and M a finitely generated graded R -module such that $\mathcal{F} = \tilde{M}$.

Reminder

$0 \rightarrow H_{\mathfrak{m}}^0(M) \rightarrow M \rightarrow \bigoplus_{\nu \in \mathbb{Z}} H^0(X, \mathcal{F}(\nu)) \rightarrow H_{\mathfrak{m}}^1(M) \rightarrow 0$ is exact, and $H_{\mathfrak{m}}^{i+1}(M) \cong \bigoplus_{\nu \in \mathbb{Z}} H^i(X, \mathcal{F}(\nu))$ for $i \geq 1$.

First, let us address the following question:

When is $H^i(X, \mathcal{F})$ a flat A -module?

$S = \operatorname{Spec}(A)$, $X = \operatorname{Proj}(R) \subset \mathbb{P}_A^r$ closed subscheme, \mathcal{F} coherent sheaf on X , $\mathfrak{m} = [R]_+ \subset R$, and M a finitely generated graded R -module such that $\mathcal{F} = \tilde{M}$.

Reminder

$0 \rightarrow H_{\mathfrak{m}}^0(M) \rightarrow M \rightarrow \bigoplus_{\nu \in \mathbb{Z}} H^0(X, \mathcal{F}(\nu)) \rightarrow H_{\mathfrak{m}}^1(M) \rightarrow 0$ is exact, and $H_{\mathfrak{m}}^{i+1}(M) \cong \bigoplus_{\nu \in \mathbb{Z}} H^i(X, \mathcal{F}(\nu))$ for $i \geq 1$.

It is then equivalent to address the A -flatness of all $H_{\mathfrak{m}}^i(M)$. This problem has been studied before: **Hochster-Roberts** (1976), **Kollár** (2014), **Smith** (2018), **Chardin-CR-Simis** (2020).

Theorem (Hochster-Roberts, 1976)

Assume A is a domain. There exists $0 \neq a \in A$ such that $H_{\mathfrak{m}}^i(M \otimes_A A_a)$ is a locally free A_a -module for all i .

Fiber-full modules

Motivated by work of **Kollár-Kovács** on the flatness of the cohomologies of a relative dualizing complex (also of **Dao-De Stefani-Ma**), Varbaro obtained the following:

Theorem (Varbaro, 2021)

Let $A = \mathbb{k}[t]$, R a fin. gen. A -algebra, M a fin. gen. R -module. Assume M is A -flat and the natural map $\mathrm{Ext}_R^i(M/tM, R) \rightarrow \mathrm{Ext}_R^i(M/t^q M, R)$ is injective $\forall i, q$. Then, $\mathrm{Ext}_R^i(M, R)$ is flat over $A \forall i$.

Fiber-full modules

Motivated by work of **Kollár-Kovács** on the flatness of the cohomologies of a relative dualizing complex (also of **Dao-De Stefani-Ma**), Varbaro obtained the following:

Theorem (Varbaro, 2021)

Let $A = \mathbb{k}[t]$, R a fin. gen. A -algebra, M a fin. gen. R -module. Assume M is A -flat and the natural map $\mathrm{Ext}_R^i(M/tM, R) \rightarrow \mathrm{Ext}_R^i(M/t^q M, R)$ is injective $\forall i, q$. Then, $\mathrm{Ext}_R^i(M, R)$ is flat over $A \forall i$.

We want to address the following general setup:

(B, \mathfrak{b}) a Noetherian local ring, R fin. gen. pos. graded B -algebra. M a fin. gen. graded R -module.

Definition

We say M is **fiber-full over B** if M is B -free and the natural map

$$H_m^i(M/\mathfrak{b}^q M) \rightarrow H_m^i(M/\mathfrak{b} M) \text{ is surjective } \forall i, q.$$

(B, \mathfrak{b}) a Noetherian local ring, R fin. gen. pos. graded B -algebra. M a finitely generated graded R -module. $T = B[x_1, \dots, x_r]$ pos. graded with $T \twoheadrightarrow R$.

Theorem (CR)

Assume M is a free B -module. The following six conditions are equivalent:

- 1 $H_m^i(M)$ is a free B -module $\forall 0 \leq i \leq r$.

(B, \mathfrak{b}) a Noetherian local ring, R fin. gen. pos. graded B -algebra. M a finitely generated graded R -module. $T = B[x_1, \dots, x_r]$ pos. graded with $T \twoheadrightarrow R$.

Theorem (CR)

Assume M is a free B -module. The following six conditions are equivalent:

- ① $H_{\mathfrak{m}}^i(M)$ is a free B -module $\forall 0 \leq i \leq r$.
- ② $\text{Ext}_T^i(M, T)$ is a free B -module $\forall 0 \leq i \leq r$.

(B, \mathfrak{b}) a Noetherian local ring, R fin. gen. pos. graded B -algebra. M a finitely generated graded R -module. $T = B[x_1, \dots, x_r]$ pos. graded with $T \twoheadrightarrow R$.

Theorem (CR)

Assume M is a free B -module. The following six conditions are equivalent:

- ① $H_{\mathfrak{m}}^i(M)$ is a free B -module $\forall 0 \leq i \leq r$.
- ② $\text{Ext}_T^i(M, T)$ is a free B -module $\forall 0 \leq i \leq r$.
- ③ $H_{\mathfrak{m}}^i(M/\mathfrak{b}^q M)$ is a free B/\mathfrak{b}^q -module $\forall 0 \leq i \leq r, q \geq 1$.

(B, \mathfrak{b}) a Noetherian local ring, R fin. gen. pos. graded B -algebra. M a finitely generated graded R -module. $T = B[x_1, \dots, x_r]$ pos. graded with $T \twoheadrightarrow R$.

Theorem (CR)

Assume M is a free B -module. The following six conditions are equivalent:

- ① $H_m^i(M)$ is a free B -module $\forall 0 \leq i \leq r$.
- ② $\text{Ext}_T^i(M, T)$ is a free B -module $\forall 0 \leq i \leq r$.
- ③ $H_m^i(M/\mathfrak{b}^q M)$ is a free B/\mathfrak{b}^q -module $\forall 0 \leq i \leq r, q \geq 1$.
- ④ $\text{Ext}_{T/\mathfrak{b}^q T}^i(M/\mathfrak{b}^q M, T/\mathfrak{b}^q T)$ is a free B/\mathfrak{b}^q -module $\forall 0 \leq i \leq r, q \geq 1$.

(B, \mathfrak{b}) a Noetherian local ring, R fin. gen. pos. graded B -algebra. M a finitely generated graded R -module. $T = B[x_1, \dots, x_r]$ pos. graded with $T \twoheadrightarrow R$.

Theorem (CR)

Assume M is a free B -module. The following six conditions are equivalent:

- ① $H_m^i(M)$ is a free B -module $\forall 0 \leq i \leq r$.
- ② $\text{Ext}_T^i(M, T)$ is a free B -module $\forall 0 \leq i \leq r$.
- ③ $H_m^i(M/\mathfrak{b}^q M)$ is a free B/\mathfrak{b}^q -module $\forall 0 \leq i \leq r, q \geq 1$.
- ④ $\text{Ext}_{T/\mathfrak{b}^q T}^i(M/\mathfrak{b}^q M, T/\mathfrak{b}^q T)$ is a free B/\mathfrak{b}^q -module $\forall 0 \leq i \leq r, q \geq 1$.
- ⑤ $H_m^i(M/\mathfrak{b}^q M) \rightarrow H_m^i(M/\mathfrak{b} M)$ is surjective $\forall 0 \leq i \leq r, q \geq 1$.

(B, \mathfrak{b}) a Noetherian local ring, R fin. gen. pos. graded B -algebra. M a finitely generated graded R -module. $T = B[x_1, \dots, x_r]$ pos. graded with $T \twoheadrightarrow R$.

Theorem (CR)

Assume M is a free B -module. The following six conditions are equivalent:

- ① $H_m^i(M)$ is a free B -module $\forall 0 \leq i \leq r$.
- ② $\text{Ext}_T^i(M, T)$ is a free B -module $\forall 0 \leq i \leq r$.
- ③ $H_m^i(M/\mathfrak{b}^q M)$ is a free B/\mathfrak{b}^q -module $\forall 0 \leq i \leq r, q \geq 1$.
- ④ $\text{Ext}_{T/\mathfrak{b}^q T}^i(M/\mathfrak{b}^q M, T/\mathfrak{b}^q T)$ is a free B/\mathfrak{b}^q -module $\forall 0 \leq i \leq r, q \geq 1$.
- ⑤ $H_m^i(M/\mathfrak{b}^q M) \rightarrow H_m^i(M/\mathfrak{b} M)$ is surjective $\forall 0 \leq i \leq r, q \geq 1$.
- ⑥ $\text{Ext}_{T/\mathfrak{b}^q T}^i(M/\mathfrak{b} M, \omega_{T/\mathfrak{b}^q T}) \rightarrow \text{Ext}_{T/\mathfrak{b}^q T}^i(M/\mathfrak{b}^q M, \omega_{T/\mathfrak{b}^q T})$ is injective $\forall 0 \leq i \leq r, q \geq 1$.

(B, \mathfrak{b}) a Noetherian local ring, R fin. gen. pos. graded B -algebra. M a finitely generated graded R -module. $T = B[x_1, \dots, x_r]$ pos. graded with $T \twoheadrightarrow R$.

Theorem (CR)

Assume M is a free B -module. The following six conditions are equivalent:

- ① $H_m^i(M)$ is a free B -module $\forall 0 \leq i \leq r$.
- ② $\text{Ext}_T^i(M, T)$ is a free B -module $\forall 0 \leq i \leq r$.
- ③ $H_m^i(M/\mathfrak{b}^q M)$ is a free B/\mathfrak{b}^q -module $\forall 0 \leq i \leq r, q \geq 1$.
- ④ $\text{Ext}_{T/\mathfrak{b}^q T}^i(M/\mathfrak{b}^q M, T/\mathfrak{b}^q T)$ is a free B/\mathfrak{b}^q -module $\forall 0 \leq i \leq r, q \geq 1$.
- ⑤ $H_m^i(M/\mathfrak{b}^q M) \rightarrow H_m^i(M/\mathfrak{b} M)$ is surjective $\forall 0 \leq i \leq r, q \geq 1$.
- ⑥ $\text{Ext}_{T/\mathfrak{b}^q T}^i(M/\mathfrak{b} M, \omega_{T/\mathfrak{b}^q T}) \rightarrow \text{Ext}_{T/\mathfrak{b}^q T}^i(M/\mathfrak{b}^q M, \omega_{T/\mathfrak{b}^q T})$ is injective $\forall 0 \leq i \leq r, q \geq 1$.

(B, \mathfrak{b}) a Noetherian local ring, R fin. gen. pos. graded B -algebra. M a finitely generated graded R -module. $T = B[x_1, \dots, x_r]$ pos. graded with $T \twoheadrightarrow R$.

Theorem (CR)

Assume M is a free B -module. The following six conditions are equivalent:

- ① $H_m^i(M)$ is a free B -module $\forall 0 \leq i \leq r$.
- ② $\text{Ext}_T^i(M, T)$ is a free B -module $\forall 0 \leq i \leq r$.
- ③ $H_m^i(M/\mathfrak{b}^q M)$ is a free B/\mathfrak{b}^q -module $\forall 0 \leq i \leq r, q \geq 1$.
- ④ $\text{Ext}_{T/\mathfrak{b}^q T}^i(M/\mathfrak{b}^q M, T/\mathfrak{b}^q T)$ is a free B/\mathfrak{b}^q -module $\forall 0 \leq i \leq r, q \geq 1$.
- ⑤ $H_m^i(M/\mathfrak{b}^q M) \rightarrow H_m^i(M/\mathfrak{b} M)$ is surjective $\forall 0 \leq i \leq r, q \geq 1$.
- ⑥ $\text{Ext}_{T/\mathfrak{b}^q T}^i(M/\mathfrak{b} M, \omega_{T/\mathfrak{b}^q T}) \rightarrow \text{Ext}_{T/\mathfrak{b}^q T}^i(M/\mathfrak{b}^q M, \omega_{T/\mathfrak{b}^q T})$ is injective $\forall 0 \leq i \leq r, q \geq 1$.

Moreover, when any of these conditions is satisfied, we have $H_m^i(M) \otimes_B C \xrightarrow{\cong} H_m^i(M \otimes_B C)$, $\text{Ext}_T^i(M, T) \otimes_B C \xrightarrow{\cong} \text{Ext}_{T \otimes_B C}^i(M \otimes_B C, T \otimes_B C)$ and $H_m^i(M) \cong {}^* \text{Hom}_B(\text{Ext}_T^{r-i}(M, T(-\delta)))$ where $\delta = \deg(x_1) + \dots + \deg(x_r)$.

Break?



A Noetherian, $S = \operatorname{Spec}(A)$, $X \subset \mathbb{P}_A^r$ closed subscheme, \mathcal{F} coherent sheaf on X .

Definition

\mathcal{F} is **fiber-full over S** if $H^i(X, \mathcal{F}(\nu))$ is a locally free A -module $\forall i, \nu$.

A Noetherian, $S = \operatorname{Spec}(A)$, $X \subset \mathbb{P}_A^r$ closed subscheme, \mathcal{F} coherent sheaf on X .

Definition

\mathcal{F} is **fiber-full over S** if $H^i(X, \mathcal{F}(\nu))$ is a locally free A -module $\forall i, \nu$.

Corollary

The following conditions are equivalent:

- 1 \mathcal{F} is fiber-full over S .
- 2 \mathcal{F} is a locally free \mathcal{O}_S -module and $H^i\left(X \times_S \operatorname{Spec}\left(\frac{A_{\mathfrak{p}}}{\mathfrak{p}^q A_{\mathfrak{p}}}\right), \mathcal{F}(\nu) \otimes_A \frac{A_{\mathfrak{p}}}{\mathfrak{p}^q A_{\mathfrak{p}}}\right)$ is a free $\frac{A_{\mathfrak{p}}}{\mathfrak{p}^q A_{\mathfrak{p}}}$ -module $\forall \mathfrak{p} \in S, i, q, \nu$.

- 3 \mathcal{F} is a locally free \mathcal{O}_S -module and the natural map

$$H^i\left(X \times_S \operatorname{Spec}\left(\frac{A_{\mathfrak{p}}}{\mathfrak{p}^q A_{\mathfrak{p}}}\right), \mathcal{F}(\nu) \otimes_A \frac{A_{\mathfrak{p}}}{\mathfrak{p}^q A_{\mathfrak{p}}}\right) \rightarrow H^i\left(X \times_S \operatorname{Spec}(\kappa(\mathfrak{p})), \mathcal{F}(\nu) \otimes_A \kappa(\mathfrak{p})\right)$$

is surjective for all $\forall \mathfrak{p} \in S, i, q, \nu$.

Fiber-full functor (again)

The fiber-full functor $\underline{\text{Fib}}_{\mathcal{F}/X/S}$

$$\underline{\text{Fib}}_{\mathcal{F}/X/S}(T) = \{\text{coherent } \mathcal{F}_T \rightarrow \mathcal{G} \mid \mathcal{G} \text{ is fiber-full over } S\}$$

Remark

When T is connected, we have the decomposition

$$\underline{\text{Fib}}_{\mathcal{F}/X/S}(T) = \bigsqcup_{\mathbf{h}: \mathbb{Z}^{r+1} \rightarrow \mathbb{N}^{r+1}} \underline{\text{Fib}}_{\mathcal{F}/X/S}^{\mathbf{h}}(T).$$

Therefore, if all the subfunctors $\underline{\text{Fib}}_{\mathcal{F}/X/S}^{\mathbf{h}}$ are representable, then $\underline{\text{Fib}}_{\mathcal{F}/X/S}$ is also representable (and a disjoint union of them).

Our main result

A Noetherian, $S = \operatorname{Spec}(A)$, $X \subset \mathbb{P}_A^r$ closed subscheme, \mathcal{F} coherent sheaf on X .

Theorem (CR - Ramkumar)

Let $\mathbf{h} = (h_0, \dots, h_r) : \mathbb{Z}^{r+1} \rightarrow \mathbb{N}^{r+1}$ be a tuple of functions. Assume $P_{\mathbf{h}} = \sum_{i=0}^r (-1)^i h_i \in \mathbb{Q}[m]$ is a numerical polynomial. Then, there is a quasi-projective S -scheme $\operatorname{Fib}_{\mathcal{F}/X/S}^{\mathbf{h}}$ that represents the functor $\underline{\operatorname{Fib}}_{\mathcal{F}/X/S}^{\mathbf{h}}$ and that is a locally closed subscheme of the Quot scheme $\operatorname{Quot}_{\mathcal{F}/X/S}^{P_{\mathbf{h}}}$.

We call $\operatorname{Fib}_{\mathcal{F}/X/S}^{\mathbf{h}}$ the **fiber-full scheme**. When $\mathcal{F} = \mathcal{O}_X$, we simply write $\operatorname{Fib}_{X/S}^{\mathbf{h}} \subset \operatorname{Hilb}_{X/S}^{\mathbf{h}}$, instead of $\operatorname{Fib}_{\mathcal{O}_X/X/S}^{\mathbf{h}}$.

General ideal of the proof

Reminder

The proof of the existence of the Quot scheme consists of two steps:

- 1 One embeds the Quot functor into a Grassmannian functor (not so deep, but it contains some tricky computations: Castelnuovo-Mumford regularity, etc...). The Grassmannian scheme represents the Grassmannian functor.
- 2 One applies a flattening stratification over the universal sheaf of the Grassmannian (this is the deeper part of the proof).

Theorem (Grothendieck, Mumford)

S locally Noetherian scheme, $X \subset \mathbb{P}_S^r$ closed subscheme, \mathcal{F} coherent sheaf on X . $P \in \mathbb{Q}[m]$ a numerical polynomial. There is a locally closed subscheme $\iota : V_{\mathcal{F}}^P \hookrightarrow S$ such that for any morphism $g : T = \operatorname{Spec}(B) \rightarrow S$, \mathcal{F}_T is T -flat with Hilbert polynomial P **if and only if** g can be factored as

$$T \rightarrow V_{\mathcal{F}}^P \xrightarrow{\iota} S.$$

S locally Noetherian scheme, $X \subset \mathbb{P}_S^r$ closed subscheme, \mathcal{F} coherent sheaf on X .
 $\mathbf{h} = (h_0, \dots, h_r) : \mathbb{Z}^{r+1} \rightarrow \mathbb{N}^{r+1}$ a tuple of functions.

Theorem (CR - Ramkumar)

Assume \mathcal{F} is flat over S . There is a locally closed subscheme $\iota : \text{FDir}_{\mathcal{F}}^{\mathbf{h}} \hookrightarrow S$ such that for any morphism $g : T = \text{Spec}(B) \rightarrow S$, $H^i(X_T, \mathcal{F}_T(\nu))$ is a locally free B -module of rank $h_i(\nu) \forall i, \nu$ **if and only if** g can be factored as

$$T \rightarrow \text{FDir}_{\mathcal{F}}^{\mathbf{h}} \xrightarrow{\iota} S.$$

S locally Noetherian scheme, $X \subset \mathbb{P}_S^r$ closed subscheme, \mathcal{F} coherent sheaf on X .
 $\mathbf{h} = (h_0, \dots, h_r) : \mathbb{Z}^{r+1} \rightarrow \mathbb{N}^{r+1}$ a tuple of functions.

Theorem (CR - Ramkumar)

Assume \mathcal{F} is flat over S . There is a locally closed subscheme $\iota : \text{FDir}_{\mathcal{F}}^{\mathbf{h}} \hookrightarrow S$ such that for any morphism $g : T = \text{Spec}(B) \rightarrow S$, $H^i(X_T, \mathcal{F}_T(\nu))$ is a locally free B -module of rank $h_i(\nu) \forall i, \nu$ **if and only if** g can be factored as

$$T \rightarrow \text{FDir}_{\mathcal{F}}^{\mathbf{h}} \xrightarrow{\iota} S.$$

Proof of the existence of $\text{Fib}_{\mathcal{F}/X/S}^{\mathbf{h}}$

- 1 We already have $\underline{\text{Fib}}_{\mathcal{F}/X/S}^{\mathbf{h}} \hookrightarrow \underline{\text{Quot}}_{\mathcal{F}/X/S}^{P_{\mathbf{h}}}$. If $\mathcal{G} \in \underline{\text{Quot}}_{\mathcal{F}/X/S}^{P_{\mathbf{h}}}(T)$, then $\mathcal{G} = (\mathcal{W}_{\mathcal{F}/X/S}^{P_{\mathbf{h}}})_T = (1_X \times_S g_{\mathcal{G}})^* \mathcal{W}_{\mathcal{F}/X/S}^{P_{\mathbf{h}}}$ where $g_{\mathcal{G}} : T \rightarrow \underline{\text{Quot}}_{\mathcal{F}/X/S}^{P_{\mathbf{h}}}$.
- 2 The theorem above yields that, $H^i(X_T, \mathcal{G}(\nu))$ is a locally free B -module of rank $h_i(\nu) \forall i, \nu$ **if and only if** $g_{\mathcal{G}}$ can be factored as

$$T \rightarrow \text{FDir}_{\mathcal{W}_{\mathcal{F}/X/S}^{P_{\mathbf{h}}}}^{\mathbf{h}} \xrightarrow{\iota} \underline{\text{Quot}}_{\mathcal{F}/X/S}^{P_{\mathbf{h}}}.$$

Therefore, we have $\text{Fib}_{\mathcal{F}/X/S}^{\mathbf{h}} := \text{FDir}_{\mathcal{W}_{\mathcal{F}/X/S}^{P_{\mathbf{h}}}}^{\mathbf{h}}$.

A Noetherian, $S = \operatorname{Spec}(A)$, R fin. gen. graded A -algebra. M fin. gen. graded R -module.

Theorem (CR - Ramkumar)

Assume M is flat over A . There is a locally closed subscheme $\iota : \operatorname{FLoc}_M^h \hookrightarrow S$ such that for any morphism $g : T = \operatorname{Spec}(B) \rightarrow S = \operatorname{Spec}(A)$, $[H_m^i(M \otimes_A B)]_\nu$ is a locally free B -module of rank $h_i(\nu) \forall i, \nu$ **if and only if** g can be factored as

$$T \rightarrow \operatorname{FLoc}_{\mathcal{F}}^h \xrightarrow{\iota} S.$$

Step 0 in the proof of our stratification theorem

Lemma (Grothendieck's complex)

A Noetherian, $S = \operatorname{Spec}(A)$, $X \subset \mathbb{P}_A^r$ closed subscheme, \mathcal{F} coherent sheaf on X that is flat over S . There is a complex K^\bullet of finitely generated free A -modules such that

$$H^i(X, \mathcal{F} \otimes_A N) \cong H^i(K^\bullet \otimes_A N)$$

for any A -module N .

Step 0 in the proof of our stratification theorem

Lemma (Grothendieck's complex)

A Noetherian, $S = \operatorname{Spec}(A)$, $X \subset \mathbb{P}_A^r$ closed subscheme, \mathcal{F} coherent sheaf on X that is flat over S . There is a complex K^\bullet of finitely generated free A -modules such that

$$H^i(X, \mathcal{F} \otimes_A N) \cong H^i(K^\bullet \otimes_A N)$$

for any A -module N .

Lemma (Jouanolou's complex)

A Noetherian, $R = A[x_1, \dots, x_r]$, M a finitely generated graded R -module that is flat over A . Let $F_\bullet : \cdots \rightarrow F_1 \rightarrow F_0$ be a graded free resolution of M by modules of finite rank. Consider the complex $L_\bullet = H_m^r(F_\bullet)$ (Note: each graded strand $[L_\bullet]_\nu$ is a complex of finitely generated free A -modules). Then

$$H_m^i(M \otimes_A N) \cong H_{r-i}(L_\bullet \otimes_A N)$$

for any A -module N .

Example (Twisted cubics)

$\text{Hilb}_{\mathbb{P}^3_{\mathbb{k}}}^{3m+1} = \text{Fib}_{\mathbb{P}^3_{\mathbb{k}}}^{\mathbf{h}} \sqcup \text{Fib}_{\mathbb{P}^3_{\mathbb{k}}}^{\mathbf{g}}$, where $\text{Fib}_{\mathbb{P}^3_{\mathbb{k}}}^{\mathbf{h}} = H - H \cap H'$ is open and $\text{Fib}_{\mathbb{P}^3_{\mathbb{k}}}^{\mathbf{g}} = H'$ is closed (we explicitly saw \mathbf{h} and \mathbf{g}).

Example (Twisted cubics)

$\text{Hilb}_{\mathbb{P}^3_{\mathbb{k}}}^{3m+1} = \text{Fib}_{\mathbb{P}^3_{\mathbb{k}}}^{\mathbf{h}} \sqcup \text{Fib}_{\mathbb{P}^3_{\mathbb{k}}}^{\mathbf{g}}$, where $\text{Fib}_{\mathbb{P}^3_{\mathbb{k}}}^{\mathbf{h}} = H - H \cap H'$ is open and $\text{Fib}_{\mathbb{P}^3_{\mathbb{k}}}^{\mathbf{g}} = H'$ is closed (we explicitly saw \mathbf{h} and \mathbf{g}).

Example (Points)

Let $\mathbf{h} = (c, 0, \dots, 0) : \mathbb{Z}^{r+1} \rightarrow \mathbb{N}^{r+1}$ and so $P_{\mathbf{h}} = c \in \mathbb{Q}[m]$. Then, we have

$$\text{Hilb}_{\mathbb{P}^r_{\mathbb{k}}}^c = \text{Fib}_{\mathbb{P}^r_{\mathbb{k}}}^{\mathbf{h}}.$$

Example (Twisted cubics)

$\text{Hilb}_{\mathbb{P}^3}^{3m+1} = \text{Fib}_{\mathbb{P}^3}^{\mathbf{h}} \sqcup \text{Fib}_{\mathbb{P}^3}^{\mathbf{g}}$, where $\text{Fib}_{\mathbb{P}^3}^{\mathbf{h}} = H - H \cap H'$ is open and $\text{Fib}_{\mathbb{P}^3}^{\mathbf{g}} = H'$ is closed (we explicitly saw \mathbf{h} and \mathbf{g}).

Example (Points)

Let $\mathbf{h} = (c, 0, \dots, 0) : \mathbb{Z}^{r+1} \rightarrow \mathbb{N}^{r+1}$ and so $P_{\mathbf{h}} = c \in \mathbb{Q}[m]$. Then, we have

$$\text{Hilb}_{\mathbb{P}^r}^c = \text{Fib}_{\mathbb{P}^r}^{\mathbf{h}}.$$

Example (Smooth Hilbert schemes)

$P \in \mathbb{Q}[m]$ such that $\text{Hilb}_{\mathbb{P}^r}^P$ is smooth, $L \subset \mathbb{k}[\mathbb{P}^r]$ be the corresponding saturated lexicographic ideal and $\mathbf{h} = (h_0, \dots, h_r)$ with $h_i(\nu) = \dim_{\mathbb{k}} (H^i(\mathbb{P}^r, \mathcal{O}_{V(L)}))$. By using the classification of [Skjeltnes - Smith \(2021\)](#), we can prove that

$$\text{Fib}_{\mathbb{P}^r}^{\mathbf{h}} = \text{Hilb}_{\mathbb{P}^r}^P.$$

Parametrizing ACM and AG schemes

$Y \subset \mathbb{P}_{\mathbb{k}}^r$ is said to be **arithmetically Cohen-Macaulay** or **arithmetically Gorenstein** when the homogeneous coordinate ring is Cohen-Macaulay or Gorenstein, resp.

Parametrizing ACM and AG schemes

$Y \subset \mathbb{P}_{\mathbb{k}}^r$ is said to be **arithmetically Cohen-Macaulay** or **arithmetically Gorenstein** when the homogeneous coordinate ring is Cohen-Macaulay or Gorenstein, resp.

$S = \text{Spec}(A)$ with A Noetherian. Let $d \in \mathbb{N}$, and $h_0, h_d : \mathbb{Z} \rightarrow \mathbb{N}$ be two functions. As all intermediate cohomologies vanish, we want to consider the functors

$$\underline{\mathcal{ACM}}_{X/S}^{h_0, h_d}(T) = \{Z \in \underline{\mathcal{Fib}}_{X/S}^{\mathbf{h}}(T) \mid Z_t \text{ is ACM for all } t \in T\}$$

and

$$\underline{\mathcal{AG}}_{X/S}^{h_0, h_d}(T) = \{Z \in \underline{\mathcal{Fib}}_{X/S}^{\mathbf{h}}(T) \mid Z_t \text{ is AG for all } t \in T\}$$

where $\mathbf{h} = (h_0, 0, \dots, 0, h_d, 0, \dots, 0) : \mathbb{Z}^{r+1} \rightarrow \mathbb{N}^{r+1}$.

Theorem (CR - Ramkumar)

$\underline{\mathcal{ACM}}_{X/S}^{h_0, h_d}$ and $\underline{\mathcal{AG}}_{X/S}^{h_0, h_d}$ are represented by open S -subschemes $\mathcal{ACM}_{X/S}^{h_0, h_d}$ and $\mathcal{AG}_{X/S}^{h_0, h_d}$ of $\text{Fib}_{X/S}^{\mathbf{h}}$.

Square-free Gröbner degenerations

Theorem (Conca - Varbaro)

$R = \mathbb{k}[x_1, \dots, x_r]$, $>$ monomial order on R and $I \subset R$ homogeneous ideal. If $\text{in}_{>}(I)$ is square-free, then

$$\dim_{\mathbb{k}} ([H_{\mathbf{m}}^i(R/I)]_{\nu}) = \dim_{\mathbb{k}} ([H_{\mathbf{m}}^i(R/\text{in}_{>}(I))]_{\nu})$$

for all i, ν .

Square-free Gröbner degenerations

Theorem (Conca - Varbaro)

$R = \mathbb{k}[x_1, \dots, x_r]$, $>$ monomial order on R and $I \subset R$ homogeneous ideal. If $\text{in}_{>}(I)$ is square-free, then

$$\dim_{\mathbb{k}} ([H_{\mathbf{m}}^i(R/I)]_{\nu}) = \dim_{\mathbb{k}} ([H_{\mathbf{m}}^i(R/\text{in}_{>}(I))]_{\nu})$$

for all i, ν .

Corollary

Let $\text{hom}_{\omega}(I) \subset R[t]$ with special fiber equal to $\text{in}_{>}(I)$. Let $Z = \text{Proj}(S/I) \subset \mathbb{P}_{\mathbb{k}}^r$. Let $\mathbf{h} = (h_0, \dots, h_r) : \mathbb{Z}^{r+1} \rightarrow \mathbb{N}^{r+1}$ given by $h_i(\nu) := \dim_{\mathbb{k}} (H^i(Z, \mathcal{O}_Z(\nu)))$. For each $\alpha \in \mathbb{k}$, let $Z_{\alpha} = \text{Proj}(R[t]/\text{hom}_{\omega}(I) \otimes_{\mathbb{k}[t]} \mathbb{k}[t]/(t - \alpha)) \subset \mathbb{P}_{\mathbb{k}}^r$. Then, we have that

Z_{α} corresponds with a point in $\text{Fib}_{\mathbb{P}_{\mathbb{k}}^r/\mathbb{k}}^{\mathbf{h}}$

for all $\alpha \in \mathbb{k}$.

Future directions

- 1 **A compactification of the fiber-full scheme.** Find the most natural compactification of the fiber-full scheme.
- 2 **Deformation theory on the fiber-full scheme.** For instance, computing the tangent space $T_{[Z]} \text{Fib}_{\mathbb{P}_{\mathbb{k}}^r}^h$ at $Z \in \mathbb{P}_{\mathbb{k}}^r$ is equivalent to find all $Z' \subset \mathbb{P}_{\mathbb{k}[\epsilon]}^r$ such that $Z \cong Z' \times_{\text{Spec}(\mathbb{k}[\epsilon])} \text{Spec}(\mathbb{k})$ and $H^i(Z', \mathcal{O}_{Z'}(\nu))$ is a $\mathbb{k}[\epsilon]$ -flat $\forall i, \nu$, where $\mathbb{k}[\epsilon] = \mathbb{k}[t]/(t^2)$.
- 3 **Understand small neighborhoods of monomial ideals in the fiber-full scheme.**



Thanks!