# Pre-Calculus 11 Factoring Trinomials

Created by Yi-Chen Lin

June 16, 2025

# Factoring Trinomials

#### Introduction

- In this section, you will be factoring trinomials where the coefficient of  $x^2$  is not equal to one.
- Examples:
  - $5x^2 + 17x + 6$  (Can't factor out common factors like the previous section.)
  - $7x^2 18x + 8$
- There are 3 different methods for factoring Trinomials:
  - B.U.M. Method
    - Easiest, straight-forward, Long
  - Criss-Cross Method
    - Fast, Quick with Numbers, Hard
  - Grouping Method
    - Textbook, standard method



2/30

## BUM Method - Example

## Example: Factor using BUM Method

Factor the following Trinomial using the BUM Method:

$$7x^2 - 18x + 8$$

# BUM Method - Example Solution

#### **Detailed Solution**

Factor  $7x^2 - 18x + 8$  using the BUM Method:

Bring the First term to the Last term and Multiply them

$$= x^2 - 18x + 56$$

Factor, two numbers that multiply to 56 and adds to -18

$$=(x-14)(x-4)$$

Bring the First term back in front of each  $\times$ 

$$= (7x - 14)(7x - 4)$$

Factor/Bum out any common factors in each binomial:

$$= \left(\frac{7x - 14}{7}\right) \left(\frac{7x - 4}{1}\right)$$
$$= (x - 2)(7x - 4)$$

## BUM Method - Practice

## Practice: Factor using BUM Method

Factor each of the following Trinomials using the BUM Method:

$$15x^2 + 22x + 8$$

$$214x^2 - 27x + 9$$

## BUM Method - Solutions Part 1

#### **Detailed Solutions**

**1**  $5x^2 + 22x + 8$  **Solution:** 

$$15x^{2} + 22x + 8$$

$$= x^{2} + 22x + 120$$

$$= (x + 12)(x + 10)$$

$$= (15x + 12)(15x + 10)$$

$$= \left(\frac{15x + 12}{3}\right) \left(\frac{15x + 10}{5}\right)$$

$$= (5x + 4)(3x + 2)$$

6/30

## BUM Method - Solutions Part 2

#### **Detailed Solutions**

**2**  $14x^2 - 27x + 9$  **Solution:** 

$$14x^{2} - 27x + 9$$

$$= x^{2} - 27x + 126$$

$$= (x - 21)(x - 6)$$

$$= (14x - 21)(14x - 6)$$

$$= \left(\frac{14x - 21}{7}\right) \left(\frac{14x - 6}{2}\right)$$

$$= (2x - 3)(7x - 3)$$

# Grouping Method - Example

## Example: Factor using Grouping Method

Factor the following Trinomial using the Grouping Method:

$$9x^2 + 15x + 4$$

# Grouping Method - Example Solution (Part 1)

#### **Detailed Solution**

Factor  $9x^2 + 15x + 4$  using the Grouping Method:

Multiply the First & Last Numbers

Find 2 numbers that MULTIPLY to 36 and ADDS to 15

$$3\times12\rightarrow Adds$$
 to  $15$ 

$$=9x^2+3x+12x+4$$

Split the 15x into the two factors

9/30

# Grouping Method - Example Solution (Part 2)

## Detailed Solution (Cont.)

Factor  $9x^2 + 15x + 4$  using the Grouping Method (Cont.):

$$= (9x^2 + 3x) + (12x + 4)$$

Group the First 2 and Last 2 terms

$$=3x(3x+1)+4(3x+1)$$

Factor out any common factors from each bracket

$$=(3x+1)(3x+4)$$

The Binomial is a GCF. Factor it out

10 / 30

# Grouping Method - Practice

## Practice: Factor using Grouping Method

Factor each of the following using the Grouping Method:

$$12x^2 + 13x - 14$$

$$9x^2 + 21x - 8$$

# Grouping Method - Solutions Part 1

#### **Detailed Solutions**



$$12x^2 + 13x - 14$$

Multiply 
$$12 \times -14 = -168$$

Find two numbers that multiply to -168 and add to 13: 21, -8

$$= 12x^2 + 21x - 8x - 14$$

$$= (12x^2 + 21x) + (-8x - 14)$$

$$=3x(4x+7)-2(4x+7)$$

$$=(4x+7)(3x-2)$$

# Grouping Method - Solutions Part 2

#### **Detailed Solutions**



**2** 
$$9x^2 + 21x - 8$$
 **Solution:**

$$9x^2 + 21x - 8$$

Multiply 
$$9 \times -8 = -72$$

Find two numbers that multiply to -72 and add to 21: 24, -3

$$= 9x^2 + 24x - 3x - 8$$

$$= (9x^2 + 24x) + (-3x - 8)$$

$$=3x(3x+8)-1(3x+8)$$

$$=(3x+8)(3x-1)$$

13 / 30

## Criss-Cross Method - Example

## Example: Factor using Criss-Cross Method

Factor the following using the Criss-Cross Method:

$$24x^2 + 2x - 15$$

# Criss-Cross Method - Example Solution (Part 1)

#### **Detailed Solution**

Factor  $24x^2 + 2x - 15$  using the Criss-Cross Method:

Pick 2 numbers that multiply to the FIRST term  $(24x^2)$ : Multiply sides ways or Criss-Cross:

4x and 6x  $4x \rightarrow -3 = -12x$ 

Pick 2 numbers that multiply to the LAST term (-15):  $6x \rightarrow 5 = 30x$ 

-3 and 5 18x (Sum must equal the middle term)

15/30

# Criss-Cross Method - Example Solution (Part 2)

## Detailed Solution (Cont.)

The sum 18x does not equal the middle term 2x. Let's try different factors:

Pick 2 numbers that multiply to the FIRST term  $(24x^2)$ : Multiply sides ways or Criss-Cross:

4x and 6x  $4x \rightarrow 5 = 20x$ 

Pick 2 numbers that multiply to the LAST term (-15):  $6x \rightarrow -3 = -18x$ 

5 and -3 2x (Sum must equal the middle term)

(4x-3)(6x+5)

Numbers on the left go in front of each bracket

16 / 30

## Criss-Cross Method - Practice

## Practice: Factor using Criss-Cross Method

Factor the following using the Criss-Cross Method:

$$0 8x^2 - 26x + 15$$

$$6x^2 - 17x + 5$$

# Criss-Cross Method - Solutions Part 1 (Cont.)

#### **Detailed Solutions**

**1** 
$$8x^2 - 26x + 15$$
 **Solution:**

$$\begin{array}{cccc}
2x & \rightarrow & -5 & = -10x \\
4x & \rightarrow & -3 & = -12x
\end{array}$$

 $\rightarrow$  -3 = -12x -22x (Incorrect sum) Therefore, the factors are (2x - 5)(4x - 3).

(

# Criss-Cross Method - Solutions Part 2 (Cont.)

#### **Detailed Solutions**

**1** 
$$8x^2 - 26x + 15$$
 **Solution:**

$$\begin{array}{cccc}
2x & \rightarrow & -3 & = -6x \\
4x & \rightarrow & -5 & = -20x
\end{array}$$

Therefore, the factors are (2x-5)(4x-3).

-26x (Correct sum)

(

## Mixed Practice: Factor Trinomials

## Factor Each of the Following Trinomials

$$20x^3 - 80x^2 + 35x$$

$$6x^4 - 17x^2y - 10y^2$$

## Mixed Practice: Solutions Part 1

#### **Detailed Solutions**

**1** 
$$20x^3 - 80x^2 + 35x$$
 **Solution:**

$$20x^3 - 80x^2 + 35x$$
$$= 5x(4x^2 - 16x + 7)$$

$$= 5x(4x - 10x + 1)$$

Factor 
$$4x^2 - 16x + 7$$
 (BUM Method:  $4 \times 7 = 28$ ; add to -16: -14, -2)  $= 5x(4x - 14)(4x - 2)$ 

$$=5x\left(\frac{4x-14}{2}\right)\left(\frac{4x-2}{2}\right)$$

$$=5x(2x-7)(2x-1)$$



 Created by Yi-Chen Lin
 Pre-Calculus 11
 June 16, 2025
 21 / 30

## Mixed Practice: Solutions Part 2

#### **Detailed Solutions**

② 
$$6x^4 - 17x^2y - 10y^2$$
 Solu

$$6x^4 - 17x^2y - 10y^2$$

Treat  $x^2$  as A and y as  $B: 6A^2 - 17AB - 10B^2$ 

BUM Method:  $6 \times -10 = -60$ ; add to -17: -20, 3

$$=(6A-20B)(6A+3B)$$

$$= \left(\frac{6A - 20B}{2}\right) \left(\frac{6A + 3B}{3}\right)$$

$$= (3A - 10B)(2A + B)$$

Substitute back  $A = x^2$  and B = y

$$= (3x^2 - 10y)(2x^2 + y)$$

# Mixed Practice: Factor Each of the Following

## Factor Each of the Following

$$14x^2 - 23x + 3$$

$$2 18x^2 + 27x + 10$$

$$30x^2 + x - 1$$

$$12x^2 + 29xy + 14y^2$$

$$5x^2 + 11xy + 6y^2$$

**6** 
$$4x^4 - 25x^2y^2 + 36y^4$$

## Mixed Practice: Solutions Part 3

#### **Detailed Solutions**

**1** 
$$14x^2 - 23x + 3$$
 **Solution:**  $(7x - 1)(2x - 3)$ 

② 
$$18x^2 + 27x + 10$$
 Solution:  $(3x + 2)(6x + 5)$ 

**3** 
$$30x^2 + x - 1$$
 **Solution**:  $(5x + 1)(6x - 1)$ 

 Created by Yi-Chen Lin
 Pre-Calculus 11
 June 16, 2025
 24 / 30

#### **Detailed Solutions**

**1** 
$$12x^2 + 29xy + 14y^2$$
 **Solution:**  $(3x + 2y)(4x + 7y)$ 

**5** 
$$5x^2 + 11xy + 6y^2$$
 **Solution:**  $(x + y)(5x + 6y)$ 

**6** 
$$4x^4 - 25x^2y^2 + 36y^4$$
 **Solution:**  $(x - 2y)(x + 2y)(2x - 3y)(2x + 3y)$ 

 Created by Yi-Chen Lin
 Pre-Calculus 11
 June 16, 2025
 25 / 30

# Application Problem: Area and Perimeter

#### Area and Perimeter Application

Ex: Given that the area of a rectangle is  $12x^2 + 23x + 10$ , then which of the following expressions is the perimeter?

- 14x + 12
- ② 14x + 14
- 37x + 7
- 0 28x + 28

26 / 30

## Application Problem: Area and Perimeter - Solution

#### **Detailed Solution**

Ex: Given that the area of a rectangle is  $12x^2 + 23x + 10$ , then which of the following expressions is the perimeter?

#### Solution:

Factor the area: 
$$12x^2 + 23x + 10$$

Multiply 
$$12 \times 10 = 120$$

Find two numbers that multiply to 120 and add to 23: 8,15

$$= (12x + 8)(12x + 15)$$

$$= \left(\frac{12x + 8}{4}\right) \left(\frac{12x + 15}{3}\right)$$

$$= (3x + 2)(4x + 5)$$

So, Length = 4x + 5 and Width = 3x + 2.

Perimeter = 2(Length + Width)

# Application Problem: Integral Values of k

## Integral Values of k

Ex: For which integral values of k can  $6x^2 + kx + 1$  be factored?

- 5,70nly
- $2 \pm 5, \pm 7$  only
- **3** -5, -7 *only*
- all integers between -7 and 5, inclusive

# Application Problem: Integral Values of k - Solution

#### **Detailed Solution**

Ex: For which integral values of k can  $6x^2 + kx + 1$  be factored?

**Solution:** We need to find combinations of factors for  $6x^2$  and 1 that produce integral values for k.

- Factors of  $6x^2$ : (x, 6x), (2x, 3x)
- Factors of 1: (1,1), (-1,-1)

Possible cross products for k (middle term):

• 
$$(x+1)(6x+1) \rightarrow 1x + 6x = 7x \rightarrow k = 7$$

• 
$$(x-1)(6x-1) \rightarrow -1x - 6x = -7x \rightarrow k = -7$$

• 
$$(2x+1)(3x+1) \rightarrow 2x+3x=5x \rightarrow k=5$$

• 
$$(2x-1)(3x-1) \rightarrow -2x-3x = -5x \rightarrow k = -5$$

The possible integral values for k are  $\pm 5, \pm 7$ . The correct option is **b)**  $\pm 5, \pm 7$  only.



 Created by Yi-Chen Lin
 Pre-Calculus 11
 June 16, 2025
 29 / 30

# Summary

## **Key Concepts**

- Understanding Trinomial Factoring when  $a \neq 1$
- B.U.M. Method for factoring trinomials
- Grouping Method for factoring trinomials
- Criss-Cross Method for factoring trinomials
- Mixed practice involving various factoring scenarios

30 / 30