

# **MATH 101 Study Notes**

## **Integral Calculus with**

## **Applications**

Yecheng Liang

## Contents

|                                |   |
|--------------------------------|---|
| 1. General Principles .....    | 3 |
| 1.1. Sigma Notation .....      | 3 |
| 2. Definite Integral .....     | 4 |
| 2.1. Estimation .....          | 4 |
| 2.2. Signed Area .....         | 5 |
| 2.3. Precise Calculation ..... | 5 |

# 1. General Principles

In this course,

- log is ln

## 1.1. Sigma Notation

$$\sum_{i=1}^n i \tag{1}$$

means “the sum of  $i$  from 1 to  $n$ , where  $i$  is an integer”.


## 2. Definite Integral

$$\int_a^b f(x) \, dx \quad (2)$$

means “the integral of  $f(x)$  from  $a$  to  $b$ ”.

$$\int_{-1}^1 |x| \, dx \quad (3)$$

and looks like:



### 2.1. Estimation

Right Riemann Sum (RRS) is an estimation of the area under the curve using rectangles with the right endpoint as the height.

For example,


$$\int_0^8 \sqrt{x} \, dx. \quad (4)$$

Using RRS with 4 rectangles, we have:

1. each rectangle has width  $\frac{8-0}{4} = 2$ ,
2. the right endpoints are 2, 4, 6, 8,
3. the heights are  $\sqrt{2}, \sqrt{4}, \sqrt{6}, \sqrt{8}$ .

That gives us the estimation:

$$\begin{aligned} \sum_{i=1}^4 2\sqrt{2i} &= 2\sqrt{2} + 2\sqrt{4} + 2\sqrt{6} + 2\sqrt{8} \\ &\approx 2.83 + 4 + 5.29 + 5.66 \\ &= 17.78. \end{aligned} \quad (5)$$



Similarly, Left Riemann Sum (LRS), Midpoint Riemann Sum (MRS), and Trapezoidal Riemann Sum (TRS) exist.

The generalized formula with  $n$  rectangles/trapeziums from  $a$  to  $b$  are:

$$\begin{aligned}
 \text{RRS}(a, b, n) &= \sum_{i=1}^n f(x_i) \Delta x \\
 \text{LRS}(a, b, n) &= \sum_{i=1}^n f(x_{i-1}) \Delta x \\
 \text{MRS}(a, b, n) &= \sum_{i=1}^n f\left(\frac{x_{i-1} + x_i}{2}\right) \Delta x \\
 \text{TRS}(a, b, n) &= \sum_{i=1}^n (f(x_{i-1}) + f(x_i)) \Delta \frac{x}{2},
 \end{aligned} \tag{6}$$

where  $\Delta x = \frac{b-a}{n}$  and  $x_i = a + i\Delta x$ .

For an increasing function ( $f'(x) > 0$ ), RRS is an overestimation, and LRS is an underestimation. For a function concave up ( $f''(x) > 0$ ), TRS is an overestimation.

## 2.2. Signed Area

If the 'area under the curve' is below the  $x$ -axis, it can be called 'negative'. Hence, the integral of a function can be interpreted as the signed area of a curve, which can be positive or negative.

## 2.3. Precise Calculation

Using Riemann Sums, we can see that the more rectangles we use, the closer the estimation is to the actual value.