
Complex Analysis I:

Problem Set IV

Youngduck Choi
CILVR Lab
New York University
yc1104@nyu.edu

Abstract

This work contains the solutions to the problem set IV of Complex Analysis I 2015 at Courant Institute of Mathematical Sciences.

Question 1. Brown p.147-2.

Solution. (b) We first have that $\cos(z/2)$ is continuous everywhere on the complex plane. Therefore, any contour from 0 to $\pi + 2i$ will have the same value of $F(\pi + 2i) - F(0)$, where F denotes the antiderivative of $\cos(z/2)$. We can compute the exact value as follows:

$$\begin{aligned}\int_0^{\pi+2i} \cos\left(\frac{z}{2}\right) dz &= \left[2\sin\left(\frac{z}{2}\right)\right]_0^{\pi+2i} \\ &= 2\sin\left(\frac{\pi}{2} + i\right) \\ &= 2\cos(i) \\ &= e + \frac{1}{e},\end{aligned}$$

as desired. \square

Question 5. Brown p.147-5.

Solution. Let C be a contour from -1 to 1 that lies above the x-axis. We wish to compute the following integral:

$$\int_C z^i dz,$$

where z^i denotes the principal branch $\exp(i\operatorname{Log}(z))$ for $|z| > 0, -\pi < \operatorname{Arg}(z) < \pi$. Notice that under the principal branch, $z = -1$ is not defined. The following branch, however, agrees with the integrand along C and has anti-derivative along C :

$$z^i = \exp(i\log(z)) \text{ for } (|z| > 0, -\frac{\pi}{2} < \arg(z) < \frac{3\pi}{2}).$$

We then can compute the integral as follows:

$$\begin{aligned}
\int_C z^i dz &= \left[\frac{z^{i+1}}{i+1} \right]_{-1}^1 \\
&= \frac{1}{i+1} (e^{(i+1)\log 1} - e^{(i+1)\log(-1)}) \\
&= \frac{1}{i+1} (e^{(i+1)(\ln 1)} - e^{(i+1)\ln(1+i\pi)}) \\
&= \frac{1}{i+1} (1 + e^{i\pi}) \\
&= \frac{1 + e^{-\pi}}{2} (1 - i),
\end{aligned}$$

as desired. \square

Question 1. Brown p.159-2.

Solution. (b) Let C_1 denote the positively oriented boundary of the square whose sides lie along the line $x = \pm 1, y = \pm 1$ and let C_2 be the positively oriented circle $|z| = 4$. Observe that C_1 is interior to C_2 and the given function $\frac{z+2}{\sin(\frac{z}{2})}$ is analytic in the closed region consisting of the C_1 and C_2 contours and all points between them. Hence, by the corollary, we have

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz,$$

for $f(z) = \frac{z+2}{\sin(\frac{z}{2})}$. \square

Question 2. Brown 159-4.

Solution. (a) Observe that along the lower horizontal leg, we have $z = x$ ($-a \leq x \leq a$). Hence, the integral along the lower horizontal leg from $-a$ to a can be written as

$$2 \int_0^a e^{-x^2} dx.$$

For the upper horizontal leg, we have $z = x + ib$ ($-a \leq x \leq a$). Hence, the integral along the upper horizontal from a to $-a$ can be written as

$$\int_a^{-a} e^{-(x+ib)^2} dx,$$

which can be simplified as follows:

$$\begin{aligned}
\int_a^{-a} e^{-(x+ib)^2} dx &= -e^{b^2} \int_{-a}^a e^{-x^2 - 2ibx} dx \\
&= -2e^{b^2} \int_0^a e^{-x^2} \cos(2bx) dx.
\end{aligned}$$

Hence, we have

$$2 \int_0^a e^{-x^2} dx + -2e^{b^2} \int_0^a e^{-x^2} \cos(2bx) dx,$$

for the sum of contour integrals along each horizontal leg. Now, observe that along the right vertical leg, we have $z = a + iy$ ($0 \leq y \leq b$). Hence,

Question 1. Brown p.170-1.

Solution. Let C denote the positively oriented boundary of the square whose sides lie along the lines $x = \pm 2$ and $y = \pm 2$. We evaluate the following integrals.

(b) We are given the following integral:

$$\int_C \frac{\cos(z)}{z(z^2 + 8)} dz,$$

which can be written as

$$\int_C \frac{\frac{\cos(z)}{(z^2+8)}}{z} dz.$$

As $\frac{\cos(z)}{(z^2 + 8)}$ is analytic everywhere inside and on the given contour, which is simple and closed, taken in the positive sense, by the Cauchy Integral formula, we obtain

$$\frac{\cos(0)}{8} = \frac{1}{2\pi i} \int_C \frac{\frac{\cos(z)}{(z^2+8)}}{z} dz,$$

which simplifies to

$$\int_C \frac{\frac{\cos(z)}{(z^2+8)}}{z} dz = \frac{\pi i}{4}.$$

(d) We are given the following integral:

$$\int_C \frac{\cosh(z)}{z^4} dz.$$

As $\frac{\cosh(z)}{z^4}$ is analytic everywhere inside and on the given contour, which is simple and closed, taken in the positive sense, by the extended Cauchy Integral formula, we obtain

$$\cosh^{(3)}(z_0) = \frac{3!}{2\pi i} \int_C \frac{\cosh(z)}{(z - z_0)^4} dz,$$

for z_0 inside and on the given contour. Observe that $\cosh^{(3)} = \sinh$. Hence, taking $z_0 = 0$ yields

$$0 = \frac{3!}{2\pi i} \int_C \frac{\cosh(z)}{z^4} dz,$$

which simplifies to

$$\int_C \frac{\cosh(z)}{z^4} dz = 0.$$

(e) We are given the following integral:

$$\int_C \frac{\tan(\frac{z}{2})}{(z - x_0)^2} dz,$$

for $-2 < x_0 < 2$. Notice that x_0 is inside the given contour. As $\frac{\tan(\frac{z}{2})}{(z - x_0)^2}$ is analytic everywhere inside and on the given contour, which is simple and closed, taken in the positive sense, by the extended Cauchy Integral formula, we obtain

$$\frac{1}{2} \sec^2\left(\frac{x_0}{2}\right) = \frac{1!}{2\pi i} \int_C \frac{\tan(\frac{z}{2})}{(z - x_0)^2} dz,$$

which simplifies to

$$\int_C \frac{\tan(\frac{z}{2})}{(z - x_0)^2} dz = i\pi \sec^2\left(\frac{x_0}{2}\right),$$

for $-2 < x_0 < 2$. \square

Question 2. Brown 170.3.

Solution. Let C be the circle $|z| = 3$, described in the positive sense. As $2s^2 - s - 2$ is analytic everywhere inside and on the given contour, which is simple and closed, taken in the positive sense, by the extended Cauchy Integral formula, we obtain

$$2z^2 - z - 2 = \frac{1}{2\pi i} \int_C \frac{2s^2 - s - 2}{s - z} ds,$$

for $|z| < 3$. As $g(z) = \int_C \frac{2s^2 - s - 2}{s - z} ds$, we have

$$g(z) = 2\pi i(2z^2 - z - 2),$$

for $|z| < 3$. Hence, it follows that $g(2) = 8\pi i$. For $|z| > 3$, we have that $\frac{2s^2 - s - 2}{s - z}$ is analytic at all points interior to and on C . Hence, by the Cauchy-Goursat theorem, we obtain

$$\int_C \frac{2s^2 - s - 2}{s - z} dz = 0,$$

for $|z| > 3$. Therefore, $g(z) = 0$ when $|z| > 3$, which completes the solution for the problem. \square

Question 3. Brown 170-4.

Solution. Let C be any simple closed contour, described in the positive sense in the z plane. As $s^3 + 2s$ is entire, by the extended Cauchy Integral formula, we obtain

$$6z = \frac{2!}{2\pi i} \int_C \frac{s^3 + 2s}{(s - z)^3} ds,$$

for z at the interior of C . As $g(z) = \int_C \frac{s^3 + 2s}{(s - z)^3} ds$, we have

$$g(z) = 6\pi iz,$$

for z inside C . Now, if z is outside of C , then $\frac{s^3 + 2s}{s - z}$ is analytic at points interior to and on C . Hence, by the Cauchy-Goursat Theorem, we have that

$$\int_C \frac{s^3 + 2s}{(s - z)^3} ds = 0,$$

for z outside of C . Hence $g(z) = 0$ when z is outside. \square

Question 4. Brown 170-7.

Solution. Let C be the unit circle. As e^{az} is entire, by the Cauchy Integral formula, we obtain

$$e^{az_0} = \frac{1}{2\pi i} \int_C \frac{e^{az}}{z - z_0} dz,$$

for z_0 inside C . By taking $z_0 = 0$, we get

$$1 = \frac{1}{2\pi i} \int_C \frac{e^{az}}{z} dz,$$

which simplifies to

$$\int_C \frac{e^{az}}{z} dz = 2\pi i.$$

Question 5. Brown 170-8.

Solution. The Legendre polynomials are defined by

$$P_n(z) = \frac{1}{2^{n+1}\pi i} \int_C \frac{(s^2 - 1)^n}{(s - z)^{n+1}} ds,$$

for any simple closed contour surrounding z . For $z = -1$, and by having C to be any arbitrary simple closed contour that surrounds $z = -1$, it follows that

$$P_n(-1) = \frac{1}{2^{n+1}\pi i} \int_C \frac{(s^2 - 1)^n}{(s + 1)^{n+1}} ds,$$

which, by using the suggestion, simplifies to

$$P_n(z) = \frac{1}{2^{n+1}\pi i} \int_C \frac{(s - 1)^n}{s + 1} ds.$$

Since $(s - 1)^n$ is entire, $(s - 1)^n$ is analytic inside and on C . Hence, by the Cauchy Integral formula, we have

$$(-2)^n 2\pi i = \int_C \frac{(s - 1)^n}{s + 1} ds.$$

Substituting the above equality into the simplified formula of Legendre polynomials yields

$$\begin{aligned} P_n(z) &= \frac{(-2)^n 2\pi i}{2^{n+1}\pi i} \\ &= (-1)^n, \end{aligned}$$

as desired. \square