

---

# Complex Analysis I:

## Problem Set VII

---

Youngduck Choi  
CILVR Lab  
New York University  
yc1104@nyu.edu

### Abstract

This work contains the solutions to the problem set VII of Complex Analysis I 2015 at Courant Institute of Mathematical Sciences.

#### Question 1. 247-7.

**Solution.** (a) Observe that

$$\frac{1}{z^2}f\left(\frac{1}{z}\right) = \frac{(3+2z)^2}{z(1-z)(2+5z)}.$$

As  $\frac{1}{z^2}f\left(\frac{1}{z}\right)$  has a simple pole at  $z = 0$ , we have

$$\begin{aligned}\int_C \frac{(3+2z)^2}{z(z-1)(2z+5)} dz &= 2\pi i \cdot \text{Res}_{z=0} \left[ \frac{1}{z^2}f\left(\frac{1}{z}\right) \right] \\ &= 2\pi i \cdot \frac{9}{2} = 9\pi i.\end{aligned}$$

(b) Observe that

$$\frac{1}{z^2}f\left(\frac{1}{z}\right) = \frac{e^z}{z^2(1+z^3)}.$$

As  $\frac{1}{z^2}f\left(\frac{1}{z}\right)$  has a pole of order 2 at  $z = 0$ , we have

$$\int_C \frac{z^3 e^{\frac{1}{z}}}{1+z^3} dz = 2\pi i \cdot \text{Res}_{z=0} \left[ \frac{1}{z^2}f\left(\frac{1}{z}\right) \right],$$

where  $\phi(z) = \frac{e^z}{1+z^3}$ . We have

$$\phi'(z) = \frac{(1+z^3)e^z - e^z 3z^2}{(1+z^3)^2}.$$

By substituting  $z = 0$ , we see that  $\phi'(0) = 1$ , which is the residue at  $z = 0$ . It follows that

$$\int_C \frac{z^3 e^{\frac{1}{z}}}{1+z^3} dz = 2\pi i.$$

□

**Question 2. 254-5.**

**Solution.** (a) The given integral can be written as

$$\begin{aligned}\int_C \tan(z) dz &= \int_C \frac{p(z)}{q(z)} dz \\ &= \int_C \frac{\sin(z)}{\cos(z)} dz.\end{aligned}$$

As the zeros of  $\cos(z)$  are  $z = \frac{\pi}{2} + n\pi$  and  $C$  is the positively oriented circle  $|z| = 2$ , there are two isolated singularities of  $\tan(z)$  interior to  $C$ , namely  $z = \pm \frac{\pi}{2}$ . It follows that

$$\begin{aligned}\text{Res}_{z=\frac{\pi}{2}} \tan(z) &= \frac{p(\frac{\pi}{2})}{q'(\frac{\pi}{2})} = \frac{\sin(\frac{\pi}{2})}{-\sin(\frac{\pi}{2})} = -1, \\ \text{Res}_{z=-\frac{\pi}{2}} \tan(z) &= \frac{p(-\frac{\pi}{2})}{q'(-\frac{\pi}{2})} = \frac{\sin(-\frac{\pi}{2})}{-\sin(-\frac{\pi}{2})} = -1.\end{aligned}$$

Consequently, by the residue theorem, we have

$$\int_C \tan(z) dz = -4\pi i.$$

(b) We wish to evaluate the integral  $\int_C \frac{dz}{\sinh(2z)}$ . As  $\sinh(z) = 0$  for  $\frac{n\pi i}{2}$ , we can conclude that the isolated singularities of the integrand happens at  $z = 0$  and  $z = \pm \frac{\pi i}{2}$ . It follows that

$$\begin{aligned}\text{Res}_{z=0} \frac{1}{\sinh(2z)} &= \frac{1}{2 \cosh(0)} = \frac{1}{2} \\ \text{Res}_{z=\frac{\pi i}{2}} \frac{1}{\sinh(2z)} &= \frac{1}{2 \cosh(\pi i)} = -\frac{1}{2} \\ \text{Res}_{z=-\frac{\pi i}{2}} \frac{1}{\sinh(2z)} &= \frac{1}{2 \cosh(-\pi i)} = -\frac{1}{2}\end{aligned}$$

Consequently, by the residue theorem, we have

$$\int_C \frac{1}{\sinh(2z)} dz = -\pi i.$$

□

**Question 3. 254-6.**

**Solution.** Observe that interior to  $C_N$ , the function  $\frac{1}{z^2 \sin(z)}$  has singularities at  $z = 0$  and  $z = \pm n\pi$  for  $n = 1, 2, \dots, N$ . The residue at  $z = 0$  can be found by finding the  $\frac{1}{z}$  coefficient of  $\frac{1}{z^2 \sin(z)}$ , which was found to be  $\frac{1}{6}$  in a previous homework problem. For the residues at  $z = \pm n\pi$  for  $n = 1, 2, \dots, N$ , we have

$$\frac{1}{z^2 \sin(z)} = \frac{p(z)}{q(z)} \text{ where } p(z) = 1, q(z) = z^2 \sin(z).$$

As  $q'(z) = z^2 \cos(z)$ , we have for  $z = \pm n\pi$ , we have  $q'(\pm n\pi) = (-1)^n n^2 \pi^2$ . By the residue theorem, it follows that

$$\int_{C_N} \frac{1}{z^2 \sin(z)} dz = 2\pi i \left[ \frac{1}{6} + 2 \sum_{n=1}^N \frac{(-1)^n}{n^2 \pi^2} \right],$$

which can be re-written as

$$\sum_{n=1}^N \frac{(-1)^{n+1}}{n^2} = \frac{\pi^2}{12} - \frac{\pi}{4i} \int_{C_N} \frac{1}{z^2 \sin(z)} dz.$$

We have that as  $N \rightarrow \infty$ , the integral goes to 0. Therefore, it follows that

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^2} = \frac{\pi^2}{12}.$$

□

**Question 254-8.**

**Solution.**

**Question 264.2.**

**Solution.** Consider the function  $f(z) = \frac{1}{(z^2 + 1)^2}$  and the simple, closed, and positively oriented contour of a half-circle above the real-axis, centered around the origin with the radius  $R > 1$ . It follows that

$$\int_{-R}^R \frac{1}{(x^2 + 1)^2} dx + \int_{C_R} \frac{1}{(z^2 + 1)^2} dz = 2\pi B,$$

where  $C_R$  denotes the contour of the curve part of the half-circle, and  $B$  is the residue of the complex integrand at  $z = i$ . For computing the residue, we have

$$\frac{1}{(z^2 + 1)} = \frac{\phi(z)}{(z - i)^2},$$

where  $\phi(z) = \frac{1}{(z + i)^2}$ . It follows that  $B = \phi'(i) = \frac{1}{4i}$ . Therefore, we obtain

$$\int_{-R}^R \frac{dx}{(x^2 + 1)^2} = \frac{\pi}{2} - \int_{C_R} \frac{1}{(z^2 + 1)^2} dz.$$

For  $z$  on  $C_R$ , we have  $|z^2 + 1| \geq R^2 - 1$ . Therefore, we have

$$\left| \int_{C_R} \frac{1}{(z^2 + 1)^2} dz \right| \leq \frac{\pi R}{(R^2 - 1)^2}.$$

Consequently, we have as  $R \rightarrow \infty$ ,  $\frac{\pi R}{(R^2 - 1)^2} \rightarrow 0$ . It follows that

$$\int_{-\infty}^{\infty} \frac{1}{(x^2 + 1)^2} dx = \frac{\pi}{2}.$$

As the integrand is an even function, we obtain that

$$\int_0^{\infty} \frac{1}{(x^2 + 1)^2} dx = \frac{\pi}{4}.$$

□

**Question 264.6.**

**Solution.** Consider the function  $f(z) = \frac{z^2}{(z^2 + 9)(z + 4)^2}$  and the simple, closed, and positively oriented contour of a half-circle above the real-axis, centered around the origin with the radius  $R > 3$ . It follows that

$$\int_{-R}^R \frac{x^2}{(x^2 + 9)(x^2 + 4)^2} dx + \int_{C_R} \frac{z^2}{(z^2 + 9)(z + 4)^2} dz = 2\pi(B_1 + B_2).$$

where  $C_R$  denotes the contour of the curve part of the half-circle, and  $B_1$  and  $B_2$  are the residues of the complex integrand at  $z = 3i$  and  $z = 2i$  respectively. First, observe that

$$\begin{aligned} B_1 &= \operatorname{Res}_{z=3i} \frac{z^2}{(z^2+9)(z^2+4)^2} \\ &= \frac{z^2}{(z+3i)(z^2+4)^2} \Big|_{z=3i} = -\frac{3}{50i}. \end{aligned}$$

For computing the  $B_2$  term, observe that

$$\begin{aligned} \frac{z^2}{(z^2+9)(z^2+4)^2} &= \frac{\phi(z)}{(z-2i)^2}, \\ \text{where} \\ \phi(z) &= \frac{z^2}{(z^2+9)(z+2i)^2}. \end{aligned}$$

It follows that

$$B_2 = \phi'(2i) = \frac{13}{200i}.$$

Hence, we have

$$\int_{-R}^R \frac{x^2}{(x^2+9)(x^2+4)^2} dx = \frac{\pi}{100} - \int_{C_R} \frac{z^2}{(z^2+9)(z^2+4)^2} dz.$$

Observe that

$$\left| \int_{C_R} \frac{z^2}{(z^2+9)(z^2+4)^2} dz \right| \leq \frac{R^2}{(R^2-9)(R^2-4)^2} \pi R$$

The RHS of the inequality goes to 0 as  $R \rightarrow \infty$ . With the fact that the integrand is even, we finally obtain

$$\int_0^\infty \frac{x^2}{(x^2+9)(x^2+4)^2} dx = \frac{\pi}{200}.$$

□

#### Question 264.8.

**Solution.** Consider the function  $f(z) = \frac{z}{(z^2+1)(z^2+2z+2)}$ , and the simple, closed, and positively oriented contour of a semi-circle with the radius strictly larger than  $\sqrt{2}$  and the center of the circle at the origin. It follows that the isolated singularities are at  $z = i$  and  $z = -1 + i$ . By the residue theorem, it follows that

$$\int_{-R}^R f(x) dx + \int_{C_R} f(z) dz = 2\pi i(B_0 + B_1),$$

where  $B_0$  and  $B_1$  are the residues at the singularities respectively. It follows that

$$\begin{aligned} B_0 &= \operatorname{Res}_{z=i} f(z) \\ &= \left[ \frac{z}{(z+i)(z^2+2z+2)} \right]_{z=i} = \frac{1}{10} - \frac{1}{5}i \\ B_1 &= \operatorname{Res}_{z=-1+i} f(z) \\ &= \left[ \frac{z}{(z^2+1)(z+1+i)} \right]_{z=-1+i} = -\frac{1}{10} - \frac{3}{10}i. \end{aligned}$$

By the residue theorem, we obtain

$$\int_{-R}^R \frac{x}{(x^2+1)(x^2+2x+2)} dx = -\frac{\pi}{5} - \int_{C_N} \frac{z}{(z^2+1)(z^2+2z+2)} dz.$$

Observe that by the RL-inequality,

$$\left| \int_{C_R} \frac{z}{(z^2 + 1)(z^2 + 2z + 2)} \right| \leq \frac{\pi R^2}{(R^2 - 1)(R - \sqrt{2})^2}.$$

Hence, the integral over the  $C_N$  contour tends to 0, as  $R \rightarrow \infty$ . Therefore, we have

$$\lim_{R \rightarrow \infty} \int_{-R}^R \frac{x}{(x^2 + 1)(x^2 + 2x + 2)} = -\frac{\pi}{5}.$$

□

**Question 264.9.**

**Solution.** Observe that interior to